

EARLY STAGE EXPERIMENT
Science Description

Experiment/Module: Analysis of Intensity Change Processes Experiment (AIPEX)

Investigator(s): Jon Zawislak, Rob Rogers, Jason Dunion, Josh Alland (NCAR), Rosimar Rios-Berrios (NCAR), George Bryan (NCAR), Falko Judt (NCAR), Michael Fischer, Jun Zhang, Paul Reasor, Joe Cione, Trey Alvey, Xiaomin Chen, Ghassan Alaka, Heather Holbach, and Josh Wadler (UM/RSMAS)

Requirements: TD, TS, Category 1

Plain Language Description: Predicting the timing and rate of tropical cyclone (TC) strengthening events remains one of the most challenging aspects of hurricane forecasting. In their early stages, the structure of developing storms is often disorganized such that their circulations are tilted in the vertical, have prominent dry air masses that can be transported into the inner circulation, and lack rainfall coverage all around the center. These are all conditions that would otherwise be considered unfavorable for further strengthening and are often a consequence of the storm experiencing unfavorable winds in its environment. Storms with these characteristics can, however, strengthen and the goal of this experiment is to understand the physical processes and structures that govern whether storms will intensify in this type of environment.

Early Stage Science Objective(s) Addressed:

1. Collect datasets that can be used to improve the understanding of intensity change processes, as well as the initialization and evaluation of 3-D numerical models, particularly for TCs experiencing moderate vertical wind shear [*IFEX Goals 1, 3*].
2. Obtain a quantitative description of the kinematic and thermodynamic structure and evolution of intense convective systems (convective bursts) and the nearby environment to examine their role in TC intensity change [*IFEX Goals 1, 3*].
3. Improve our understanding of the physical processes responsible for the formation and evolution of arc clouds, as well as their impacts on TC structure and intensity in the short-term [*IFEX Goals 1, 3*].
4. Test new (or improved) technologies with the potential to fill gaps, both spatially and temporally, in the existing suite of airborne measurements in early stage TCs. These measurements include improved three-dimensional representation of the TC wind field, more spatially dense thermodynamic sampling of the boundary layer, and more accurate measurements of ocean surface winds [*IFEX Goal 2*].

Motivation: While some improvements in operational tropical cyclone (TC) intensity forecasting have been made in recent years (DeMaria et al. 2014), predicting changes in TC intensity (as defined by the 1-min. maximum sustained wind) remains problematic. In particular, the operational prediction of rapid intensification (RI) has proven to be especially difficult (Kaplan et al. 2010). The significant impact of such episodes has prompted the Tropical Prediction

EARLY STAGE EXPERIMENT

Science Description

Center/National Hurricane Center (TPC/NHC) to declare it as its top forecast priority (Rappaport et al. 2009).

One specific challenge in forecasting TC intensity change is understanding how vertical wind shear (VWS) influences the storm (DeMaria and Kaplan 1994). While the response of storms to low magnitude VWS (favoring strengthening) and high VWS (favoring weakening) are well accepted, storm response in the moderate shear range (i.e., 4.5 to 11 m s⁻¹) can vary greatly. Consider the intensity forecasts for Hurricane Michael (2018), a storm within an environment of moderate VWS. On 7 October, when Michael was a weak tropical depression in the Caribbean Sea, NHC forecasted tropical storm intensity at landfall; two days later, and a day before landfall, NHC forecasted major hurricane intensity. Michael made landfall over the Florida panhandle as a Category 5 hurricane. For under-predicted intensity forecasts like Michael, emergency personnel may have limited time to prepare and respond, and the public may have limited time to evacuate. After the sinking of cargo vessel El Faro event in Hurricane Joaquin (2014), the National Transportation Safety Board specifically mandated that NOAA improve their forecasts of TCs under the influence of moderate VWS as the disaster was attributed to, in part, by uncertain forecasts of Joaquin as it interacted with VWS (Sumwalt et al. 2017). When VWS works together with dry environmental air, TC development is further limited (Tang and Emanuel 2012). However, the pathways by which VWS and dry air work together to limit TC development remain unclear.

VWS is only one of many factors that affect processes that govern TC intensification, which span spatial and temporal scales from the environmental to vortex to convective and smaller scales. Other environmental impacts (e.g., mid-tropospheric relative humidity (RH), boundary layer equivalent potential temperature and static stability, and sea surface temperature (SST) on the TC inner core latent heating structure and distribution are also key to understanding RI. Understanding the communication of those impacts from the TC near-environment (~150–300 km from the TC center) to the inner core (within 150 km), how they vary across time scales, and their impact on inner-core RI processes is critical.

Background: Prior studies have found a number of large-scale environmental factors that are generally favorable for TC intensification, including low environmental vertical wind shear, high ocean heat content, and elevated low- to mid-tropospheric humidity. Thus far, statistically-based prediction schemes that employ predictors derived from large-scale environmental fields and GOES infrared satellite imagery have generally been shown to provide the most skillful objective RI guidance (Kaplan et al. 2015). These schemes include the SHIPS rapid intensification index (SHIPS-RII) (Kaplan et al. 2010) and the more recently developed Bayesian and logistic regression RI models (Rozoff and Kossin 2011). Kaplan et al. (2015) showed that these statistical models are capable of explaining roughly 20% of the skill of Atlantic basin RI forecasts at a lead-time of 24-h. The remaining 80% of the skill not explained by the statistical models is assumed to be attributable either to processes not explicitly accounted for by those models or by limitations in the predictability of RI events.

EARLY STAGE EXPERIMENT

Science Description

On the vortex-scale, a number of observational studies have found that intensifying TCs have more precipitation and convective bursts occurring within the high inertial stability region inside the radius of maximum wind, or RMW (e.g., Rogers et al. 2013, 2015, 2016). This configuration is favorable for TC intensification for two hypothesized reasons: 1) In the high inertial stability region, heat energy is much more efficiently converted to kinetic energy (Schubert and Hack 1982; Vigh and Schubert 2009), and 2) Diabatic heating within the high inertial stability region enables angular momentum surfaces to be drawn inward at the RMW, resulting in tangential wind spin-up (Smith and Montgomery 2016).

Observational studies have also found that intensifying TCs typically have more symmetrically distributed precipitation and deep convection than non-intensifying TCs (e.g., Rogers et al. 2013; Alvey et al. 2015; Tao and Jiang 2015; Wadler et al. 2018a). This is consistent with idealized modeling studies that show that TC intensification is most sensitive to the axisymmetric, azimuthal wavenumber-0 component of diabatic heating (e.g., Nolan et al. 2007). One principal environmental factor that can prevent the development of this symmetry is VWS. The interaction of TCs with environmental VWS typically results in a wavenumber-1 asymmetry in vertical motion and precipitation, in which upward vertical motion and deep convection is favored in the downshear semicircle, while downward motion and suppression of deep convection is observed in the upshear semicircle (e.g., Marks et al. 1992; Reasor et al. 2000, 2013; Rogers et al. 2016; Zawislak et al. 2016). Subsidence in the upshear quadrants can increase the temperature and decrease the relative humidity of the middle troposphere, effectively capping (stabilizing) the lower troposphere. An increase in asymmetry can lead to the decrease in the projection of diabatic heating onto the axisymmetric, azimuthal wavenumber-0 component that has been shown to be important for TC intensification. However, the magnitude of this asymmetry can exhibit considerable variability, particularly within the moderate shear regime (4.5 to 11 m s^{-1}) that has been shown to be problematic for operational intensity forecasts (Bhatia and Nolan 2013).

This suggests the importance of identifying and understanding the environmental and internal (within the inner core) mechanisms that govern the azimuthal and radial distribution of precipitation, as well as its modes (i.e., shallow, moderately deep, and deep convection, as well as stratiform rain), could improve the understanding of the intensification process. Recent studies indicate that these mechanisms may include: the interaction of the vortex with environmental VWS and dry air, ventilation, vortex-scale subsidence, surface enthalpy fluxes from the underlying ocean, tropospheric stability, as well as the accompanying aeriially-averaged vertical mass flux profile. For example, Rogers et al. (2020) identified several important factors that led to the intensification of Hermine (2016): the co-location of deep convection once the mid- and low-level circulations aligned, moistening of the middle troposphere, a stabilization of the lower troposphere, and subsequent lowering of the peak vertical mass flux as more moderately-deep convection became the dominant precipitation mode.

VWS and dry air can work together to limit TC development via ventilation, defined as the flux of low-equivalent potential temperature (θ_E) environmental air into the TC inner core (Simpson and Riehl 1958; Cram et al. 2007; Riemer et al. 2010; Tang and Emanuel 2012; Munsell et al. 2013). Dry air can ventilate the subcloud layer via convective downdrafts (downdraft ventilation)

EARLY STAGE EXPERIMENT

Science Description

(Riemer et al. 2010). Dry air can also ventilate rising air in the eyewall via the inward radial transport (radial ventilation). Didlake and Houze (2013) documented a potentially separate ventilation pathway in Hurricane Rita (2005), which had strong descending radial inflow associated with a stratiform rainband. This inflow was associated with low- θ_E air at middle and low levels, and may transport dry air into convection (e.g., radial ventilation) or downward into the subcloud layer (e.g., downdraft ventilation). Under certain oceanic conditions, the air-sea enthalpy fluxes can lead to recovery of low- θ_E air in the boundary layer (Molinari et al. 2013; Zhang et al. 2013, 2017; Wadler et al. 2018b; Nguyen et al. 2019). In particular, Nguyen et al. (2019) showed that initially weak TCs that underwent intensification tend to have larger enthalpy fluxes in the upshear quadrants than weakening TCs.

Overall, ventilation pathways can modulate the spatial structure of precipitation, especially upshear, which has been shown to be associated with TC intensity (e.g., Rogers et al. 2013; Alvey et al. 2015; Tao and Jiang 2015; Rios-Berrios 2016a, b; Nguyen et al. 2017; Fischer et al. 2018, Rios-Berrios et al. 2018). The majority of studies investigating the effects of ventilation on TC development has focused on mature TCs, but the effects for weak tropical cyclones have not received much attention. Recent work has used idealized modeling to document the effects of ventilation pathways on TC development for weak TCs (Alland et al. 2020a, b). These experiments showed the modulating effects on TC development to be:

1. Downdraft ventilation, which transported low- θ_E air left-of-shear and upshear between heights of 0 and 3 km associated with the principal rainband.
2. Radial ventilation, which transported low- θ_E air left-of-shear and upshear between heights of 0 and 3 km associated with the principal rainband (coupled with downdraft ventilation), and upshear and right-of-shear between heights of 5 and 9 km associated with a vertically-tilted TC vortex.

Both of these ventilation pathways aided in reducing the area of strong upward motions ($w > 0.5 \text{ m s}^{-1}$), reducing the vertical mass flux in the inner core, and limiting TC development. However, these experiments were simplified (e.g., linear wind profile with height, constant SST, dry air surrounding the TC). How do these ventilation pathways modulate TC development for a real storm? Observations are necessary to document the importance of these ventilation pathways, and their spatial structures, on TC development so the atmospheric science community can verify models and better predict TC intensity change.

Goal(s): Collect aircraft observations that will allow us to characterize the precipitation and vortex-scale kinematic and thermodynamic structures of TCs experiencing moderate vertical shear. Understanding the reasons behind these structures, particularly greater azimuthal coverage of precipitation, vortex alignment, and boundary layer ventilation and recovery, will contribute toward a greater understanding of the physical processes that govern whether TCs will intensify, (especially those that undergo RI) in this type of environment.

Hypotheses: The following hypotheses are guided by the theory that the thermodynamic and kinematic variability in the region of the TC near-environment (i.e., ~150–300 km from the TC

EARLY STAGE EXPERIMENT
Science Description

center) are communicated to the inner-core (<150 km from the center), which subsequently impacts the precipitation (i.e., latent heating) distribution in a way that may be favorable for TC (rapid) intensification.

Inner core Processes

1. The local kinematic (e.g., shear, vertical alignment of the vortex, inertial stability), thermodynamic (e.g., SST, static stability, and moisture/RH), and boundary layer properties (e.g., strength and depth of radial inflow, SST, surface enthalpy flux) are key in governing the precipitation modes and whether that precipitation is symmetrically distributed and primarily inside the RMW, which is a favorable configuration for intensification.
 - a. The middle troposphere is moistened upshear due to detrainment from mid-tropospheric congestus, evaporation of falling stratiform rain, or reduced lateral advection of dry air from the environment
 - b. The lower troposphere is convectively unstable in the upshear quadrants due to enhanced surface enthalpy fluxes from the underlying ocean and/or reduced convective downdrafts
2. The presence of dry air and VWS alone does not guarantee that intensification will be limited. Downdraft and/or radial ventilation must occur for the combined effects of dry air and VWS to limit TC development.
 - a. Downdraft ventilation influences TC development by transporting low- θ_E air downward left-of-shear and upshear (cyclonically downwind of convection). This low- θ_E air modulates convection left-of-shear and upshear.
 - b. Radial ventilation influences TC development by modulating convection in two ways:
 - i. At low levels, radial ventilation is coupled with downdraft ventilation and is associated with rainband activity.
 - ii. At middle and upper levels, radial ventilation, due to storm-relative flow associated with a tilted vortex and/or a weak vortex circulation, transports low- θ_E air upshear and right-of-shear. Convection occurring downshear is relatively protected from this dry air, unless the VWS magnitude is sufficiently strong.

Near-environment Processes:

1. Boundary layer inflow strength and depth in the near-environment varies diurnally with maxima (minima) in the early morning (afternoon) that represents a relatively favorable (unfavorable) period for intensification.
2. TC diurnal pulses exhibit squall-line-like behaviors as they propagate away from the TC each day and can initiate strong convectively-driven downdrafts in the near-environment, particularly in dry air positioned upshear, that temporarily disrupt the boundary layer via drying and enhanced static stability.

EARLY STAGE EXPERIMENT
Science Description

Objectives:

1. Quantify inner core and near-environment thermodynamic and kinematic variability, specifically as they relate to the precipitation modes and distributions observed within the inner core during intensity change periods
2. Observe diurnal variability in the TC planetary boundary layer, in particular changes to the radial inflow strength, and its relationship with the timing of intensity change
3. Use observations within the inner core and near-environment to assess whether hypothesized downdraft and radial ventilation is occurring, and, if so, which specific pathways are affecting the inner convective structure and distribution; such as from the downward transport of low- θ_e air (downdraft ventilation), inward transport of low- θ_e air (radial ventilation), and the subsequent PBL recovery from surface enthalpy fluxes.

Overall, these objectives convey the need for a systematic, three-dimensional examination of physical processes near and within the TC inner core that act to modulate the spatial structure of convection, the thermodynamic and kinematic structures, and potentially, the intensity of a TC. Observations can also be compared with model and satellite-derived quantities (e.g., wind, moisture) to improve satellite algorithms and evaluate model representations (e.g., parameterizations) (IFEX Goal #1). TCs that are experiencing moderate vertical wind shear (4.5 to 11 m s⁻¹) over a deep layer (850–200 hPa) are of particular interest, since this range of shear values is often associated with considerable uncertainty with respect to the prospect for TC intensification (Bhatia and Nolan 2013).

Aircraft Pattern/Module Descriptions (see *Flight Pattern* document for more detailed information):

Missions will be targeted for systems that have a reasonable chance of undergoing intensification based on statistical and numerical model forecast guidance. When possible (i.e., subject to range, timing, and other logistical constraints), missions will begin at least 24 h prior to the expected onset of intensification, while the TC is still at tropical depression or tropical storm intensity. This enables the documentation of TC structure during the time leading up to intensification onset (if it indeed occurs). Ideally missions will continue every 12 to 24 h, (depending on the *Scenario* below) as long as feasible. Although all intensification rates are of interest, ***priority will be given to those with a high potential for RI*** according to model guidance and/or are forecast to experience at least moderate (4.5 to 11 m s⁻¹) vertical wind shear over a deep layer. There are a few possible configurations for the execution of this experiment:

P-3 Pattern #1: A standard Figure-4 pattern centered on the estimated low-level center, oriented such that the radial passes are aligned through approximately the upshear, downshear, left-of-shear, and right-of-shear directions. This pattern can be flown oriented within the quadrants; i.e., 45° [downshear right], 315° [downshear left], 225° [upshear left], 135° [upshear right]) and rotated to complete a full rotated Figure-4 or flown with repeated azimuths to complete a repeated Figure-4.

EARLY STAGE EXPERIMENT

Science Description

P-3 Pattern #2: A standard Butterfly pattern centered on the estimated low-level center. The butterfly should be oriented such that the upshear semicircle (or downshear if precipitation sampling is preferred in an asymmetric precipitation configuration) contains the most radial legs.

P-3 Module #1 (“Upshear Circumnavigation Module”): Sample the upshear semicircle, including the boundary between no convection and convection, if such a boundary exists, at up to three possible radii: 90 n mi (167 km), 60 n mi (111 km), and 40 n mi (74 km). Release up to 8 dropsondes per radius, as equally spaced as possible. The P-3 should operate as high as possible. This high-altitude circumnavigation allows for increased azimuthal and vertical dropsonde data coverage, particularly in the critical, precipitation-free upshear region that may fill in as intensification commences.

P-3 Module #2 (“Rainband Module”): Follow the principal rainband inward towards the center by paralleling the band (either radially inside or outside), keeping the band within range [~10 n mi (19 km)] of the TDR, and releasing dropsondes approximately every 20 n mi (37 km).

P-3 Module #3 (“High Density Eyewall Drops”): A sequence of up to 8 dropsondes, released every 2.5-5 n mi (5–10 km, approximately every 30 seconds to 1 minute), distributed such that it covers inbound/outbound of the RMW and convection in the developing eyewall region.

G-IV Pattern #1: A standard octagonal circumnavigation pattern centered on the estimated low-level center. Radius of each circumnavigation is: 150 n mi (277 km), 90 n mi (167 km), 60 n mi (111 km).

G-IV Pattern #2: A standard Star with Circumnavigation pattern. If desired, the outer points can be oriented to align with the VWS vector. Supplemental observations can also be made when model sensitivity regions are indicated (e.g., derived from ECMWF and the COAMPS-TC model ensembles) that could positively impact forecasts of TC track, intensity and/or structure.

G-IV Module #1: A Figure-4 pattern centered on the low-level center, with radial legs extending up to 105 n mi (195 km). If desired, the pattern can be centered on the estimated low-level center, oriented such that the radial passes are aligned through approximately the upshear, downshear, left-of-shear, and right-of-shear directions.

Scenario #1: Both P-3s (Back-to-Back) and G-IV

One P-3 will fly Pattern #1 or Pattern #2 with the option to either follow the initial Figure-4 / butterfly with Module #1 (“Upshear Circumnavigation Module”, depending on the extent of the precipitation in the upshear semicircle that may pose limitations on flight altitudes above the freezing level), Module #2 (“Rainband Module”), or rotate and complete a second Figure-4, or fly the same oriented Figure-4 as the initial. Arrival of the second P-3 should be timed to arrive when the first P-3 mission departs the storm environment. The second P-3 mission should complete the same pattern as the first. If available, a sUAS can be released during either or both P-3 missions to complete an “Inflow Module” outlined in RICO SUAVE, ideally being flown in the PBL below

EARLY STAGE EXPERIMENT

Science Description

1000 ft (305 m), and released left-of-shear, flown cyclonically to the downshear right quadrant, and subsequently penetrating the eyewall into the eye.

In this scenario, the G-IV will observe the outer to near-TC environment [~60 n mi (~100 km) and outward to ~160 n mi (~300 km) from the center] using the circumnavigation in Pattern #1 or Pattern #2 (Star with Circumnavigation, which should be oriented with two outer points being upshear), with an option to complete Module #1, if no P-3 has achieved higher-altitude dropsonde releases near the TC center. The flight should be timed such that the mission overlaps some of each P-3 mission.

Scenario #2: One P-3 and the G-IV, coordinated

One P-3 will fly Pattern #1 (Figure-4) or Pattern #2 (Butterfly) with the option to either follow the initial Figure-4 / butterfly with Module #1 (“Upshear Circumnavigation Module”, depending on the extent of the precipitation in the upshear semicircle that may pose limitations on flight altitudes above the freezing level), Module #2 (“Rainband Module”), or rotate and complete a second Figure-4, or fly the same oriented Figure-4.. If available, a sUAS can be released during either or both P-3 missions to complete an “Inflow Module” outlined in RICO SUAVE, ideally being flown in the PBL below 1000 ft (305 m), and released left-of-shear, flown cyclonically to the downshear right quadrant, and subsequently penetrating the eyewall into the eye.

In this scenario, the G-IV will observe the outer to near-TC environment [~60 n mi (100 km) and outward to ~160 n mi (~300 km) from the center] using the circumnavigation in Pattern #1 or Pattern #2 (Star with Circumnavigation, which should be oriented with two outer points being upshear), with an option to complete Module #1, if no P-3 has achieved higher-altitude dropsonde releases near the TC center. This scenario leaves the possibility open to 12-hourly (twice-a-day) coordinated missions.

Scenario #3: Only one P-3 available (no G-IV)

When the G-IV and 2nd P-3 is not available for coordinated operations, either because of operational tasking requirements or aircraft unavailability, P-3 targeted observations in the near environment and inner core can still contribute towards the objectives of the experiment. In this scenario there are two possible strategies for sampling, which depend on whether the precipitation distribution is asymmetric:

a. TC is highly asymmetric:

This option will be chosen when the precipitation distribution in the targeted TC is expected to be highly asymmetric during the mission. Such an asymmetric configuration would allow for a high-altitude P-3 circumnavigation pattern (P-3 Module #1) to at least target the precipitation-free upshear semicircle, and when hazard avoidance is possible, to extend the circumnavigation to the downshear quadrants.

Indications of an appropriate magnitude of asymmetry may include:

EARLY STAGE EXPERIMENT
Science Description

- 1) Visible, infrared, or microwave satellite imagery indicates an exposed or partially-exposed low-level circulation center (see example below).
- 2) The environmental vertical wind shear, as indicated by SHIPS, is expected to be sufficient ($> 4.5 \text{ m s}^{-1}$) during the mission to result in an asymmetric precipitation structure.
- 3) High-resolution numerical guidance (i.e. HWRF) forecast a lack of precipitation in the upshear semicircle of the TC during the mission.




Figure-4 in green, circumnavigation in orange, shear vector in black, 'X' is a dropsonde location).

Given this scenario, the P-3 will sample the near environment and inner core with a pattern that includes a high-altitude circumnavigation and, optimally, a rotated Figure-4 in P-3 Pattern #1 or butterfly in P-3 Pattern #2. If time doesn't permit for a complete rotated Figure-4, then a single Figure-4 can be substituted.

b. TC is relatively symmetric:

This scenario applies to a targeted TC that has the potential for intensification, but the precipitation is expected to be too symmetric during the mission for the P-3 high-altitude circumnavigation to be conducted safely. Here, the P-3 will sample the inner-core vortex structure with a standard rotated Figure-4 (P-3 Pattern #1) or Butterfly (P-3 Pattern #2) pattern.

Scenario #4: Only G-IV available (no P-3s)

This option is less preferable as targeted observations of the vortex structure are also important towards the objectives of the experiment. This option applies to any targeted TC that has the potential for intensification, regardless of asymmetric structure. Under this option, the G-IV will sample the outer and near environments with either the circumnavigation (G-IV Pattern #1) or star with circumnavigation (G-IV Pattern #2), and requires that hazard avoidance permit the G-IV to obtain measurements within/very near the inner core, possibly (if time permits), a single Figure-4 center on the low-level center (G-IV Module #1).

EARLY STAGE EXPERIMENT

Science Description

Links to Other Early Stage Experiments/Modules:

If the opportunity arises during the execution of AIPEX, fly the Convective Burst Module or Arc Cloud Module (see accompanying discussion in Field Program plan). The Convective Burst Module would be optimal for determining the structure and evolution of deep convection within the framework of the broader vortex-scale circulation as it interacts with vertical shear (if appropriate), while the Arc Cloud Module would be ideal for documenting locations within the vortex circulation encountering significant low- to mid-level dry air and determining the impact of the associated outflow boundaries on the boundary layer temperature and moisture distribution (an example of the communication that occurs between the inner core and near environment). AIPEX can also be flown in conjunction with the following *Early Stage* experiments and modules: SFMR Module, TDR Experiment, Synoptic Flow Experiment, NESDIS JPSS Satellite Validation Experiment, and ADM-Aeolus Satellite Validation Module, or with RICO SUAVE in the *Mature Stage*.

Link to the ONR TCRI Program:

TCRI Investigator(s): James Doyle (NRL-Monterey), Jon Moskaitis (NRL-Monterey), Michael Bell (CSU), George Bryan (NCAR), Joseph Cione (NOAA/AOML/HRD), Jason Dunion (Univ. of Miami/CIMAS-NOAA/AOML/HRD), Russell Elsberry (UCCS/Naval Postgraduate School), Joel Feldmeier (Naval Postgraduate School), Ralph Foster (Univ. of Washington), Zeljka Fuchs (New Mexico Tech), Chanh Kieu (Indiana University), Sharan Majumdar (Univ. of Miami/RSMAS), David Nolan (Univ. of Miami/RSMAS), David Raymond (New Mexico Tech), David Richter (Notre Dame), Robert Rogers (NOAA/AOML/HRD), C.G. Rossby (Univ. Chicago), Tony Wimmers (Univ. of Wisconsin/CIMSS), Christopher Velden (Univ. of Wisconsin/CIMSS), Jonathan Zawislak (Univ. of Miami/CIMAS-NOAA/AOML/HRD), and Jun Zhang (Univ. of Miami/CIMAS-NOAA/AOML/HRD)

In addition, beginning in 2020, IFEX will collaborate with the Office of Naval Research (ONR) Departmental Research Initiative (DRI) Tropical Cyclone Rapid Intensification (TCRI) Program. As part of this program, TCRI will utilize the P-3s and possibly the G-IV in coordinated research missions with IFEX. The overarching goal of TCRI is to utilize multiple tools (observations, basic theory, and numerical modeling experiments) to better understand, and subsequently predict, TC intensity change — with an emphasis on rapid intensification events.

AIPEX and the TCRI program share many common scientific objectives, and thus the observing strategies outlined here for AIPEX have been designed in such a way that integrates the science questions and observing strategies of the TCRI Science Team for P-3 and G-IV airborne data collection. The close collaboration with TCRI, and their involvement in the execution of AIPEX and G-IV Synoptic Flow missions, will be equally beneficial towards the success of both IFEX and TCRI. Listed below are some specific science questions outlined by the TCRI Science Team:

Identify the key multi-scale processes and dynamics governing the onset of RI and how this influences RI predictability. Specifically:

EARLY STAGE EXPERIMENT

Science Description

- How does TC intensification evolve from the initial three-dimensional asymmetric structure of a TC? What key processes governing eyewall formation?
- How do air-sea interaction and planetary boundary layer structure and dynamics modulate TC intensification and RI, such as PBL height, radial inflow/convergence, agradient force, air-sea disequilibrium, and recovery? How does the radial structure of the tangential wind in the boundary layer evolve before and during the onset of intensification? Where is the radial location of maximum frictional convergence relative to the RMW? How do the surface fluxes of heat and moisture influence the onset of RI, particularly where the OHC varies substantially?
- Do properties of the TC boundary layer (including the azimuthal asymmetry of these properties) provide valuable information about TC intensification that precedes other observable properties such as convective bursts and latent heating?
- What are the most favorable spatial and temporal arrangements of diabatic heating and cooling for rapid intensification or weakening? Does the location of the latent heating relative to the RMW within the mid- and upper-troposphere matter for intensification onset? How do the microphysical processes govern the horizontal and vertical latent heating distribution? Is the moist neutrality condition assumed in the current WISHE theory for TC development strictly valid during RI?
- Do diabatic or boundary layer processes dominate during the onset of intensification? Why does the eyewall form? How are the organizational changes before and during RI distinct from cases of more gradual intensification?
- How do thermodynamic conditions in a cyclone affect the eyewall mass flux?
- What is the impact of VWS on RI immediately following TC formation and RI in already mature TCs?
- What are the multiscale (environmental, vortex, PBL, convective, microphysical) factors that govern the radial, azimuthal, vertical, and temporal distribution of TC inner-core latent heating? How are these factors modulated by the near environment [~ 80 - 160 n mi (~ 150 - 300 km)] and how is this modulation communicated from the near environment to the inner core?
- To what extent are RI predictions limited by uncertainties in the near-storm environment and the vortex? Do additional observations in sensitive regions (e.g., derived from ECMWF and the COAMPS-TC model ensembles) improve forecasts of the onset of intensification or RI?

In addition, the TCRI Science Team seeks to use a combination of aircraft and satellite observations to analyze the 3-dimensional TC wind field and answer the following science questions:

- How much of the TC near-surface wind field evolution can be captured in multispectral satellite image information (e.g., GMI, AMSR2, SSMIS, ATMS)?
- Can satellite-borne Synthetic Aperture Radar (SAR) be used to improve surface wind vector retrievals? Can they be used to estimate physically-realistic vertical profiles of the mean wind profiles in the boundary layer for the full sampling of the storm?

EARLY STAGE EXPERIMENT

Science Description

Analysis Strategy: The general analysis strategy follows that performed in recent observational and modeling studies (e.g., Reasor et al. 2013; Zhang et al. 2013; Rogers et al. 2016; Zawislak et al. 2016; Nguyen et al. 2017; Rios-Berrios et al. 2018, Nguyen et al. 2019, Rogers et al. 2020, Alland et al. 2020a, b). The analysis strategy includes assessing and documenting the time evolution of the following:

- *Azimuthal and radial distribution of inner-core precipitation and deep convection* (P-3 TDR/LF, possibly G-IV TDR). The inner-core precipitation asymmetry, and its projection onto the axisymmetric, azimuthal wavenumber-0 component will be assessed quantitatively (assuming sufficient azimuthal coverage). The location of precipitation and convective bursts relative to the RMW will be examined.
- *Precipitation mode, particularly upshear* (P-3 TDR/LF, possibly G-IV TDR). An analysis of the precipitation mode (shallow, moderately deep, deep convection, as well as stratiform rain), using the vertical velocity and reflectivity structure, will allow for an assessment as to whether moistening of the inner core occurs through upscale growth of convection (moistening from convective detrainment at gradually higher altitudes), or from the top-down via stratiform rain as hydrometeors produced downshear are transported azimuthally upshear.
- *Characteristics of vertical mass flux profiles* (P-3 TDR, possibly G-IV TDR). Literature (Raymond et al. 2011; Gjorgjievska and Raymond 2014; Rogers et al. 2020) has identified a key transition from a “top heavy” vertical mass flux profile (convergence preferentially in the middle troposphere) to a “bottom heavy” profile (convergence preferentially in the lower troposphere) in the development of a TC. To further understand this evolution, the relationship between the mass flux profiles and the contributions of various precipitation modes (through vertical velocity, reflectivity) to overall precipitation will be analyzed. In addition, measurements within 2–3x the RMW using TDR analyses will be used to determine if there is a relationship between the vertical mass flux, ventilation pathways, and TC intensity.
- *Low-wavenumber thermodynamic and kinematic structure of the boundary layer* (P-3/G-IV dropsondes, DWL for kinematic only), including surface fluxes of heat, moisture, and momentum. The thermodynamic focus will be on the boundary layer cooling by convective downdrafts and the subsequent recovery via surface enthalpy fluxes from the underlying ocean in the downstream (upshear-left through downshear-right) quadrants. Surface enthalpy fluxes will be calculated where dropsondes are paired with AXBTs that provide SST, and/or from sUAS and IRsonde measurements. The kinematic focus will be on obtaining measurements of the strength and depth of boundary layer inflow and convergence in the boundary layer, both in a symmetric sense and relative to the shear vector (when relevant). Additionally, the gradient and agradient flow in the boundary layer will be calculated.
- *Low-wavenumber thermodynamic and kinematic structure above the boundary layer* (P-3/G-IV dropsondes, P-3/G-IV TDR and DWL for kinematic only). The presence of mid-tropospheric dry air is of particular interest. Assuming mid-tropospheric dry air is present (most likely in the upshear quadrants), the potential sources of this dry air (vortex-scale subsidence or lateral advection from the environment) and how this upshear dry air is

EARLY STAGE EXPERIMENT

Science Description

removed (i.e., through detrainment from congestus or evaporation of stratiform precipitation) will be assessed.

- *Vortex tilt* (P-3 TDR, possibly G-IV TDR). Assuming sufficient TDR coverage, the vortex tilt will be examined quantitatively by merging TDR analyses from each Figure-4. If the vortex tilt appears to decrease rapidly during a flight, individual TDR analyses can be used to qualitatively examine the time evolution of vortex tilt during the alignment process.
- *Vertical wind shear and upper-level divergence* (G-IV dropsondes). These quantities will be computed and compared with global model analyses. The vertical distribution of shear will also be evaluated (i.e., VWS calculations in different vertical levels outside of the standard 850 to 200 hPa will be calculated, c.f. Finocchio et al. 2016, 2017), as upper-level shear is hypothesized to be less deleterious than low-level shear.
- *Ventilation* will be quantified by combining the thermodynamic data from dropsondes with wind analyses from dropsondes and the TDR. Radial and downdraft ventilation will both be calculated. It will be important to obtain deep vertical profiles of thermodynamic and kinematic data in the upshear semicircle to determine the vertical extent of ventilation pathways. The necessity for thermodynamic and kinematic coverage upshear is why it is recommended to release more dropsondes than the standard flight patterns/modules.

The overarching hypothesis is that, by performing the above analyses for multiple AIPEX data sets collected during both RI and non-RI events, it will be possible to determine the conditions that are triggers for RI. This analysis strategy can also assist in the evaluation of 3-D numerical models, including the sufficiency (or lack thereof) of the horizontal resolution, and the microphysical and planetary boundary layer parameterization schemes. Dropsonde, flight-level, super-obbed Doppler radar, and SFMR data are made available over the GTS and assimilated in real time, while full Doppler fields and lower fuselage radar will be available post-flight.

References:

- Alland, J. J., B. H. Tang, K. L. Corbosiero, and G. H. Bryan, 2020a: Combined effects of midlevel dry air and vertical wind shear on tropical cyclone development. Part I: Downdraft ventilation. *in review*.
- Alland, J. J., B. H. Tang, K. L. Corbosiero, and G. H. Bryan, 2020b: Combined effects of midlevel dry air and vertical wind shear on tropical cyclone development. Part I: Radial ventilation. *in review*.
- Alvey III, G. R., J. Zawislak, and E. Zipser, 2015: Precipitation Properties Observed during Tropical Cyclone Intensity Change. *Mon. Wea. Rev.*, **143**, 4476–4492. doi: 10.1175/MWR-D-15-0065.1.
- Bhatia, K. T., and D. S. Nolan, 2013: Relating the skill of tropical cyclone intensity forecasts to the synoptic environment. *Wea. Forecasting*, **28**, 961–980, doi: 10.1175/WAF-D-12-00110.1.
- Cram, T. A., J. Persing, M. T. Montgomery, and S. A. Braun, 2007: A Lagrangian trajectory view on transport and mixing processes between the eye, eyewall, and environment using a high-resolution simulation of Hurricane Bonnie (1998). *J. Atmos. Sci.*, **64**, 1835–1856, doi:10.1175/JAS3921.1.

EARLY STAGE EXPERIMENT

Science Description

- DeMaria, M., and J. Kaplan, 1994: A statistical hurricane intensity prediction scheme (SHIPS) for the Atlantic basin. *Wea. Forecasting*, **9**, 209–220, doi:10.1175/1520-0434(1994)0092.0.CO;2.
- DeMaria, M., M. Mainelli, L. K. Shay, J. A. Knaff, and J. Kaplan, 2005: Further improvements to the statistical hurricane intensity prediction scheme (SHIPS). *Wea. Forecasting*, **20**, 531–543, doi:10.1175/WAF862.1.
- DeMaria, M., C. R. Sampson, J. A. Knaff, and K. D. Musgrave, 2014: Is tropical cyclone intensity guidance improving? *Bull. Amer. Meteor. Soc.*, **95**, 387–398, doi:10.1175/BAMS-D-12-00240.1.
- Didlake, A. C., and R. A. Houze, 2013: Dynamics of the stratiform sector of a tropical cyclone rainband. *J. Atmos. Sci.*, **70**, 1891–1911, doi:10.1175/JAS-D-12-0245.1.
- Finocchio, P. M., S. J. Majumdar, D. S. Nolan, and M. Iskandarani, 2016: Idealized tropical cyclone responses to the height and depth of environmental vertical wind shear. *Mon. Wea. Rev.*, **144**, 2155–2175, doi:10.1175/MWR-D-15-0320.1.
- Finocchio, P. M., and S. J. Majumdar, 2017: A statistical perspective on wind profiles and vertical wind shear in tropical cyclone environments of the Northern Hemisphere. *Mon. Wea. Rev.*, **145**, 361–378, doi:10.1175/MWR-D-16-0221.1.
- Gjorgjievska, S., and D. J. Raymond, 2014: Interaction between dynamics and thermodynamics during tropical cyclogenesis. *Atmos. Chem. and Phys.*, **14**, 3065–3082, doi:10.5194/784acp-14-3065-2014.
- Kaplan, J., M. DeMaria, and J. A. Knaff, 2010: A revised tropical cyclone rapid intensification index for the Atlantic and eastern North Pacific basins. *Wea. Forecasting*, **25**, 220–241, doi: 10.1175/2009WAF2222280.1.
- Kaplan, J., and Coauthors, 2015: Evaluating environmental impacts on tropical cyclone rapid intensification predictability utilizing statistical models. *Wea. Forecasting*, **30**, 1374–1396, doi: 10.1175/WAF-D-15-0032.1.
- Marks, F. D., Jr., R. A. Houze Jr., and J. F. Gamache, 1992: Dual-aircraft investigation of the inner core of Hurricane Norbert. Part I: Kinematic structure. *J. Atmos. Sci.*, **49**, 919–942, doi: 10.1175/1520-0469(1992)049<0919:DAIOTI>2.0.CO;2.
- Molinari, J., J. Frank, and D. Vollaro, 2013: Convective Bursts, Downdraft Cooling, and Boundary Layer Recovery in a Sheared Tropical Storm. *Mon. Wea. Rev.*, **141**, 1048–1060, doi: 10.1175/MWR-D-12-00135.1.
- Munsell, E. B., F. Zhang, and D. P. Stern, 2013: Predictability and dynamics of a nonintensifying tropical storm: Erika (2009). *J. Atmos. Sci.*, **70**, 2505–2524, doi:10.1175/JAS-D-12-0243.1.
- Nguyen, L. T., R. Rogers, and P. Reasor 2017: Thermodynamic and kinematic influences on precipitation symmetry in sheared tropical cyclones: Bertha and Cristobal (2014). *Mon. Wea. Rev.*, **145**, 4423–4446, <https://doi.org/10.1175/MWR-D-17-0073.1>
- Nguyen, L.T., R. Rogers, J. Zawislak, and J.A. Zhang, 2019: Assessing the Influence of Convective Downdrafts and Surface Enthalpy Fluxes on Tropical Cyclone Intensity Change in Moderate Vertical Wind Shear. *Mon. Wea. Rev.*, **147**, 3519–3534, <https://doi.org/10.1175/MWR-D-18-0461.1>

EARLY STAGE EXPERIMENT

Science Description

- Nolan, D. S., Y. Moon, and D. P. Stern, 2007: Tropical cyclone intensification from asymmetric convection: Energetics and efficiency. *J. Atmos. Sci.*, **64**, 3377–3405, doi: 10.1175/JAS3988.1.
- Rappaport, E. N., and Coauthors, 2009: Advances and Challenges at the National Hurricane Center. *Wea. Forecasting*, **24**, 395–419.
- Raymond, D. J., S. L. Sessions, and C. L. Carrillo, 2011: Thermodynamics of tropical cyclogenesis in the northwest pacific. *J. Geophys. Res. Atmos.*, **116** (D18), doi:10.1029/2011JD015624
- Reasor, P. D., M. T. Montgomery, F. D. Marks, and J. F. Gamache, 2000: Low-wavenumber structure and evolution of the hurricane inner core observed by airborne dual-Doppler radar. *Mon. Wea. Rev.*, **128**, 1653–1680, doi:10.1175/1520-0493(2000)1282.0.CO;2.
- Reasor, P. D., R. F. Rogers, and S. Lorsolo, 2013: Environmental flow impacts on tropical cyclone structure diagnosed from airborne Doppler radar composites. *Mon. Wea. Rev.*, **141**, 2949–2969, doi: 10.1175/MWR-D-12-00334.1.
- Riemer, M., M. T. Montgomery, and M. E. Nicholls, 2010: A new paradigm for intensity modification of tropical cyclones: Thermodynamic impact of vertical wind shear on the inflow layer. *Atmos. Chem. Phys.*, **10**, 3163–3188, doi:10.5194/acp-10-3163-2010.
- Rios-Berrios, R., R. D. Torn, and C. A. Davis, 2016a: An ensemble approach to investigate tropical cyclone intensification in sheared environments. Part I: Katia (2011). *J. Atmos. Sci.*, **73**, 71–93, doi:10.1175/jas-d-15-0052.1.
- Rios-Berrios, R., R. D. Torn, and C. A. Davis, 2016b: An ensemble approach to investigate tropical cyclone intensification in sheared environments. Part II: Ophelia (2011). *J. Atmos. Sci.*, **73**, 1555–1575, doi:10.1175/jas-d-15-0245.1.
- Rios-Berrios, R., C. A. Davis, and R. D. Torn, 2018: A hypothesis for the intensification of tropical cyclones under moderate vertical wind shear. *J. Atmos. Sci.*, **75**, 4149–4173, doi:10.1175/jas-d-18-0070.1.
- Simpson, R., and R. Riehl, 1958: Mid-tropospheric ventilation as a constraint on hurricane development and maintenance. Tech. Conf. on Hurricanes, Amer. Meteor. Soc., Miami Beach, FL, D4–1–D4–10.
- Smith, R.K. and Montgomery, M.T. (2016), The efficiency of diabatic heating and tropical cyclone intensification. *Q.J.R. Meteorol. Soc.*, **142**, 2081–2086, doi:10.1002/qj.2804
- Sumwalt, R. L., C. A. Hart, E. F. Weener, and T. B. Dinh-Zahr, 2017: Tropical cyclone information for mariners. National Transportation Safety Board safety recommendation report, 21 pp, <https://www.nts.gov/investigations/AccidentReports/Reports/MSR1702.pdf>.
- Rogers, R., P. Reasor, and S. Lorsolo, 2013: Airborne Doppler observations of the inner-core structural differences between intensifying and steady-state tropical cyclones. *Mon. Wea. Rev.*, **141**, 2970–2991, doi: 10.1175/MWR-D-12-00357.1.
- Rogers, R. F., P. D. Reasor, and J. A. Zhang, 2015: Multiscale structure and evolution of Hurricane Earl (2010) during rapid intensification. *Mon. Wea. Rev.*, **143**, 536–562, doi: 10.1175/MWR-D-14-00175.1.
- Rogers, R., J. Zhang, J. Zawislak, H. Jiang, G. Alvey, E. Zipser, and S. Stevenson, 2016: Observations of the structure and evolution of Hurricane Edouard (2014) during intensity change. Part II: Kinematic structure and the distribution of deep convection. *Mon. Wea. Rev.*, **144**, 3355–3376, doi: 10.1175/MWR-D-16-0017.1.

EARLY STAGE EXPERIMENT

Science Description

- Rogers, R. F., P. D. Reasor, J. A. Zawislak, and L. T. Nguyen, 2020: Precipitation Processes and Vortex Alignment during the Intensification of a Weak Tropical Cyclone in Moderate Vertical Shear. *Mon. Wea. Rev.*, **148**, 1899–1929, <https://doi.org/10.1175/MWR-D-19-0315.1>
- Rozoff, C. M., and J. P. Kossin, 2011: New probabilistic forecast models for the prediction of tropical cyclone rapid intensification. *Wea. Forecasting*, **26**, 677–689, doi: 10.1175/WAF-D-10-05059.1.
- Schubert, W. H., and J. J. Hack, 1982: Inertial stability and tropical cyclone development. *J. Atmos. Sci.*, **39**, 1687–1697, doi:10.1175/1520-0469(1982)039<1687:ISATCD>2.0.CO;2.
- Tang, B., and K. Emanuel, 2012: Sensitivity of tropical cyclone intensity to ventilation in an axisymmetric model. *J. Atmos. Sci.*, **69**, 2394–2413, doi:10.1175/jas-d-11-0232.1.
- Tao, C., and H. Jiang, 2015: Distributions of shallow to very deep precipitation–convection in rapidly intensifying tropical cyclones. *J. Climate*, **28**, 8791–8824, doi: 10.1175/JCLI-D-14-00448.1.
- Vigh, J. L., and W. H. Schubert, 2009: Rapid development of the tropical cyclone warm core. *J. Atmos. Sci.*, **66**, 3335–3350, doi: 10.1175/2009JAS3092.1.
- Wadler, J.B., R.F. Rogers, and P.D. Reasor, 2018a: The Relationship Between Spatial Variations in the Structure of Convective Bursts and Tropical Cyclone Intensification as Determined by Airborne Doppler Radar. *Mon. Wea. Rev.*, **146**, 761–780, <https://doi.org/10.1175/MWR-D-17-0213.1>
- Wadler, J.B., J.A. Zhang, B. Jaimes, and L.K. Shay, 2018b: Downdrafts and the Evolution of Boundary Layer Thermodynamics in Hurricane Earl (2010) before and during Rapid Intensification. *Mon. Wea. Rev.*, **146**, 3545–3565, <https://doi.org/10.1175/MWR-D-18-0090.1>
- Zawislak, J., H. Jiang, G. Alvey, E. Zipser, R. Rogers, J. Zhang, and S. Stevenson, 2016: Observations of the structure and evolution of Hurricane Edouard (2014) during intensity change. Part I: Relationship between the thermodynamic structure and precipitation. *Mon. Wea. Rev.*, **144**, 3333–3354, doi: 10.1175/MWR-D-16-0017.1.
- Zhang, J. A., R. F. Rogers, P. Reasor, E. Uhlhorn, and F. D. Marks Jr., 2013: Asymmetric hurricane boundary layer structure from dropsonde composites in relation to the environmental wind shear. *Mon. Wea. Rev.*, **141**, 3968–3984, doi: 10.1175/MWR-D-12-00335.1.
- Zhang, J.A., J. J. Cione, E. A. Kalina, E.W. Uhlhorn, T. Hock, and J.A. Smith, 2017: Observations of infrared sea surface temperature and air-sea interaction in Hurricane Edouard (2014) using GPS dropsondes. *J. Atmos. Oceanic Technol.*, **0**, doi: 10.1175/JTECH-D-16-0211.1.