

**Quantitative estimates of species composition and
abundance of fishes, and fish species/habitat
associations in St. Croix, U.S. Virgin Islands**

Dr. Wesley Toller

Division of Fish and Wildlife
Department of Planning and Natural Resources
US Virgin Islands

Final Report: Patterns of habitat utilization by reef fish on St. Croix
F-7-17, Study3
Period: 1 October 2001 to 30 September 2002

Quantitative estimates of species composition
and abundance of fishes, and fish
species/habitat associations in
St. Croix, U.S. Virgin Islands

Dr. Wesley Toller
Fisheries Biologist II

Division of Fish and Wildlife
Department of Planning and Natural Resources
US Virgin Islands

November 2002

ABSTRACT

Reef fish communities around St. Croix, U.S. Virgin Islands were studied by visual census methods in order to identify fisheries habitat links, to document the role of local coral reefs as essential fish habitat, and to provide a baseline for monitoring studies. Seven reef sites were surveyed by stationary fish censuses during spring and fall of 2002. In general, the fish communities were similar across sites and survey periods: all reefs supported a diverse and abundant fish assemblage that was largely replicated between spring and fall surveys. Planktivorous fish (primarily labrids and pomacentrids) dominated the fauna numerically. Herbivorous fish (acanthurids and scarids) were common and abundant at all sites. Piscivores and other specialized feeders were least abundant. Serranids were common and relatively diverse, however most observed fish were either diminutive or small species (e.g. harlequin bass, hamlets, coney, and graysby). Lutjanids were of larger size, but less common, less abundant, and less diverse than serranids. Very few large reef-associated piscivorous species were seen, and the significance of this observation is discussed.

Some variation in fish assemblages was observed among reefs. Comparison of species richness and diversity (H') among reefs showed a suggestive positive relationship to vertical relief, but no obvious relation to percentage live coral cover or percentage turf alga cover. Planktivorous pomacentrids predominated at the two reef sites with the highest coral cover. Acanthurids and scarids, while both nearly ubiquitous, showed a reciprocal density relationship, with scarids predominating on well-developed reefs. Chaetodontids were more abundant and diverse near a submarine canyon (Salt River). Lutjanids were not observed at Lang Bank — an offshore platform with sparse coral cover.

These results imply that at least some members of reef fish communities respond to presently unidentified biotic or abiotic differences among the coral reef habitats of St. Croix, and that the source of this variation warrants additional investigation. Additional studies are recommended to monitor changes in these reef fish communities over time, to improve assessments of rare (but recreationally and commercially important) species, and to further elucidate critical fish-habitat links.

| | Page No. |
|---|-----------------|
| Abstract | ii |
| Table of Contents | iii |
| List of Tables | iv |
| List of Figures | iv |
| | |
| INTRODUCTION | 1 |
| METHODS | 1 |
| Description of Study Sites | 1 |
| Visual Census Methods | 2 |
| | |
| RESULTS | 3 |
| DISCUSSION | 5 |
| Planktivores | 6 |
| Herbivores | 6 |
| Piscivores, Carnivores and Specialized Invertebrate Feeders | 7 |
| Recommendations | 8 |
| | |
| ACKNOWLEDGEMENTS | 8 |
| | |
| LITERATURE CITED | 8 |
| | |
| APPENDICES | |
| 1. Distribution of observations on 85 fish species across seven St. Croix reef sites: | 19-23 |
| | |
| 2. A. Fish Census Data from Lang Bank | 24-25 |
| B. Fish Census Data from Jacks & Isaacs Bay | 26-28 |
| C. Fish Census Data from East Wall, Salt River Canyon | 29-31 |
| D. Fish Census Data from West Wall, Salt River Canyon | 32-34 |
| E. Fish Census Data from Eagle Ray | 35-38 |
| F. Fish Census Data from Cane Bay | 39-41 |
| G. Fish Census Data from Sprat Hole | 42-44 |

| List of Tables | | Page No. |
|-----------------------|---|----------|
| 1. | Description of St. Croix reef sites for fish census study. | 15 |
| 2. | Fish census effort | 15 |
| 3. | Total number of fish and species observed during stationary fish censuses | 16 |
| 4. | The 25 most abundant fish species observed in this study. | 17-18 |

| List of Figures | | |
|------------------------|--|----|
| 1. | Map of St. Croix showing the seven reef sites where fish visual censuses were conducted. | 11 |
| 2. | Summary of fish community structure at seven St. Croix reef sites. | |
| A. | Diversity (H') | 12 |
| B. | Species Richness | 12 |
| C. | Fish Abundance | 12 |
| 3. | Fish abundance by family at seven St. Croix reef sites. | 12 |
| 4. | Abundance of fishes representing 8 families at 7 St. Croix reef sites. | |
| A. | Labridae (wrasses) | 13 |
| B. | Pomacentridae (damsel fishes) | 13 |
| C. | Acanthuridae (surgeonfishes) | 13 |
| D. | Scaridae (parrotfishes) | 13 |
| E. | Serranidae (groupers) | 14 |
| F. | Lutjanidae (snappers) | 14 |
| G. | Haemulidae (grunts) | 14 |
| H. | Chaetodontidae (butterflyfishes) & Pomacanthidae (angelfish) | 14 |

INTRODUCTION

Coral reefs of the U.S. Virgin Islands support a diverse assemblage of fishes. In addition to their ecological roles on reefs, these fish have economic importance to local recreational and commercial fisheries (Appeldoorn et al. 1992).

Management of reef fish resources depends upon sound information about the status of local populations, and knowledge of critical links between habitat and fish life-history (i.e. essential fish habitats). Thus a first step is to simply establish the fine-scale patterns of fish distribution among habitats. A descriptive study, such as this one, can provide valuable information about habitat-species associations, which may ultimately lead to identification of critical links.

A second objective of management must be to monitor resources, so that the trajectory of populations through time can be evaluated. A thorough and descriptive baseline survey, whether it represents a natural or disturbed community (Jackson 1997), is the essential starting point for any monitoring program. On St. Croix, for example a proposed Marine Protected Area (MPA) will encompass the nearshore, eastern end of the island. Implementation of a broad "No-Take" zone may conceivably influence the structure of entire reef fish communities located in this area. Monitoring will be instrumental in gauging the effects of this proposed MPA on fish communities. Alternatively, on a scale of individual species, monitoring may allow managers to determine whether specific fisheries regulations, such as localized closures to protect spawning aggregations, are effective at restoring depleted stocks (*sensu* Tobias et al. 1988).

METHODS

Description of Study Sites

Seven reef sites around the island of St. Croix, U. S. Virgin Islands were selected for this study (see Figure 1): Lang Bank (LB), Jacks and Isaacs Bay (JI), East Wall (EW) of Salt River Canyon, West Wall (WW) of Salt River Canyon, Eagle Ray (ER) on Long Reef, Cane Bay (CB), and Sprat Hole (SH). Each of these reefs is surveyed annually by the Center for Marine and Environmental Studies, University of the Virgin Islands, as part of their coral reef monitoring program. Those studies provide much of the descriptive information on sessile reef biota, as presented in Table 1 (Nemeth and Herzlieb 2002). Detailed geological, biological, and spatial descriptions of these reef sites have been made previously, either in a general sense (Hubbard 1989a), in detail (at CB - Hubbard 1989b; at EW and WW - Hubbard 1989c) or at comparable nearby reefs (at JI Hubbard 1989d). Briefly, the seven reef sites all represent shallow water (< 15 m depth) hard bottom substrates situated on the insular shelf of St. Croix. Despite geographic proximity, there is considerable variation among reefs in terms of benthic flora and fauna, physical structure, exposure to prevailing currents and wave forces, and adjacent habitats (to name but a few of the variables). For example, the development of scleractinian

coral communities varies considerably among sites, from about 5% living coral cover at LB to >24% at SH (Nemeth and Herzlieb 2002). Some of the reefs have previously been classified (Hubbard 1989a-d) as barrier-type reefs (ER), fringing reefs (JI), shelf edge reefs (CB, SH), or submarine hardgrounds (LB). In the present study, only those attributes presented in Table 1 were considered in relation to reef fish community structure.

Visual Census Methods

The stationary fish census method (Bohnsack and Bannerot 1986) was used to assess reef fish community structure. Compared to other visual census methods (belt transects, timed random swim methods), the stationary count offers the advantage of estimating fish community structure as a function of density per unit area for each observed species, while enabling size (length) estimates of individuals. Generally, results from stationary point counts are comparable to results obtained from belt transects (e.g. Bortone et al. 1989), and the two methods differ primarily in duration (belt transects are faster) and area surveyed per replicate. A typical belt transect of 30 m x 2 m yields 60 m² whereas a 15 m wide stationary census (used in this study) yields 176 m² per replicate. However neither of these two methods is likely to enumerate all of the species present within a given area (i.e. to generate a species list) - that purpose is best served using replicated timed random diver swims (Kimmel 1985) or related methods (see Rogers et al 2001).

The census protocol is only slightly modified from Bohnsack and Bannerot (1986), and a brief description is provided here. A 15 m diameter census "cylinder" was defined by transect tape with the observer positioned in the center (7.5 m mark). Fish within this cylinder were censused as follows. During an initial 5-minute "listing" period, the names of all observed fish species were recorded onto pre-printed data forms. At the end of the listing period, the observer began enumerating all individuals of each species, working from the bottom of the list upward and making one 360° sweep. Strict adherence was made to the 5-minute listing period. New species observed during the enumeration period (e.g. fish that swam into the cylinder after 5 minutes of listing) were not included in the stationary count data. Divers estimated fish total length to the nearest cm, using a measuring "T-bar" for underwater visual reference. Maximum, minimum, and average size was recorded for each species. Divers also recorded a brief description/sketch of habitat features within the census area. All fish were identified to species, however gobies (Gobiidae) and blennies (Blenniidae) were excluded from census counts. To insure that our fish census areas coincided with ongoing UVI coral surveys, but that replicates did not cover overlapping areas, census stations were marked with a small piece of rebar driven into nonliving reef substrate.

Surveys were conducted during two sampling periods: Spring (May - June, 2002) and Fall (September, 2002). For each reef site, six replicate censuses were made [this is a level of replication thought to adequately sample the species representing > 90% of all individuals at a site (Bohnsack and Bannerot 1986)] except for LB (Spring) where only four replicate censuses were conducted due to unfavorable weather conditions. Six replicate censuses surveyed a total area of 1,056 m² per reef site. A summary of census dates, number of replicates, and census duration is presented in Table 2.

Data from this study were analyzed using Microsoft Excel software.

RESULTS

A total of 35,322 fish were sighted during the surveys (Table 3). More fish were observed in fall (16,859) than spring (18,463). The number of fish observed at each site ranged from 1,385 (JI, fall) to 5,443 (CB, fall). A total of 85 fish species representing 27 families were sighted (Table 3; Appendix 1). More species were seen during fall (77 species) than during spring (72 species; Table 3). The number of species observed at each site during spring or fall ranged from 24 (LB, spring) to 42 (WW, ER, CB; all fall) and the cumulative number of species observed at each reef site ranged from 32 (LB) to 52 (ER).

In general, fish assemblages present at each reef site were similar between spring and fall. Average fish abundance (Figure 2A) remained relatively constant at 6 of the reef sites. At CB, an increase in average abundance in fall was due to a large influx of small (~3 to 5 cm, TL) creole wrasses (*Clepticus parrae*; see Appendix 2F). Average species richness also remained relatively constant between spring and fall (Figure 2B). However the Shannon-Weiner diversity index (H'), computed as an average per census, showed greater variability between survey periods (Figure 2C; LB and ER), suggesting that community composition had changed in some cases. At LB, change was attributed to a large (30%) decrease in abundance of the numerically dominant species, the bluehead wrasse, *Thalassoma bifasciatum* and a 3-fold increase in bicolor damselfish (*Stegastes partitus*) abundance. At ER, the fall increase in diversity corresponded to a 40% decrease in abundance of the bluehead wrasse and the appearance of large schools of brown chromis, *Chromis multilineata* (an increase from 1.7 to 43.3 fish/census; Appendix 2E).

In contrast to temporal comparisons (i.e. spring vs. fall), comparisons across reef sites showed greater dissimilarity in fish assemblages (Figure 2). Average abundance (Figure 2A) was highest at CB and SH, sites with the highest coral cover, but also at LB where coral cover is sparse. Average species richness varied from ~15 to ~22 species observed/census (Figure 2B) and LB had the fewest species while ER, CB and SH had the most. Diversity indices showed a similar trend (Figure 2C).

The 25 most abundant species observed in this study are present in Table 4. The bluehead wrasse alone accounted for 38.6% of all fish sightings. The five most abundant species were the bluehead wrasse, creole wrasse, bicolor damselfish, brown chromis, and blue chromis (*Chromis cyanea*) - all primarily planktivorous species. Together, they accounted for 87.2% of all fish sightings. Despite comparably high abundance levels shared among these five species, their distribution and frequency of occurrence were different (Table 4). While the strongly reef-associated bluehead wrasse and bicolor damselfish were ubiquitous (present at each reef site) and common (frequency > 97% of censuses), the schooling creole wrasse and brown chromis were more patchy in distribution (Appendix 1) and less common (frequency < 45% of censuses).

A detailed account of species is beyond the scope of the present work (see Appendix 2A to G): Here, an analysis of reef fish community composition is presented based upon taxonomic

divisions (families) that delineate eight commercially, recreationally or ecologically important fish groups. These families are: Labridae (wrasses), Pomacentridae (damselfishes), Acanthuridae (surgeonfishes), Scaridae (parrotfishes), Serranidae (groupers), Lutjanidae (snappers), Haemulidae (grunts), Chaetodontidae (butterflyfishes) and Pomacanthidae (angelfishes). The latter two families were combined as a single group (butterflyfish and angelfish) for these analyses. Use of these taxonomic divisions is justified because: (1) each family loosely represents a trophic (ecological) group, and (2) the families collectively represented over 95% (usually >98%) of fish observed at each reef site (e.g. Table 4; Appendices 2A to G)

Fish abundance by family is summarized in Figure 3, where data were pooled from all seven reef sites. Average fish abundance was similar between spring and fall, however abundance varied considerably among families. Labrids and Pomacentrids were over 10 times more abundant than all other families (~ 100 fish observed/census). Acanthurids and scarids had similar abundance levels of ~10 fish/census. The abundance of serranids, lutjanids, haemulids, chaetodontids and pomacanthids was generally low (1 to 4 fish/census) and typically quite variable within and among sites (see below).

To examine fish community structure across the seven reef sites, the same eight fish families were utilized as a basis for analyses (Figure 4). Labrids were abundant at all sites (average 242 fish/census) but highly variable within and among sites (Figure 4a) [as noted above, an exceptionally high abundance of creole wrasses was recorded at CB in spring]. Pomacentrids were also abundant (average 148 fish/census), and showed the greatest variation among sites. The highest abundances were seen at CB and SH, where > 300 fish/census were recorded in spring and fall (Figure 4B), Blue chromis and brown chromis were the most abundant pomacentrids at CB and SH (Appendix 2F and G). Herbivorous fish were less abundant than planktivores. The abundance of acanthurids (average 10.7 fish/census) and scarids (average 9.6 fish/census) was also variable across the reef sites (Figures 4C and D). Acanthurids had their highest abundances at WW, EW, JI, and LB and were least abundant at CB and SH. Scarids had their highest abundance at SH.

Predatory fishes (piscivores, carnivores) are represented by the serranids, lutjanids and haemulids. As a family, serranids had low abundance (average 4 fish/census) that showed little variation among reef sites (Figure 4E). Serranid diversity was variable among reef sites. The coney, *Cephalopholis fulvus*, was uniformly distributed across reefs (Appendix 1) and relatively common (Appendix 2A-G), however the graysby, *C. cruentatus*, was more frequent in habitats of high topographic complexity (ER, CB, SH). The red hind, *Epinephelus guttatus*, was observed only infrequently at 4 sites (Appendices 1 and 2). Hamlets (genus *Hypoplectrus*) also showed highest diversity levels at SH, ER, EW and WW but not CB. In contrast to serranids, lutjanids showed greater variation to in abundance and distribution. The yellowtail snapper, *Ocyurus chrysurus*, and the mahogany snapper, *Lutjanus mahogoni*, were the most common and abundant snappers, however the schoolmaster, *L. apodus*, and the mutton snapper, *L. analis*, were also observed (Appendix 1). No lutjanids were recorded at LB. Haemulid abundance was relatively uniform (average 2.4 fish/census) across reef sites (Figure 4G).

The combined group of Chaetodontidae and Pomacanthidae represent long-lived reef residents with specialized feeding habits (e.g. feeding on cnidarians, sponges). Abundance and diversity

of this group varied among reef sites (Figure 4H) and was generally low (2.3 fish/census). Chaetodontids were most frequent and abundant at EW, WW and ER. It is interesting that EW and WW had the highest diversity of butterflyfish (4 and 3 species, respectively) whereas other reef sites had 2 or 1 species. Only the banded butterflyfish, *Chaetodon striatus*, was recorded from LB and JI, whereas only the foureye butterflyfish, *C. capistratus*, was recorded from CB and SH (Appendix 2).

DISCUSSION

The reefs investigated in this study supported diverse and abundant fish communities. This is not surprising - previous studies have documented well over 200 reef-associated species in St. Croix waters (Clavijo et al. 1980), and over 100 species may occur on a single patch reef (Gladfelter and Gladfelter 1978). For the following discussion, it is important to note that the stationary visual census method does not capture total reef fish diversity (i.e. Appendices 2A to G do not represent complete species lists). Rather, these are quantitative data on the most conspicuous and abundant components of each fish community. The absence of a given species from our observations should be considered "relative" absence. Nonetheless, the average number of species observed per census is a quantitative estimate of community species richness, and a useful measure for comparing reef sites.

The goal of the present study was to document, in an exploratory manner, species-habitat links of coral reef fish populations. This experimental design allows only a preliminary analysis of the relationship between individual fish species (or fish communities) and specific reef attributes (e.g. % live coral, vertical relief, benthic invertebrate community composition, wave exposure, prevailing currents, etc). When fish communities were compared across the seven reef sites, there was no obvious relation of average fish abundance to percentage live coral nor percentage turf algal cover (not shown). Planktivore abundance (blue chromis and brown chromis), on the other hand, may be positively related to percentage live coral cover (e.g. CB and SH; Figure 4B). Species richness showed a more suggestive positive relation to coral cover, although ER represents an important exception. At ER, coral cover is relatively low, but vertical relief is high (a spur and groove area formed of *Montastraea* skeletons). A relatively high number of fish species was observed at ER. This suggests that the richness of St. Croix reef fish communities may be influenced more by three-dimensional architecture (topographic complexity) of the reef than by the presence of living (versus dead) corals. This idea is not new (Roberts and Ormond 1987), and it has been suggested by results from long-term studies of fish populations at Buck Island, St. Croix (Gladfelter et al. 1992), where hurricane damage and disease permanently reduced live coral cover (*Acropora palmata*), but left habitat structural complexity relatively intact. Despite the loss of living coral, resident fish populations recovered to pre-hurricane densities.

Nemeth and Herzlieb (2002) suggested that reef sites along St. Croix's northern coast harbor a greater abundance of reef fish than the other reef sites. Our results did not corroborate this finding (see Figure 2 and Figure 4). Neither that study nor this one included sufficient replication of reef sites, adequate spatial coverage of insular reefs, nor control over confounding variables (e.g. topographic complexity) to reach a definitive conclusion. Nonetheless, previous studies indicate that recruitment patterns are highest on St. Croix's northwestern and southeastern shores (Caselle and Warner 1996, and Swearer 2002). Recruitment studies, in

conjunction with community assessments, may clarify a postulated relationship between larval supply and resident reef fish abundance.

Planktivores:

The reef fish communities were numerically dominated by labrids (bluehead wrasse, creole wrasse) and planktivorous pomacentrids (bicolor damselfish, blue chromis, and brown chromis), which were collectively 10 to 100-fold more abundant than all other species. The bluehead wrasse, perhaps the most abundant of all West Indian reef fishes, is a facultative planktivore, and the remaining species feed primarily on zooplankton (Randall 1968).

Large populations of planktivores form a major trophic link between coral reef habitats and open-water communities (e.g. Hobson 1991). Undoubtedly, a large number of these fish fall prey to resident and reef associated predatory fishes. Between spring and fall, bluehead wrasse abundance decreased by ~ 20%, which is suggestive of heavy predation. However we found no relation between planktivore abundance and piscivore abundance (compare Figure 4A and B with 4E and F). Some authors have argued that, rather than predation, the primary trophic link between planktivores and coral reefs is through production of feces, which is then consumed by a wide variety of reef fish and invertebrates (Robertson 1982). Nonetheless, the role of planktivores as forage for commercially and recreationally important species should be investigated further.

Herbivores:

Herbivorous fish were common, moderately abundant, and relatively diverse on all seven reefs. For acanthurids and scarids, average abundance was similar (9.6 fish/176 m² census and 10.7 fish/176 m² census, respectively). In 2001, Nemeth and Herzlieb (2002) censused fish from the same sites using belt transects. When normalized to 100m², the density of acanthurids observed in our study was slightly lower than in theirs (6.0 vs 9.0 fish/100 m²) and the density of scarids observed in this study was substantially lower (5.5 vs 13 fish/100m²). The apparent decline in parrotfish populations is unexplained. Methodological differences between the studies may account for this discrepancy. Alternatively, scarid populations may be in decline due to overfishing – parrotfishes are actively sought by a local gillnet fishery. Given the important trophic role that scarids play in maintaining scleractinian coral growth and diversity (e.g. Pennings 1997), local parrotfish populations should be monitored carefully in the future.

In this study, parrotfishes (especially the redband parrotfish, *Sparisoma aurofrenatum*, and the princess parrotfish, *Scarus taeniopterus*) and surgeonfishes (ocean surgeonfish, *Acanthurus bahianus*, and blue tang, *A. coeruleus*) were nearly ubiquitous among the reef sites. Analysis of their abundance patterns (Figure 4C and D) suggests that their densities were reciprocal. Acanthurids were more abundant on low relief, low coral cover sites (LB, JI, EW, WW) while Scarids were more abundant on reefs with greater topographic complexity (ER, CB, SH). This pattern may arise from differences in feeding mode: acanthurids tend to browse over soft bottoms while scarids are scrapers of hardbottom substrates (e.g. Choat 1991).

Piscivores, Carnivores and Specialized Invertebrate Feeders:

A diversity of predatory fishes was recorded during censuses. The observed species varied in their degree of reef association. For example carangids (bar jack, *Carynx ruber*, and blue runner, *C. crysos*) and scombrids (cero mackerel, *Scomberomorus regalis*) were infrequently observed, in part due to their high mobility and their loose association with reefs. On the other hand, the coney appears to represent a strongly reef-associated resident species that displayed no habitat preference. This species was the most abundant serranid, occurring with moderate to high frequency at all studied reef sites. The coney is important to both the recreational and commercial fishery of the USVI (Appeldoorn et al. 1992; Bolden 1994). The graysby, another frequently observed serranid, showed habitat preference for reefs with a high degree of topographic complexity.

As adults, lutjanids are largely residents of reef habitats. However, the juveniles of many lutjanid species, as well as haemulids and chaetodontids, depend upon "nursery habitats" during their early post-settlement life. On St. Croix for example, Adams and Tobias (1993) documented the importance of an estuarine environment (Salt River) for juvenile schoolmasters (*L. apodus*), french grunts (*Haemulon flavolineatum*), and four-eye butterflyfish (*C. capistratus*). Likewise, St. Croix embayments (comprised of patch reef, sand and seagrass habitats) appear to function as important nursery habitats for French grunts and yellowtail snapper (Mateo and Tobias 2001). In our study, adults of these species (but not juveniles) were commonly observed on reefs. For example, adult French grunts were among the most frequently observed fish (~80% of all censuses), but no juveniles were observed these reef sites. For such species, these results serve to underscore the connectivity between adult populations on reefs and juvenile populations residing in nursery habitats.

Nonetheless, it is clear that many of the long-lived reef species were relatively rare. Commercially and recreationally important species such as the red hind (*E. guttatus*) or mutton snapper (*L. analis*) were observed only infrequently (6.1% and 3.7% of censuses, respectively). Some of the larger non-piscivorous reef species such as the french angelfish (*Pomacanthus paru*), queen angelfish (*Holacanthus ciliaris*), or puddingwife (*Halichoeres radiatus*) were also rare in our studies. Formerly, all of the above species were considered "common" on St. Croix reefs (Clavijo et al. 1980). Overfishing is the likely explanation for the relative rarity of these species. Establishment of a marine protected area should augment populations of long-lived fish species with high susceptibility to fishing pressures (Bohnsack 1993), such as those discussed above.

Crosby and Reese (1996) have suggested that butterflyfish density is an indicator of reef health. In our study, butterflyfish reached their greatest abundance and diversity at Salt River canyon. This distribution pattern was unique among fishes in our study. Localized abundance at Salt River Canyon may be due to reduced fishing pressure in the immediate vicinity – chaetodontids are readily caught in fish traps (DFW, unpubl. SEAMAP study) or to the proximity of nursery habitat (Salt River Estuary; see Adams and Tobias 1993), which may augment local butterflyfish populations. Alternatively, the chaetodontids species may share a habitat preference for some unidentified feature of the submarine canyon itself.

Recommendations:

Continued monitoring of these seven reef sites should be conducted semi-annually. One or two additional reef sites should be added to include reef fish communities along the southern insular shelf of St. Croix. The visual census method used here is robust and reproducible and should therefore be used in future monitoring efforts. However, a second assessment method, such as the tuned random diver swim (TRDS) method, should also be included in the protocol. Sufficient TRDS replicates (performed on an annual or biannual basis) would establish a species lists for each reef site and enable more powerful inferences regarding rare species.

Population assessments of large, resident reef fishes were compromised by the rarity of these fish. This is particularly troublesome because many are commercially and recreationally important species. However, for species such as the Nassau grouper (*Epinephelus striata*), goliath grouper (*E. itajara*), tiger grouper (*Mycteroperca tigris*), black grouper (*M. bonaci*), yellowfin grouper (*M. venenosa*), mutton snapper (*Lutjanus analis*), and cubera snapper (*L. cyanopterus*), populations are probably too low to adequately assess them by any of the aforementioned visual census methods. Instead, future monitoring efforts should be directed towards censusing these species during their annual spawning aggregations (e.g. Colin 1996). Aggregation sites are already known for at least two species on St. Croix (*L. analis* and *E. guttatus*) and these could be monitored using diver surveys or ROV.

ACKNOWLEDGMENTS

William Tobias provided technical guidance and assisted in many of the field surveys. Hector Rivera and Willy Ventura were boat operators. A special thanks goes to Dianne "Dee" Osinski and Robert Daniel Stone for their participation in (and enthusiasm for) the fish counts.

LITERATURE CITED

- Adams, A.J., and W.J. Tobias. 1993. Red mangrove prop-root habitat as a finfish nursery area. A case study of Salt River, St. Croix, U.S.V.I. *Proc. Gulf Carib. Fish. Inst.* 46:22-47
- Appeldoorn, R., J. Beets, J. Bohnsack, S. Bolden, D. Matos, S. Meyers, A. Rosario, Y. Sadovy, and W. Tobias. 1992. Shallow water reef fish stock assessment for the U.S. Caribbean. NOAA Tech. Memo. Niv S:SEFSC-304, 70 pp.
- Bohnsack, J.A. 1993. Marine reserves: they enhance fisheries, reduce conflicts, and protect resources. *Oceanus* 36(3): 63-71
- Bohnsack, J.A., and S.P. Bannerot. 1986. A stationary visual census technique for quantitatively assessing community structure of coral reef fishes. NOAA Technical Report NMFS 41, U.S. Dept of Commerce, 15 pp.
- Bortone, S.A., J.J. Kimmel, and C.M. Bundrick. 1989. A comparison of three methods for visually assessing reef fish communities: time and area compensated. *Northeast Gulf Sci.* 10(2):85-96.

- Caselle, J.E., and R.R. Warner. 1996. Variability in recruitment of coral reef fishes: the importance of habitat at two spatial scales. *Ecology* 77: 2488-2504.
- Choat, J.H. 1991. The biology of herbivorous fishes on coral reefs. Pp. 120-155 *In*: P. F. Sale (ed.) *The Ecology of Fishes on Coral Reefs*. Academic Press, Inc., New York, 54 pp
- Clavijo, I.E., J.A. Yntema, and J.C. Ogden. 1980. An annotated checklist of the fishes of St. Croix U.S. Virgin Islands. West Indies Laboratory, St. Croix, USVI, 49 pp.
- Colin, P.L. 1996. Longevity of some coral reef fish spawning aggregations. *Copeia* 1996(1): 189-2.
- Crosby, M.P., and E.S. Reese. 1996. A manual for monitoring coral reefs with indicator species: butterflyfishes as indicators of change on Indo Pacific reefs. Office of Ocean and Coastal Resource Management, NOAA, Silver Springs MD, 45 pp.
- Gladfelter, W.B., and E.H. Gladfelter. 1978. Fish community structure as a function of habitat structure on West Indian patch reefs. *Rev. Biol. Trop.* (suppl. 1):65-84.
- Gladfelter, E.H., J.C. Bythell, and Z.M. Hillis. 1992. Changes in fish assemblage structure at Buck Island, St. Croix, U. S. Virgin Islands from 1980-1990: an indication of predictability in coral reef fish assemblages based on known habitat changes? Pages 14-29 *In*: *Ecological Studies of Buck Island Reef National Monument, St. Croix, U. S. Virgin Islands. Part II. Report prepared for U.S. Department of the Interior, National Park Service.*
- Hobson, E.S. 1991. Trophic relationships of fishes specialized to feed on zooplankters above coral reefs. Pp. 69-95 *In*: P.F. Sale (ed.) *The Ecology of Fishes on Coral Reefs*. Academic Press, Inc., New York, 754 pp.
- Hubbard, D. 1989a. Modern carbonate environments of St. Croix and the Caribbean: A general overview. Pages 85-94 *In*: D. Hubbard (ed.) *Terrestrial and marine geology of St. Croix, U.S. Virgin Islands. Sp. Publ. No. 8, West Indies Laboratory, St. Croix, USVI, 213 pp.*
- Hubbard D. 1989b. The shelf-edge reefs of Davis and Cane Bays, Northwestern St. Croix, U.S.V.I. Pages 167-179 *In*: D. Hubbard (ed.) *Terrestrial and marine geology of St. Croix, U. S. Virgin Islands. Sp. Publ. No. 8, West Indies Laboratory, St. Croix, USVI, 213 pp.*
- Hubbard, D. 1989c. Depositional environments of Salt River Estuary and submarine canyon, St. Croix, U. S:V.I. Pages 181-196 *In*: D. Hubbard (ed.) *Terrestrial and marine geology of St. Croix, U.S. Virgin Islands. Sp, Publ. No. 8, West Indies Laboratory, St. Croix, USVI, 213 pp.*
- Hubbard, D. 1989d. Depositional environments of the Turner Hole reef complex. Pages 155-160 *In*: D. Hubbard (ed.) *Terrestrial and marine geology of St. Croix, U.S. Virgin Islands. Sp. Publ. No. 8, West Indies Laboratory, St. Croix, USVI, 213 pp.*
- Jackson, J.B.C. 1997. Reefs since Columbus. *Coral Reefs* 16:S23-S32.

- Kimmel, J.J. 1985. A new species-time method for visual assessment of fishes and its comparison with established methods. *Env. Biol. Fishes* 12:23-32.
- Mateo, I. and W.J. Tobias. 2001. The role of nearshore habitats as nursery grounds for juvenile fishes on the northeast coast of St. Croix. *Proc. Gulf Carib. Fish. Inst.* 52:512-530.
- Nemeth, R.S., and S. Herzlieb. 2002. Video monitoring assessment of coral reefs in proposed marine parks St. Croix, United States Virgin Islands. Center for Marine and Environmental Studies, University of the Virgin Islands, St. Thomas, USVI. *In: U.S. Virgin Islands Coral Reef Monitoring Project, Year 1 Final Report USVI Department of Planning and Natural Resources, Division of Coastal Zone Management.*
- Pennings, S.C. 1997. Indirect interactions on coral reefs. Pp. 249-272 *In: C. Birkeland (ed) Life and Death of Coral Reefs.* Chapman & Hall, New York, 536 pp.
- Randall, J.E. 1968. Caribbean Reef Fishes. T.F.H. Publications, Inc., Jersey City, NJ, 318 pp.
- Roberts, C.M. and R.F.G. Ormond. 1987. Habitat complexity and coral reef fish diversity and abundance on Red Sea fringing reefs. *Mar. Ecol. Prog. Ser.* 41:1-8.
- Robertson, D.R. 1982. Fish feces as food on a Pacific coral reef *Mar. Ecol. Prog. Ser.* 7:253-265.
- Rogers, C.S., G. Garrison, R. Grober, Z. Hillis, and M.A. Franke. 2001. Coral Reef Monitoring Manual for the Caribbean and Western Atlantic. 2nd Ed., Virgin Islands National Park.
- Swearer, S.E. 2000. Self-recruitment in coral reef fish populations. Ph.D. dissertation, University of California, Santa Barbara, 151 pp.
- Tobias, W., E. Telemaque, and M. Davis. 1988. Buck Island fish and shellfish populations. Biosphere Reserve Research Report No. 26, Virgin Islands National Park, 27 pp.

Figure 1. Map of St. Croix showing the seven reef sites where fish visual censuses were conducted. Abbreviations are as follows: Lang Bank (LB), Jacks and Isaacs Bay (JI), East Wall (EW) of Salt River Canyon, West Wall (WW) of Salt River Canyon, Eagle Ray (ER) on Long Reef, Cane Bay (CB), and Sprat Hole (SH).




Figure 2. Summary of fish community structure at seven St. Croix reef sites.




Figure 3. Fish abundance by family at seven St. Croix reef sites. Error bars show range of average abundance values across sites.




Figure 4. Abundance of fishes representing 8 families at 7 St. Croix reef sites. Families are: A. Labridae (wrasses), B. Pomacentridae (damselfishes), C. Acanthuridae (surgeonfishes), D. Scaridae (parrotfishes), E. Serranidae (groupers), F. Lutjanidae (snappers), G. Haemulidae (grunts), and H. Chaetodontidae (butterfly fishes) and Pomacanthidae (angelfishes). Error bars represent standard deviation.






Figure 4 (continued). Abundance of fishes representing 8 families at 7 St. Croix reef sites. Families are: A. Labridae (wrasses), B. Pomacentridae (damselfishes), C. Acanthuridae (surgeonfishes), D. Scaridae (parrotfishes), E. Serranidae (groupers), F. Lutjanidae (snappers), G. Haemulidae (grunts), and H. Chaetodontidae (butterflyfishes) and Pomacanthidae (angelfishes). Error bars represent standard deviation.


E.



F.



G.



H.




Table 1. Description of St. Croix reef sites for fish census study.

| Reef Site | Lang Bank | Jack's & Isaac's | East Wall (Salt River) | West Wall (Salt River) | Eagle Ray (Long Reef) | Cane Bay | Sprat Hole |
|---------------------------------|-----------------------|------------------|-----------------------------|------------------------|-----------------------|------------------|------------------|
| Abbreviation | LB | Jl | EW | WW | ER | CB | SH |
| Location | East | Southeast | North | North | North | Northwest | West |
| Latitude | N 17° 44.448' | N 17° 44.586' | N 17° 47.221' | N 17° 47.116' | N 17° 45.688' | N 17° 46.433' | N 17° 44.038' |
| Longitude | W 64° 32.186' | W 64° 34.310' | W 64° 45.445' | W 64° 45.564' | W 64° 41.929' | W 64° 48.810' | W 64° 53.722' |
| Min. Depth | 44 | 36 | 32 | 24 | 22 | 24 | 30 |
| Max. Depth | 48 | 49 | 45 | 27 | 35 | 36 | 35 |
| Avg. Depth | 45.8 | 41.7 | 36.8 | 25.3 | 28.0 | 28.8 | 32.3 |
| Reef Description ¹ | Submarine Hardgrounds | Fringing Reef | Sloping Cobble, Canyon Wall | Steep Canyon Wall | Barrier Reef | Shelf-Edge Reef | Shelf-Edge Reef |
| Vertical Relief ² | Low | Moderate | Moderate | Low & High | High | High | High |
| % Cover Live Coral ³ | ~ 5% | ~ 9% | ~ 12 % | ~ 11% | ~ 7% | ~ 23% | ~ 25% |
| % Cover Turf Algae ³ | ~ 80% | ~ 50% | ~ 72% | ~ 78% | ~ 70% | ~ 60% | ~ 60% |

*notes:

1. After Hubbard (1989a-d)
2. From diver observations (this study)
3. From Nemeth and Herzlieb (2002)

Table 2. Fish census effort

| | LB | Jl | EW | WW | ER | CB | SH |
|-------------------------|--------|--------|------------|------------|------------|---------|---------|
| Spring 2002 | | | | | | | |
| Date of Censuses | 29-May | 11-Jun | 23, 28-May | 22, 23-May | 3--May | 15-May | 16-May |
| No. of censuses | 4 | 6 | 6 | 6 | 6 | 6 | 6 |
| Total Census Time (min) | 91 | 195 | 187 | 190 | 233 | 242 | 255 |
| Avg Time / Census (min) | 22.8 | 32.5 | 31.2 | 31.7 | 38.8 | 40.3 | 42.5 |
| Fall 2002 | | | | | | | |
| Date of Censuses | 3-Sept | 4-Sept | 6-Sept | 12-Sept | 4 & 6-Sept | 10-Sept | 11-Sept |
| No. of censuses | 6 | 6 | 6 | 6 | 6 | 6 | 6 |
| Total Census Time (min) | 197 | 179 | 225 | 254 | 309 | 294 | 285 |
| Avg Time / Census (min) | 32.8 | 29.8 | 37.5 | 42.3 | 51.5 | 49.0 | 47.5 |

Table 3. Total number of fish and species observed during stationary fish censuses *1

| Reef Site | Survey | LB | JI | EW | WW | ER | CB | SH | Total |
|----------------|------------|---------|-------|-------|-------|-------|-------|-------|--------|
| No. of Fish | Spring '02 | 1,659*2 | 1,921 | 1,755 | 1,922 | 2,156 | 3,120 | 4,326 | 16,859 |
| Observed | Fall '02 | 2,491 | 1,385 | 1,456 | 2,221 | 2,148 | 5,443 | 3,319 | 18,463 |
| | Total | 4,150 | 3,306 | 3,211 | 4,143 | 4,304 | 8,563 | 7,645 | 35,322 |
| No. of Species | Spring '02 | 24 | 39 | 37 | 34 | 40 | 32 | 40 | 72 |
| Observed | Fall '02 | 28 | 37 | 38 | 42 | 42 | 42 | 40 | 77 |
| | Cumulative | 32 | 47 | 47 | 48 | 52 | 44 | 49 | 85 |

Notes:

1. Values reported for Numbers of species are not additive because there is substantial overlap in species compositions among different reef sites and between sampling periods.
2. Total number of fish from 4 replicate censuses.

Table 4. The 25 most abundant fish species observed in this study.

| Common Name | Species | Family | Spring '02 | | | Fall '02 | | | Total No. of Fish | % of All Obs* |
|-----------------------|---|----------------|----------------------------|-------------------------|---------------|----------------------------|-------------------------|---------------|-------------------|---------------|
| | | | Total No. of Fish Observed | Avg No. Fish per Census | % Freq (n=40) | Total No. of Fish Observed | Avg No. Fish per Census | % Freq (n=42) | | |
| bluehead wrasse | <i>Thalassoma bifasciatum</i> | Labridae | 7,790 | 194.8 | 97.5 | 5,830 | 138.8 | 97.6 | 13,620 | 38.6 |
| creole wrasse | <i>Clepticus parrae</i> | Labridae | 2,010 | 50.3 | 45 | 3,635 | 86.5 | 38.1 | 5,645 | 16.0 |
| bicolor damselfish | <i>Stegastes partitus</i> | Pomacentridae | 1,466 | 36.7 | 100 | 2,485 | 59.2 | 100.0 | 3,951 | 11.2 |
| brown chromis | <i>Chromis multilineata</i> | Pomacentridae | 1,661 | 41.5 | 40 | 2,172 | 51.7 | 40.5 | 3,833 | 10.9 |
| blue chromis | <i>Chromis cyanea</i> | Pomacentridae | 1,864 | 46.6 | 75 | 1,880 | 44.8 | 76.2 | 3,744 | 10.6 |
| ocean surgeonfish | <i>Acanthurus bahianus</i> | Acanthuridae | 367 | 9.2 | 97.5 | 287 | 6.8 | 97.6 | 654 | 1.9 |
| yellowhead wrasse | <i>Halichoeres garnotti</i> | Labridae | 162 | 4.1 | 72.5 | 149 | 3.5 | 81.0 | 311 | 0.9 |
| black durgon | <i>Melichthys niger</i> | Balistidae | 94 | 2.4 | 50 | 201 | 4.8 | 52.4 | 295 | 0.8 |
| redband parrotfish | <i>Sparisoma aurofrenatum</i> | Scaridae | 144 | 3.6 | 90 | 139 | 3.3 | 83.3 | 283 | 0.8 |
| princess parrotfish | <i>Scarus taeniopterus</i> | Scaridae | 138 | 3.5 | 87.5 | 138 | 3.3 | 85.7 | 276 | 0.8 |
| dusky damselfish | <i>Stegastes adustus</i> (<i>S. fuscus</i>) | Pomacentridae | 150 | 3.8 | 60 | 100 | 2.4 | 50.0 | 250 | 0.7 |
| threespot damselfish | <i>Stegastes planifrons</i> | Pomacentridae | 111 | 2.8 | 22.5 | 94 | 2.2 | 21.4 | 205 | 0.6 |
| coney | <i>Cephalopholis fulvus</i> | Serranidae | 97 | 2.4 | 72.5 | 108 | 2.6 | 81.0 | 205 | 0.6 |
| sergeant major | <i>Abudefduf saxatilis</i> | Pomacentridae | 47 | 1.2 | 32.5 | 157 | 3.7 | 42.9 | 204 | 0.6 |
| blue tang | <i>Acanthurus coeruleus</i> | Acanthuridae | 89 | 2.2 | 87.5 | 106 | 2.5 | 90.5 | 195 | 0.6 |
| french grunt | <i>Haemulon flavolineatum</i> | Haemulidae | 93 | 2.3 | 82.5 | 85 | 2.0 | 78.6 | 178 | 0.5 |
| yellow goatfish | <i>Mulloidichthys martinicus</i> | Mullidae | 14 | 0.4 | 17.5 | 158 | 3.8 | 23.8 | 172 | 0.5 |
| foureye butterflyfish | <i>Chaetodon capistratus</i> | Chaetodontidae | 59 | 1.5 | 57.5 | 61 | 1.5 | 54.8 | 120 | 0.3 |
| stoplight parrotfish | <i>Sparisoma viride</i> | Scaridae | 36 | 0.9 | 55 | 62 | 1.5 | 64.3 | 98 | 0.3 |
| yellowtail damselfish | <i>Microspathodon chrysurus</i> | Pomacentridae | 46 | 1.2 | 60 | 48 | 1.1 | 52.4 | 94 | 0.3 |

Table 4 (continued). The 25 most abundant fish species observed in this study.

| Common Name | Species | Family | Spring '02 | | | Fall '02 | | | Total No. of Fish | % of All Obs* | |
|-----------------------------|--------------------------------------|---------------|----------------------------|-------------------------|---------------|----------------------------|-------------------------|---------------|-------------------|---------------|------|
| | | | Total No. of Fish Observed | Avg No. Fish per Census | % Freq (n=40) | Total No. of Fish Observed | Avg No. Fish per Census | % Freq (n=42) | | | |
| striped parrotfish blackbar | <i>Scarus iserti (S. croicensis)</i> | Scaridae | 19 | 0.5 | 22.5 | 73 | 1.7 | 40.5 | 92 | 0.3 | |
| soldierfish yellowtail | <i>Myripristis jacobus</i> | Holocentridae | 45 | 1.1 | 35 | 43 | 1.0 | 33.3 | 88 | 0.2 | |
| snapper mahogany | <i>Ocyurus chrysurus</i> | Lutjanidae | 49 | 1.2 | 32.5 | 23 | 0.5 | 16.7 | 72 | 0.2 | |
| snapper | <i>Lutjanus mahogoni</i> | Lutjanidae | 23 | 0.6 | 12.5 | 27 | 0.6 | 23.8 | 50 | 0.1 | |
| fairy basslet | <i>Gramma loreto</i> | Grammatidae | 11 | 0.3 | 10 | 37 | 0.9 | 16.7 | 48 | 0.1 | |
| Subtotal = | | | 16,585 | | | Subtotal = | 18,098 | | Subtotal = | 34,683 | 98.2 |

* percent of all observations (n = 35,322)

Appendix I. Distribution of observations on 85 fish species across seven St. Croix reef sites.

| Common Name | Species | Family | Lang Bank | | Jacks | | East Wall | | West Wall | | Eagle Ray | | Cane Bay | | Sprat Hole | |
|-------------------------|--|----------------|-----------|------|--------|------|-----------|------|-----------|------|-----------|------|----------|------|------------|------|
| | | | Spring | Fall | Spring | Fall | Spring | Fall | Spring | Fall | Spring | Fall | Spring | Fall | Spring | Fall |
| ocean surgeonfish | <i>Acanthurus bahianus</i> | Acanthuridae | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| doctorfish | <i>Acanthurus chirurgus</i> | Acanthuridae | 1 | 1 | 1 | 1 | 1 | | 1 | | 1 | 1 | 1 | 1 | | |
| blue tang | <i>Acanthurus coeruleus</i> | Acanthuridae | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| trumpetfish | <i>Aulostomus maculatus</i> | Aulostomidae | | | | | 1 | 1 | | | 1 | 1 | 1 | 1 | 1 | 1 |
| queen triggerfish | <i>Balistes vetula</i> | Balistidae | 1 | 1 | | | | | | | | | | | | |
| black durgon | <i>Melichthys niger</i> | Balistidae | 1 | 1 | | 1 | 1 | 1 | 1 | 1 | 1 | | 1 | 1 | | |
| peacock flounder | <i>Bothus lunatus</i> | Bothidae | | | | | | | | 1 | | | | | | |
| blue runner | <i>Caranx crysos</i> (C. <i>fuscus</i>) | Carangidae | | | 1 | | | | | | | | | | | 1 |
| bar jack | <i>Caranx ruber</i> | Carangidae | | | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| longsnout butterflyfish | <i>Chaetodon aculeatus</i> | Chaetodontidae | | | | | 1 | 1 | 1 | 1 | | | | | | |
| four-eye butterflyfish | <i>Chaetodon capistratus</i> | Chaetodontidae | | | | | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| spotfin butterflyfish | <i>Chaetodon ocellatus</i> | Chaetodontidae | | | | | 1 | 1 | | | | | | | | |
| banded butterflyfish | <i>Chaetodon striatus</i> | Chaetodontidae | 1 | 1 | 1 | 1 | 1 | 1 | 1 | | 1 | | | | | |
| yellowfin mojarra | <i>Gerres cinereus</i> | Gerreidae | | | | | | | | 1 | | | | | | 1 |
| fairy basslet | <i>Gramma loreto</i> | Grammatidae | | | | | | | | | | | 1 | 1 | 1 | 1 |
| tomtate | <i>Haemulon aurolineatum</i> | Haemulidae | | | 1 | 1 | | | | | | | | | | |
| caesar grunt | <i>Haemulon carbonarium</i> | Haemulidae | | 1 | 1 | 1 | | | | | 1 | | 1 | 1 | 1 | |
| smallmouth grunt | <i>Haemulon chrysargyreum</i> | Haemulidae | | | 1 | | | | | | | | | 1 | | |
| french grunt | <i>Haemulon flavolineatum</i> | Haemulidae | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| white grunt | <i>Haemulon plumieri</i> | Haemulidae | | | | | 1 | 1 | 1 | 1 | 1 | | 1 | | | |

Appendix 1 (continued). Distribution of observations on 85 fish species across seven St. Croix reef sites.

| Common Name | Species | Family | Lang Bank | | Jacks | | East Wall | | West Wall | | Eagle Ray | | Cane Bay | | Sprat Hole | |
|-------------------------------|----------------------------------|---------------|-----------|------|--------|------|-----------|------|-----------|------|-----------|------|----------|------|------------|------|
| | | | Spring | Fall | Spring | Fall | Spring | Fall | Spring | Fall | Spring | Fall | Spring | Fall | Spring | Fall |
| bluestriped grunt | <i>Haemulon sciurus</i> | Haemulidae | 1 | | | 1 | 1 | | | | | | 1 | 1 | | |
| squirrelfish | <i>Holocentrus adensionis</i> | Holocentridae | | 1 | | 1 | | 1 | 1 | | 1 | | 1 | 1 | 1 | |
| longspine squirrelfish | <i>Holocentrus rufus</i> | Holocentridae | 1 | 1 | 1 | 1 | 1 | | 1 | 1 | 1 | 1 | | 1 | | |
| blackbar soldierfish | <i>Myripristis jacobus</i> | Holocentridae | | | 1 | 1 | | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| longjaw squirrelfish | <i>Neoniphon marianus</i> | Holocentridae | | | | | | | | | | | | 1 | | |
| spanish hogfish | <i>Bodianus rufus</i> | Labridae | | 1 | 1 | 1 | 1 | | | | 1 | 1 | 1 | 1 | 1 | 1 |
| creole wrasse | <i>Clepticus parrae</i> | Labridae | | | | 1 | | | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| slippery dick | <i>Halichoeres bivittatus</i> | Labridae | 1 | | | | | | 1 | | | | | | | |
| yellowcheek wrasse | <i>Halichoeres cyanocephalus</i> | Labridae | | 1 | | | | | | | | 1 | | | | |
| yellowhead wrasse | <i>Halichoeres garnoti</i> | Labridae | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| clown wrasse | <i>Halichoeres maculipinna</i> | Labridae | 1 | 1 | | 1 | | | | | | | | | | 1 |
| puddingwife | <i>Halichoeres radiatus</i> | Labridae | | | 1 | | | | | | | | | | | |
| | <i>Thalassoma bifasciatum</i> | Labridae | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| bluehead wrasse | <i>Lutjanus analis</i> | Lutjanidae | | | | | | | | | | 1 | | | | 1 |
| mutton snapper | <i>Lutjanus apodus</i> | Lutjanidae | | | 1 | 1 | | 1 | 1 | 1 | 1 | 1 | | 1 | 1 | 1 |
| schoolmaster mahogany snapper | <i>Lutjanus mahogoni</i> | Lutjanidae | | | 1 | 1 | | | | | | | | 1 | 1 | 1 |
| yellowtail snapper | <i>Ocyurus chrysurus</i> | Lutjanidae | | | | | 1 | 1 | 1 | 1 | 1 | 1 | | | | |
| | <i>Malacanthus plumieri</i> | Malacanthidae | | | 1 | 1 | | | | | | | | | | 1 |
| sand tilefish | <i>Ahuterus scripta</i> | Monacanthidae | | | 1 | | | | | 1 | | | | | | |
| scrawled filefish | <i>Cantherhines pullus</i> | Monacanthidae | | 1 | | | 1 | 1 | | 1 | | | 1 | 1 | | |

Appendix 1 (continued). Distribution of observations on 85 fish species across seven St. Croix reef sites.

| Common Name | Species | Family | Lang Bank | | Jacks | | East Wall | | West Wall | | Eagle Ray | | Cane Bay | | Sprat Hole | |
|-----------------------|--------------------------------------|---------------|-----------|------|--------|------|-----------|------|-----------|------|-----------|------|----------|------|------------|------|
| | | | Spring | Fall | Spring | Fall | Spring | Fall | Spring | Fall | Spring | Fall | Spring | Fall | Spring | Fall |
| yellow goatfish | <i>Mulloidichthys martinicus</i> | Mullidae | | | | 1 | 1 | 1 | | 1 | 1 | | 1 | 1 | 1 | 1 |
| spotted goatfish | <i>Pseudupeneus maculatus</i> | Mullidae | | | | | 1 | | 1 | 1 | | | | | | |
| spotted moray | <i>Gymnothorax moringa</i> | Muraenidae | | | | | | | | | | | | | | 1 |
| honeycomb cowfish | <i>Acanthostracion ploygonia</i> | Ostraciidae | | | 1 | 1 | | 1 | | | 1 | | | | | 1 |
| scrawled cowfish | <i>Acanthostracion quadricornis</i> | Ostraciidae | 1 | | | | | | | | | | | | | |
| spotted trunkfish | <i>Lactophrys bicaudalis</i> | Ostraciidae | | | | | | | | | 1 | | | 1 | | |
| smooth trunkfish | <i>Lactophrys triqueter</i> | Ostraciidae | | | | | 1 | | | | 1 | | | | | 1 |
| queen angelfish | <i>Holacanthus ciliaris</i> | Pomacanthidae | 1 | | | | | | | | | | | | | |
| rock beauty | <i>Holacanthus tricolor</i> | Pomacanthidae | 1 | 1 | 1 | 1 | | 1 | 1 | 1 | | 1 | | | 1 | 1 |
| French angelfish | <i>Pomacanthus paru</i> | Pomacanthidae | | | | | 1 | | | | | | | | | |
| sergeant major | <i>Abudefduf saxatilis</i> | Pomacentridae | | | 1 | | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | |
| blue chromis | <i>Chromis cyanea</i> | Pomacentridae | 1 | 1 | 1 | 1 | 1 | 1 | 1 | | 1 | 1 | 1 | 1 | 1 | 1 |
| brown chromis | <i>Chromis multilineata</i> | Pomacentridae | | | 1 | 1 | | | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| yellowtail damselfish | <i>Microspathodon chrysurus</i> | Pomacentridae | | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| dusky damselfish | <i>Stegastes adustus</i> (S. fuscus) | Pomacentridae | | | 1 | 1 | | | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| beaugregory | <i>Stegastes leucostictus</i> | Pomacentridae | | | 1 | | | | 1 | | 1 | 1 | | 1 | | |
| bicolor damselfish | <i>Stegastes partitus</i> | Pomacentridae | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |

Appendix 1 (continued). Distribution of observations on 85 fish species across seven St. Croix reef sites.

| Common Name | Species | Family | Lang Bank | | Jacks | | East Wall | | West Wall | | Eagle Ray | | Cane Bay | | Sprat Hole | |
|------------------------|--|---------------|-----------|------|--------|------|-----------|------|-----------|------|-----------|------|----------|------|------------|------|
| | | | Spring | Fall | Spring | Fall | Spring | Fall | Spring | Fall | Spring | Fall | Spring | Fall | Spring | Fall |
| threespot damselfish | <i>Stegastes planifrons</i> | Pomacentridae | | | | | | | | | | | 1 | 1 | 1 | 1 |
| cocoa damselfish | <i>Stegastes variabilis</i> | Pomacentridae | | | | | | | | 1 | 1 | | | | 1 | |
| striped parrotfish | <i>Scarus iserti</i> (S. <i>croicensis</i>) | Scaridae | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | | 1 | 1 | 1 |
| princess parrotfish | <i>Scarus taeniopterus</i> | Scaridae | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| queen parrotfish | <i>Scarus vetula</i> | Scaridae | | | 1 | | | | 1 | 1 | 1 | | 1 | | 1 | 1 |
| greenblotch parrotfish | <i>Sparisoma atomarium</i> | Scaridae | | | | | | 1 | | | | | | | | |
| redband parrotfish | <i>Sparisoma aurofrenatum</i> | Scaridae | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| redtail parrotfish | <i>Sparisoma chrysopteron</i> | Scaridae | | 1 | 1 | 1 | | | | 1 | 1 | | | | | |
| yellowtail parrotfish | <i>Sparisoma rubripinne</i> | Scaridae | | | | | | | | 1 | | | | | | |
| stoplight parrotfish | <i>Sparisoma viride</i> | Scaridae | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| highhat | <i>Pareques acuminatus</i> | Sciaenidae | | | | 1 | | | | | | | | | | |
| cero mackerel | <i>Scomberomorus regalis</i> | Scombridae | | | | | | | | 1 | | | | | | |
| spotted scorpionfish | <i>Scorpaena plumieri</i> | Scorpaenidae | | | 1 | | | | | | | | | | | |
| graysby | <i>Cephalopholis cruentatus</i> | Serranidae | | | | | 1 | 1 | | | 1 | 1 | 1 | 1 | 1 | 1 |
| coney | <i>Cephalopholis fulvus</i> | Serranidae | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| red hind | <i>Epinephelus guttatus</i> | Serranidae | | 1 | 1 | | 1 | 1 | | | | 1 | | | | |
| yellowtail hamlet | <i>Hypoplectrus chlorurus</i> | Serranidae | | | | | | | 1 | | 1 | | | | | 1 |
| shy hamlet | <i>Hypoplectrus guttavarius</i> | Serranidae | | | | | 1 | | | 1 | | | | | 1 | |
| black hamlet | <i>Hypoplectrus nigricans</i> | Serranidae | | | | | 1 | | | | 1 | | | 1 | | 1 |

Appendix 1 (continued). Distribution of observations on 85 fish species across seven St. Croix reef sites.

| Common Name | Species | Family | Lang Bank | | Jacks | | East Wall | | West Wall | | Eagle Ray | | Cane Bay | | Sprat Hole | |
|------------------|------------------------------|----------------|-----------|------|--------|------|-----------|------|-----------|------|-----------|------|----------|------|------------|------|
| | | | Spring | Fall | Spring | Fall | Spring | Fall | Spring | Fall | Spring | Fall | Spring | Fall | Spring | Fall |
| barred hamlet | <i>Hypoplectrus puella</i> | Serranidae | | | | | | | | 1 | | | | | 1 | 1 |
| butter hamlet | <i>Hypoplectrus unicolor</i> | Serranidae | | | | | 1 | | | | 1 | | | | | |
| greater soapfish | <i>Rypticus saponaceus</i> | Serranidae | | | | | | | | 1 | | | | | | |
| tobaccofish | <i>Serranus tabacarius</i> | Serranidae | | | | | | | | 1 | | | | | | |
| harlequin bass | <i>Serranus tigrinus</i> | Serranidae | 1 | 1 | | | 1 | 1 | 1 | 1 | 1 | 1 | | 1 | 1 | 1 |
| jolthead porgy | <i>Calamus bajonado</i> | Sparidae | | | | | | | | | 1 | | | | | |
| sand diver | <i>Synodus intermedius</i> | Synodontidae | | | | | | | 1 | | | | | | | 1 |
| sharpnose puffer | <i>Canthigaster rostrata</i> | Tetraodontidae | | | 1 | | | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| porcupinefish | <i>Diodon hystrix</i> | Tetraodontidae | | | | | | | | | | | | | | 1 |
| Grand Total = | 85 species | Subtotal = | 24 | 28 | 39 | 37 | 37 | 38 | 34 | 42 | 40 | 42 | 32 | 42 | 40 | 40 |

Appendix 2A. Fish census data from Lang Bank

| Common Name | Species | Family | Spring '02 Census | | | | | | | Fall '02 Census | | | | | | |
|-----------------------------------|----------------------------------|----------------|-------------------|-------------|------|-------|------|-----|-----|-----------------|-------------|------|-------|------|-----|-----|
| | | | % Freq | No. of Fish | | | Size | | | % Freq | No. of Fish | | | Size | | |
| | | | (n=4) | Total | Avg. | StDev | Avg | Min | Max | (n=6) | Total | Avg. | StDev | Avg | Min | Max |
| ocean surgeonfish | <i>Acanthurus bahianus</i> | Acanthuridae | 100.0 | 30 | 7.5 | 2.6 | 14.5 | 11 | 17 | 100.0 | 64 | 10.7 | 2.8 | 14.8 | 10 | 20 |
| doctorfish | <i>Acanthurus chirurgus</i> | Acanthuridae | 50.0 | 2 | 0.5 | 0.6 | 18.0 | 17 | 19 | 33.3 | 7 | 1.2 | 2.0 | 20.5 | 17 | 23 |
| blue tang queen | <i>Acanthurus coeruleus</i> | Acanthuridae | 100.0 | 11 | 2.8 | 1.7 | 16.3 | 14 | 19 | 100.0 | 24 | 4.0 | 1.5 | 14.7 | 10 | 19 |
| triggerfish | <i>Balistes vetula</i> | Balistidae | 25.0 | 1 | 0.3 | 0.5 | 27.0 | 27 | 27 | 16.7 | 1 | 0.2 | 0.4 | 20.0 | 20 | 20 |
| black durgon banded butterflyfish | <i>Melichthys niger</i> | Balistidae | 100.0 | 15 | 3.8 | 4.2 | 22.0 | 20 | 26 | 83.3 | 58 | 9.7 | 11.1 | 21.0 | 15 | 30 |
| caesar grunt | <i>Chaetodon striatus</i> | Chaetodontidae | 25.0 | 2 | 0.5 | 1.0 | 10.5 | 10 | 11 | 33.3 | 4 | 0.7 | 1.0 | 12.3 | 11 | 14 |
| french grunt | <i>Haemulon carbonarium</i> | Haemulidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 20.0 | 20 | 20 |
| bluestriped grunt | <i>Haemulon flavolineatum</i> | Haemulidae | 100.0 | 9 | 2.3 | 1.3 | 15.8 | 15 | 18 | 66.7 | 13 | 2.2 | 2.2 | 16.5 | 14 | 20 |
| squirrelfish longspine | <i>Haemulon sciurus</i> | Haemulidae | 25.0 | 1 | 0.3 | 0.5 | 26.0 | 26 | 26 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| squirrelfish spanish hogfish | <i>Holocentrus acensionis</i> | Holocentridae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 50.0 | 6 | 1.0 | 1.3 | 15.8 | 10 | 20 |
| slippery dick yellowcheek wrasse | <i>Holocentrus rufus</i> | Holocentridae | 100.0 | 5 | 1.3 | 0.5 | 17.5 | 17 | 20 | 50.0 | 5 | 0.8 | 1.2 | 16.5 | 16 | 17 |
| wrasse yellowhead wrasse | <i>Bodianus rufus</i> | Labridae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 50.0 | 3 | 0.5 | 0.5 | 14.3 | 9 | 17 |
| clown wrasse | <i>Halichoeres bivittatus</i> | Labridae | 50.0 | 8 | 2.0 | 2.4 | 6.0 | 5 | 9 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| | <i>Halichoeres cyanocephalus</i> | Labridae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 33.3 | 2 | 0.3 | 0.5 | 18.5 | 12 | 25 |
| | <i>Halichoeres garnoti</i> | Labridae | 100.0 | 11 | 2.8 | 1.0 | 10.5 | 8 | 16 | 83.3 | 25 | 4.2 | 3.1 | 9.1 | 4 | 18 |
| | <i>Halichoeres maculipinna</i> | Labridae | 25.0 | 2 | 0.5 | 1.0 | 9.5 | 8 | 11 | 33.3 | 15 | 2.5 | 4.0 | 5.5 | 3 | 9 |

Appendix 2A (continued). Fish census data from Lang Bank

| Common Name | Species | Family | Spring '02 Census | | | | | | | Fall '02 Census | | | | | | |
|------------------------|---|---------------|-------------------|-------------|-------|-------|------|-----|-----|-----------------|-------------|-------|-------|------|-----|-----|
| | | | % Freq (n=4) | No. of Fish | | | Size | | | %Freq (n=6) | No. of Fish | | | Size | | |
| | | | | Total | Avg. | StDev | Avg | Min | Max | | Total | Avg. | StDev | Avg | Min | Max |
| bluehead wrasse | <i>Thalassoma bifasciatum</i> | Labridae | 100.0 | 1340 | 335.0 | 102.5 | 3.0 | 2 | 9 | 100.0 | 1405 | 234.2 | 86.2 | 4.3 | 2 | 10 |
| orangespotted filefish | <i>Cantherhines pullus</i> | Monacanthidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 33.3 | 3 | 0.5 | 0.8 | 21.8 | 17 | 25 |
| scrawled cowfish | <i>Acanthostracion quadricornis</i> | Ostraciidae | 25.0 | 1 | 0.3 | 0.5 | 22.0 | 22 | 22 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| queen angelfish | <i>Holacanthus ciliaris</i> | Pomacanthidae | 25.0 | 1 | 0.3 | 0.5 | 25.0 | 25 | 25 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| rock beauty | <i>Holacanthus tricolor</i> | Pomacanthidae | 75.0 | 3 | 0.8 | 0.5 | 17.7 | 15 | 22 | 33.3 | 4 | 0.7 | 1.0 | 9.0 | 3 | 20 |
| blue chromis | <i>Chromis cyanea</i> | Pomacentridae | 75.0 | 7 | 1.8 | 1.5 | 4.0 | 3 | 7 | 83.3 | 45 | 7.5 | 4.4 | 4.8 | 2 | 8 |
| yellowtail damselfish | <i>Microspathodon chrysurus</i> | Pomacentridae | 0 | 0 | 0 | 0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 12.0 | 12 | 12 |
| bicolor damselfish | <i>Stegastes partitus</i> | Pomacentridae | 100.0 | 171 | 42.8 | 18.6 | 4.0 | 2 | 6 | 100.0 | 721 | 120.2 | 25.6 | 3.8 | 2 | 6 |
| striped parrotfish | <i>Scarus iserti</i> (<i>S. croicensis</i>) | Scaridae | 25.0 | 1 | 0.3 | 0.5 | 24.0 | 24 | 24 | 16.7 | 1 | 0.2 | 0.4 | 30.0 | 30 | 30 |
| princess parrotfish | <i>Scarus taeniopterus</i> | Scaridae | 100.0 | 11 | 2.8 | 1.3 | 16.8 | 12 | 21 | 100.0 | 28 | 4.7 | 1.9 | 17.2 | 10 | 30 |
| redband parrotfish | <i>Sparisoma aurofrenatum</i> | Scaridae | 75.0 | 7 | 1.8 | 1.3 | 14.7 | 12 | 18 | 83.3 | 16 | 2.7 | 1.8 | 17.0 | 10 | 20 |
| redtail parrotfish | <i>Sparisoma chrysopteron</i> | Scaridae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 22.0 | 22 | 22 |
| stoplight parrotfish | <i>Sparisoma viride</i> | Scaridae | 50.0 | 3 | 0.8 | 1.0 | 27.8 | 25 | 30 | 50.0 | 7 | 1.2 | 1.5 | 30.7 | 25 | 40 |
| coney | <i>Cephalopholis fulvus</i> | Serranidae | 100.0 | 14 | 3.5 | 2.4 | 16.6 | 14 | 20 | 100.0 | 29 | 4.8 | 2.4 | 15.2 | 10 | 22 |
| red hind | <i>Epinephelus guttatus</i> | Serranidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 25.0 | 25 | 25 |
| harlequin bass | <i>Serranus tigrinus</i> | Serranidae | 50.0 | 3 | 0.8 | 1.0 | 8.3 | 6 | 10 | 16.7 | 1 | 0.2 | 0.4 | 7.0 | 7 | 7 |
| Total= | | | | 1659 | | | | | | Total= | 2491 | | | | | |

Appendix 2B. Fish census data from Jacks & Isaacs Bay

| Common Name | Species | Family | Spring '02 Census | | | | | | Fall '02 Census | | | | | | | |
|------------------------|-------------------------------|----------------|-------------------|-------------|------|-------|------|-----|-----------------|--------------|-------------|------|-------|------|-----|-----|
| | | | % Freq (n=6) | No. of Fish | | | Size | | | % Freq (n=6) | No. of Fish | | | Size | | |
| | | | | Total | Avg. | StDev | Avg | Min | Max | | Total | Avg. | StDev | Avg | Min | Max |
| ocean surgeonfish | <i>Acanthurus bahianus</i> | Acanthuridae | 100.0 | 74 | 12.3 | 5.5 | 15.0 | 7 | 20 | 100.0 | 32 | 5.3 | 1.2 | 13.7 | 6 | 20 |
| doctorfish | <i>Acanthurus chirurgus</i> | Acanthuridae | 50.0 | 5 | 0.8 | 1.2 | 19.0 | 17 | 21 | 16.7 | 1 | 0.2 | 0.4 | 18.0 | 18 | 18 |
| blue tang | <i>Acanthurus coeruleus</i> | Acanthuridae | 100.0 | 12 | 2.0 | 0.9 | 14.7 | 8 | 20 | 100.0 | 17 | 2.8 | 1.0 | 11.2 | 3 | 17 |
| black durgon | <i>Melichthys niger</i> | Balistidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 33.3 | 2 | 0.3 | 0.5 | 17.5 | 15 | 20 |
| blue runner | <i>Caranx crysos</i> | Carangidae | 16.7 | 1 | 0.2 | 0.4 | 30.0 | 30 | 30 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| bar jack | <i>Caranx ruber</i> | Carangidae | 16.7 | 2 | 0.3 | 0.8 | 22.5 | 21 | 24 | 16.7 | 2 | 0.3 | 0.8 | 20.0 | 20 | 20 |
| banded butterflyfish | <i>Chaetodon striatus</i> | Chaetodontidae | 16.7 | 4 | 0.7 | 1.6 | 11.0 | 10 | 12 | 16.7 | 2 | 0.3 | 0.8 | 9.0 | 8 | 10 |
| tomtate | <i>Haemulon aurolineatum</i> | Haemulidae | 16.7 | 1 | 0.2 | 0.4 | 19.0 | 19 | 19 | 16.7 | 2 | 0.3 | 0.8 | 14.5 | 14 | 15 |
| caesar grunt | <i>Haemulon carbonarium</i> | Haemulidae | 16.7 | 1 | 0.2 | 0.4 | 19.0 | 19 | 19 | 16.7 | 1 | 0.2 | 0.4 | 16.0 | 16 | 16 |
| smallmouth grunt | <i>Haemulon chrysargyreum</i> | Haemulidae | 16.7 | 3 | 0.5 | 1.2 | 12.0 | 12 | 12 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| french grunt | <i>Haemulon flavolineatum</i> | Haemulidae | 83.3 | 19 | 3.2 | 2.9 | 18.0 | 14 | 23 | 66.7 | 9 | 1.5 | 1.6 | 15.5 | 13 | 16 |
| bluestriped grunt | <i>Haemulon sciurus</i> | Haemulidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 2 | 0.3 | 0.8 | 19.0 | 18 | 20 |
| squirrelfish | <i>Holocentrus acensionis</i> | Holocentridae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 33.3 | 3 | 0.5 | 0.8 | 19.3 | 19 | 20 |
| longspine squirrelfish | <i>Holocentrus rufus</i> | Holocentridae | 83.3 | 7 | 1.2 | 0.8 | 19.6 | 18 | 22 | 33.3 | 2 | 0.3 | 0.5 | 16.5 | 16 | 17 |
| blackbar soldierfish | <i>Myripristis jacobus</i> | Holocentridae | 66.7 | 7 | 1.2 | 1.2 | 16.4 | 16 | 18 | 33.3 | 6 | 1.0 | 2.0 | 13.5 | 12 | 15 |
| spanish hogfish | <i>Bodianus rufus</i> | Labridae | 33.3 | 2 | 0.3 | 0.5 | 18.5 | 17 | 20 | 16.7 | 1 | 0.2 | 0.4 | 6.0 | 6 | 6 |
| creole wrasse | <i>Clepticus parrae</i> | Labridae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 33.3 | 13 | 2.2 | 4.8 | 10.0 | 4 | 15 |

Appendix 2B (continued). Fish census data from Jacks & Isaacs Bay

| Common Name | Species | Family | Spring '02 Census | | | | | | | Fall '02 Census | | | | | | |
|-------------------------------|----------------------------------|---------------|-------------------|-------------|-------|-------|------|-----|-----|-----------------|-------------|-------|-------|------|-----|-----|
| | | | % Freq (n=6) | No. of Fish | | | Size | | | % Freq (n=6) | No. of Fish | | | Size | | |
| | | | | Total | Avg. | StDev | Avg | Min | Max | | Total | Avg. | StDev | Avg | Min | Max |
| yellowhead wrasse | <i>Halichoeres garnoti</i> | Labridae | 83.3 | 13 | 2.2 | 1.3 | 9.8 | 6 | 17 | 83.3 | 16 | 2.7 | 2.6 | 9.0 | 4 | 15 |
| clown wrasse | <i>Halichoeres maculipinna</i> | Labridae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 20 | 3.3 | 8.2 | 3.5 | 3 | 5 |
| puddingwife bluehead wrasse | <i>Halichoeres radiatus</i> | Labridae | 16.7 | 1 | 0.2 | 0.4 | 8.0 | 8 | 8 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| schoolmaster mahogany snapper | <i>Thalassoma bifasciatum</i> | Labridae | 100.0 | 995 | 165.8 | 78.3 | 5.3 | 2 | 12 | 100.0 | 640 | 106.7 | 59.6 | 4.0 | 2 | 10 |
| | <i>Lutjanus apodus</i> | Lutjanidae | 16.7 | 1 | 0.2 | 0.4 | 17.0 | 17 | 17 | 16.7 | 1 | 0.2 | 0.4 | 20.0 | 20 | 20 |
| | <i>Lutjanus mahogoni</i> | Lutjanidae | 16.7 | 3 | 0.5 | 1.2 | 22.0 | 20 | 22 | 33.3 | 3 | 0.5 | 0.8 | 18.0 | 15 | 20 |
| sand tilefish | <i>Malacanthus plumieri</i> | Malacanthidae | 50.0 | 3 | 0.5 | 0.5 | 27.3 | 26 | 30 | 16.7 | 1 | 0.2 | 0.4 | 25.0 | 25 | 25 |
| scrawled filefish | <i>Aluterus scripta</i> | Monacanthidae | 16.7 | 1 | 0.2 | 0.4 | 25.0 | 25 | 25 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| yellow goatfish | <i>Mulloidichthys martinicus</i> | Mullidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 5 | 0.8 | 2.0 | 18.0 | 16 | 20 |
| honeycomb cowfish | <i>Acanthostracion ploygonia</i> | Ostraciidae | 33.3 | 2 | 0.3 | 0.5 | 24.0 | 22 | 26 | 16.7 | 1 | 0.2 | 0.4 | 20.0 | 20 | 20 |
| smooth trunkfish | <i>Lactophrys triqueter</i> | Ostraciidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 33.3 | 3 | 0.5 | 0.8 | 17.0 | 15 | 20 |
| rock beauty sergeant major | <i>Holacanthus tricolor</i> | Pomacanthidae | 33.3 | 2 | 0.3 | 0.5 | 11.5 | 8 | 15 | 16.7 | 1 | 0.2 | 0.4 | 18.0 | 18 | 18 |
| blue chromis | <i>Abudefduf saxatilis</i> | Pomacentridae | 16.7 | 1 | 0.2 | 0.4 | 13.0 | 13 | 13 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| brown chromis | <i>Chromis cyanea</i> | Pomacentridae | 100.0 | 197 | 32.8 | 32.0 | 5.3 | 2 | 8 | 100.0 | 139 | 23.2 | 19.3 | 4.3 | 2 | 7 |
| yellowtail damselfish | <i>Chromis multilineata</i> | Pomacentridae | 16.7 | 8 | 1.3 | 3.3 | 5.5 | 5 | 6 | 16.7 | 2 | 0.3 | 0.8 | 5.5 | 5 | 6 |
| dusky damselfish | <i>Microspathodon chrysurus</i> | Pomacentridae | 50.0 | 6 | 1.0 | 1.1 | 12.2 | 11 | 13 | 66.7 | 6 | 1.0 | 0.9 | 12.3 | 12 | 13 |
| beaugregory | <i>Stegastes adustus</i> | Pomacentridae | 33.3 | 3 | 0.5 | 0.8 | 7.8 | 6 | 10 | 16.7 | 1 | 0.2 | 0.4 | 8.0 | 8 | 8 |
| | <i>(S. fuscus)</i> | Pomacentridae | 33.3 | 3 | 0.5 | 0.8 | 7.8 | 6 | 10 | 16.7 | 1 | 0.2 | 0.4 | 8.0 | 8 | 8 |
| | <i>Stegastes leucostictus</i> | Pomacentridae | 33.3 | 3 | 0.5 | 0.8 | 7.0 | 6 | 8 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |

Appendix 2B (continued). Fish census data from Jacks & Isaacs Bay

| Common Name | Species | Family | Spring '02 Census | | | | | | | Fall '02 Census | | | | | | |
|------------------------------|---|----------------|-------------------|-------------|------|-------|------|-----|-----|-----------------|-------------|------|-------|------|-----|-----|
| | | | % Freq (n=6) | No. of Fish | | | Size | | | % Freq (n=6) | No. of Fish | | | Size | | |
| | | | | Total | Avg. | StDev | Avg | Min | Max | | Total | Avg. | StDev | Avg | Min | Max |
| bicolor damselfish | <i>Stegastes partitus</i> | Pomacentridae | 100.0 | 446 | 74.3 | 22.5 | 5.1 | 3 | 7 | 100.0 | 400 | 66.7 | 31.1 | 3.8 | 2 | 6 |
| striped parrotfish | <i>Scarus iserti</i> (<i>S. croicensis</i>) | Scaridae | 16.7 | 2 | 0.3 | 0.8 | 13.0 | 12 | 14 | 16.7 | 3 | 0.5 | 1.2 | 10.0 | 6 | 18 |
| princess parrotfish | <i>Scarus taeniopterus</i> | Scaridae | 83.3 | 17 | 2.8 | 2.8 | 15.4 | 8 | 22 | 66.7 | 5 | 0.8 | 0.8 | 21.5 | 16 | 30 |
| queen parrotfish | <i>Scarus vetula</i> | Scaridae | 16.7 | 2 | 0.3 | 0.8 | 21.0 | 20 | 22 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| redband parrotfish | <i>Sparisoma aurofrenatum</i> | Scaridae | 100.0 | 24 | 4.0 | 1.3 | 14.2 | 5 | 25 | 100.0 | 17 | 2.8 | 1.6 | 16.8 | 10 | 22 |
| redtail parrotfish | <i>Sparisoma chrysopteron</i> | Scaridae | 16.7 | 12 | 2.0 | 4.9 | 23.0 | 18 | 24 | 16.7 | 1 | 0.2 | 0.4 | 22.0 | 22 | 22 |
| stoplight parrotfish | <i>Sparisoma viride</i> | Scaridae | 66.7 | 6 | 1.0 | 0.9 | 19.5 | 15 | 24 | 50.0 | 4 | 0.7 | 0.8 | 22.7 | 16 | 30 |
| highhat spotted scorpionfish | <i>Pareques acuminatus</i> | Sciaenidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 2 | 0.3 | 0.8 | 11.5 | 8 | 15 |
| scorpionfish | <i>Scorpaena plumieri</i> | Scorpaenidae | 16.7 | 1 | 0.2 | 0.4 | 27.0 | 27 | 27 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| coney | <i>Cephalopholis fulvus</i> | Serranidae | 100.0 | 32 | 5.3 | 2.7 | 18.1 | 12 | 25 | 100.0 | 19 | 3.2 | 1.6 | 16.6 | 10 | 25 |
| red hind | <i>Epinephelus guttatus</i> | Serranidae | 16.7 | 1 | 0.2 | 0.4 | 22.0 | 22 | 22 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| sharpnose puffer | <i>Canthigaster rostrata</i> | Tetraodontidae | 16.7 | 1 | 0.2 | 0.4 | 5.0 | 5 | 5 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| Total = | | | | 1921 | | | | | | | Total = | 1385 | | | | |

Appendix 2C. Fish census data from East Wall, Salt River Canyon

| Common Name | Species | Family | Spring '02 Census | | | | | | Fall '02 Census | | | | | | | |
|-------------------------|-------------------------------|----------------|-------------------|-------------|------|-------|------|-----|-----------------|--------------|-------------|------|-------|------|-----|-----|
| | | | % Freq (n=6) | No. of Fish | | | Size | | | % Freq (n=6) | No. of Fish | | | Size | | |
| | | | | Total | Avg. | StDev | Avg | Min | Max | | Total | Avg. | StDev | Avg | Min | Max |
| ocean surgeonfish | <i>Acanthurus bahianus</i> | Acanthuridae | 100.0 | 73 | 12.2 | 4.4 | 14.5 | 12 | 18 | 100.0 | 62 | 10.3 | 2.8 | 13.3 | 8 | 18 |
| doctorfish | <i>Acanthurus chirurgus</i> | Acanthuridae | 33.3 | 3 | 0.5 | 0.8 | 20.3 | 19 | 23 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| blue tang | <i>Acanthurus coeruleus</i> | Acanthuridae | 83.3 | 18 | 3.0 | 0.9 | 15.4 | 14 | 18 | 100.0 | 20 | 3.3 | 1.6 | 15.0 | 10 | 20 |
| trumpetfish | <i>Aulostomus maculatus</i> | Aulostomidae | 16.7 | 2 | 0.3 | 0.8 | 27.5 | 25 | 30 | 33.3 | 3 | 0.5 | 0.8 | 42.5 | 30 | 50 |
| black durgon | <i>Melichthys niger</i> | Balistidae | 83.3 | 11 | 1.8 | 1.5 | 23.7 | 20 | 25 | 83.3 | 31 | 5.2 | 7.4 | 21.2 | 15 | 25 |
| bar jack | <i>Caranx ruber</i> | Carangidae | 33.3 | 4 | 0.7 | 1.0 | 21.0 | 14 | 20 | 16.7 | 2 | 0.3 | 0.8 | 19.0 | 18 | 20 |
| longsnout butterflyfish | <i>Chaetodon aculeatus</i> | Chaetodontidae | 16.7 | 1 | 0.2 | 0.4 | 8.0 | 8 | 8 | 16.7 | 3 | 0.5 | 1.2 | 9.0 | 8 | 10 |
| four-eye butterflyfish | <i>Chaetodon capistratus</i> | Chaetodontidae | 100.0 | 16 | 2.7 | 1.8 | 9.3 | 7 | 11 | 83.3 | 18 | 3.0 | 2.4 | 9.0 | 5 | 11 |
| spotfin butterflyfish | <i>Chaetodon ocellatus</i> | Chaetodontidae | 16.7 | 1 | 0.2 | 0.4 | 16.0 | 16 | 16 | 16.7 | 1 | 0.2 | 0.4 | 13.0 | 13 | 13 |
| banded butterflyfish | <i>Chaetodon striatus</i> | Chaetodontidae | 50.0 | 6 | 1.0 | 1.1 | 10.5 | 8 | 12 | 16.7 | 2 | 0.3 | 0.8 | 11.0 | 11 | 11 |
| french grunt | <i>Haemulon flavolineatum</i> | Haemulidae | 100.0 | 11 | 1.8 | 0.8 | 15.3 | 12 | 17 | 100.0 | 17 | 2.8 | 1.2 | 15.6 | 10 | 20 |
| white grunt | <i>Haemulon plumieri</i> | Haemulidae | 50.0 | 3 | 0.5 | 0.5 | 22.3 | 20 | 25 | 16.7 | 1 | 0.2 | 0.4 | 20.0 | 20 | 20 |
| bluestriped grunt | <i>Haemulon sciurus</i> | Haemulidae | 16.7 | 1 | 0.2 | 0.4 | 22.0 | 22 | 22 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| squirrelfish | <i>Holocentrus acensionis</i> | Holocentridae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 3 | 0.5 | 1.2 | 19.0 | 18 | 20 |
| longspine squirrelfish | <i>Holocentrus rufus</i> | Holocentridae | 50.0 | 5 | 0.8 | 1.0 | 16.2 | 15 | 17 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| blackbar soldierfish | <i>Myripristis jacobus</i> | Holocentridae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 18.0 | 18 | 18 |
| spanish hogfish | <i>Bodianus rufus</i> | Labridae | 16.7 | 2 | 0.3 | 0.8 | 11.5 | 3 | 20 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |

Appendix 2C (continued). Fish census data from East Wall, Salt River Canyon

| Common Name | Species | Family | Spring '02 Census | | | | | | | Fall '02 Census | | | | | | |
|------------------------|---|---------------|-------------------|-------------|-------|-------|------|-----|-----|-----------------|-------------|-------|-------|------|-----|-----|
| | | | %Freq (n=6) | No. of Fish | | | Size | | | %Freq (n=6) | No. of Fish | | | Size | | |
| | | | | Total | Avg. | StDev | Avg | Min | Max | | Total | Avg. | StDev | Avg | Min | Max |
| yellowhead wrasse | <i>Halichoeres garnoti</i> | Labridae | 50.0 | 8 | 1.3 | 1.5 | 9.8 | 5 | 14 | 100.0 | 22 | 3.7 | 2.6 | 8.6 | 4 | 14 |
| bluehead wrasse | <i>Thalassoma bifasciatum</i> | Labridae | 100.0 | 1290 | 215.0 | 114.8 | 3.2 | 2 | 9 | 100.0 | 820 | 136.7 | 61.5 | 4.8 | 2 | 12 |
| schoolmaster | <i>Lutjanus apodus</i> | Lutjanidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 5 | 0.8 | 2.0 | 13.0 | 10 | 15 |
| yellowtail snapper | <i>Ocyurus chrysurus</i> | Lutjanidae | 100.0 | 19 | 3.2 | 2.1 | 22.6 | 17 | 28 | 16.7 | 3 | 0.5 | 1.2 | 21.0 | 18 | 24 |
| orangespotted filefish | <i>Cantherhines pullus</i> | Monacanthidae | 16.7 | 1 | 0.2 | 0.4 | 17.0 | 17 | 17 | 16.7 | 1 | 0.2 | 0.4 | 16.0 | 16 | 16 |
| yellow goatfish | <i>Mulloidichthys martinicus</i> | Mullidae | 16.7 | 1 | 0.2 | 0.4 | 20.0 | 20 | 20 | 16.7 | 3 | 0.5 | 1.2 | 19.0 | 18 | 20 |
| spotted goatfish | <i>Psuedupeneus maculatus</i> | Mullidae | 50.0 | 3 | 0.5 | 0.5 | 18.7 | 16 | 20 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| honeycomb cowfish | <i>Acanthostracion ploygonia</i> | Ostraciidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 26.0 | 26 | 26 |
| smooth trunkfish | <i>Lactophrys triqueter</i> | Ostraciidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 66.7 | 4 | 0.7 | 0.5 | 15.8 | 15 | 17 |
| rock beauty | <i>Holacanthus tricolor</i> | Pomacanthidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 33.3 | 2 | 0.3 | 0.5 | 17.5 | 13 | 22 |
| french angelfish | <i>Pomacanthus paru</i> | Pomacanthidae | 33.3 | 2 | 0.3 | 0.5 | 24.5 | 23 | 26 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| sergeant major | <i>Abudefduf saxatilis</i> | Pomacentridae | 16.7 | 2 | 0.3 | 0.8 | 13.0 | 12 | 14 | 83.3 | 67 | 11.2 | 10.7 | 11.8 | 8 | 15 |
| blue chromis | <i>Chromis cyanea</i> | Pomacentridae | 33.3 | 71 | 11.8 | 24.0 | 3.0 | 2 | 4 | 50.0 | 65 | 10.8 | 13.9 | 4.3 | 3 | 7 |
| yellowtail damselfish | <i>Microspathodon chrysurus</i> | Pomacentridae | 83.3 | 5 | 0.8 | 0.4 | 12.4 | 12 | 13 | 66.7 | 7 | 1.2 | 1.0 | 11.6 | 10 | 13 |
| bicolor damselfish | <i>Stegastes partitus</i> | Pomacentridae | 100.0 | 143 | 23.8 | 7.8 | 4.0 | 2 | 7 | 100.0 | 211 | 35.2 | 8.1 | 4.0 | 2 | 6 |
| striped parrotfish | <i>Scarus iserti</i> (<i>S. croicensis</i>) | Scaridae | 16.7 | 2 | 0.3 | 0.8 | 8.0 | 4 | 12 | 16.7 | 6 | 1.0 | 2.4 | 5.0 | 4 | 5 |
| princess parrotfish | <i>Scarus taeniopterus</i> | Scaridae | 83.3 | 15 | 2.5 | 2.1 | 17.0 | 14 | 25 | 83.3 | 21 | 3.5 | 2.3 | 17.1 | 7 | 25 |

Appendix 2C (continued). Fish census data from East Wall, Salt River Canyon

| Common Name | Species | Family | Spring '02 Census | | | | | | | Fall '02 Census | | | | | | |
|------------------------|---------------------------------|----------------|-------------------|-------------|------|-------|------|-----|-----|-----------------|-------------|------|-------|------|-----|-----|
| | | | % Freq (n=6) | No. of Fish | | | Size | | | % Freq (n=6) | No. of Fish | | | Size | | |
| | | | | Total | Avg. | StDev | Avg | Min | Max | | Total | Avg. | StDev | Avg | Min | Max |
| greenblotch parrotfish | <i>Sparisoma atomarium</i> | Scaridae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 4 | 0.7 | 1.6 | 14.5 | 14 | 15 |
| redband parrotfish | <i>Sparisoma aurofrenatum</i> | Scaridae | 83.3 | 13 | 2.2 | 1.7 | 16.9 | 12 | 21 | 83.3 | 21 | 3.5 | 1.9 | 17.2 | 11 | 23 |
| stoplight parrotfish | <i>Sparisoma viride</i> | Scaridae | 33.3 | 2 | 0.3 | 0.5 | 22.5 | 15 | 30 | 66.7 | 6 | 1.0 | 0.9 | 19.8 | 6 | 32 |
| graysby | <i>Cephalopholis cruentatus</i> | Serranidae | 16.7 | 1 | 0.2 | 0.4 | 17.0 | 17 | 17 | 16.7 | 1 | 0.2 | 0.4 | 18.0 | 18 | 18 |
| coney | <i>Cephalopholis fulvus</i> | Serranidae | 100.0 | 15 | 2.5 | 1.6 | 16.4 | 14 | 19 | 83.3 | 15 | 2.5 | 1.9 | 17.8 | 8 | 24 |
| red hind | <i>Epinephelus guttatus</i> | Serranidae | 16.7 | 1 | 0.2 | 0.4 | 25.0 | 25 | 25 | 16.7 | 1 | 0.2 | 0.4 | 20.0 | 20 | 20 |
| shy hamlet | <i>Hypoplectrus guttavarius</i> | Serranidae | 16.7 | 1 | 0.2 | 0.4 | 13.0 | 13 | 13 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| black hamlet | <i>Hypoplectrus nigricans</i> | Serranidae | 16.7 | 1 | 0.2 | 0.4 | 14.0 | 14 | 14 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| butter hamlet | <i>Hypoplectrus unicolor</i> | Serranidae | 16.7 | 1 | 0.2 | 0.4 | 10.0 | 10 | 10 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| greater soapfish | <i>Rypticus saponaceus</i> | Serranidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 15.0 | 15 | 15 |
| tobaccofish | <i>Serranus tabacarius</i> | Serranidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 5.0 | 5 | 5 |
| harlequin bass | <i>Serranus tigrinus</i> | Serranidae | 16.7 | 1 | 0.2 | 0.4 | 11.0 | 11 | 11 | 50.0 | 3 | 0.5 | 0.5 | 7.0 | 5 | 8 |
| sharpnose puffer | <i>Canthigaster rostrata</i> | Tetraodontidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 6.0 | 6 | 6 |
| Total= | | | | 1755 | | | | | | | Total= | | | 1456 | | |

Appendix 2D. Fish census data from West Wall, Salt River Canyon

| Common Name | Species Name | Family | Spring '02 Census | | | | | | | Fall '02 Census | | | | | | |
|-------------------------|---------------------------------|----------------|-------------------|-------------|------|-------|------|-----|-----|-----------------|-------------|------|-------|------|-----|-----|
| | | | % Freq | No. of Fish | | | Size | | | % Freq | No. of Fish | | | Size | | |
| | | | (n=6) | Total | Avg. | StDev | Avg | Min | Max | (n=6) | Total | Avg. | StDev | Avg | Min | Max |
| ocean surgeonfish | <i>Acanthurus bahianus</i> | Acanthuridae | 100.0 | 96 | 16.0 | 6.3 | 13.6 | 9 | 17 | 100.0 | 49 | 8.2 | 4.3 | 14.8 | 7 | 22 |
| doctorfish | <i>Acanthurus chirurgus</i> | Acanthuridae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 33.3 | 3 | 0.5 | 0.8 | 16.5 | 12 | 21 |
| blue tang | <i>Acanthurus coeruleus</i> | Acanthuridae | 83.3 | 14 | 2.3 | 1.6 | 16.4 | 14 | 20 | 100.0 | 12 | 2.0 | 0.6 | 16.8 | 14 | 20 |
| black durgon | <i>Melichthys niger</i> | Balistidae | 66.7 | 27 | 4.5 | 6.3 | 23.1 | 20 | 26 | 66.7 | 44 | 7.3 | 8.9 | 23.0 | 12 | 25 |
| peacock flounder | <i>Bothus lunatus</i> | Bothidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 12.0 | 12 | 12 |
| bar jack | <i>Caranx ruber</i> | Carangidae | 16.7 | 1 | 0.2 | 0.4 | 22.0 | 22 | 22 | 33.3 | 3 | 0.5 | 0.8 | 29.5 | 18 | 40 |
| longsnout butterflyfish | <i>Chaetodon aculeatus</i> | Chaetodontidae | 33.3 | 3 | 0.5 | 0.8 | 6.3 | 6 | 7 | 33.3 | 3 | 0.5 | 0.8 | 8.0 | 7 | 10 |
| foureye butterflyfish | <i>Chaetodon capistratus</i> | Chaetodontidae | 100.0 | 20 | 3.3 | 2.3 | 9.5 | 4 | 11 | 83.3 | 15 | 2.5 | 1.6 | 9.2 | 7 | 12 |
| banded butterflyfish | <i>Chaetodon striatus</i> | Chaetodontidae | 50.0 | 5 | 0.8 | 1.0 | 11.5 | 11 | 12 | 0.0 | 0 | 0.00 | 0.0 | - | - | - |
| yellowfin mojarra | <i>Gerres cinereus</i> | Gerreidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 2 | 0.3 | 0.8 | 25.0 | 24 | 26 |
| french grunt | <i>Haemulon flavolineatum</i> | Haemulidae | 100.0 | 17 | 2.8 | 1.0 | 13.8 | 9 | 17 | 83.3 | 15 | 2.5 | 2.4 | 17.2 | 15 | 20 |
| white grunt | <i>Haemulon plumieri</i> | Haemulidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 22.0 | 22 | 22 |
| squirrelfish | <i>Holocentrus adscensionis</i> | Holocentridae | 33.3 | 3 | 0.5 | 0.8 | 17.3 | 15 | 20 | 16.7 | 1 | 0.2 | 0.4 | 18.0 | 18 | 18 |
| longspine squirrelfish | <i>Holocentrus rufus</i> | Holocentridae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 33.3 | 2 | 0.3 | 0.5 | 16.5 | 16 | 17 |
| blackbar soldierfish | <i>Myripristis jacobus</i> | Holocentridae | 16.7 | 1 | 0.2 | 0.4 | 14.0 | 14 | 14 | 16.7 | 1 | 0.2 | 0.4 | 15.0 | 15 | 15 |
| creole wrasse | <i>Clepticus parrae</i> | Labridae | 66.7 | 136 | 22.7 | 27.0 | 10.5 | 8 | 15 | 66.7 | 345 | 57.5 | 58.8 | 16.3 | 10 | 20 |

Appendix 2D (continued). Fish census data from West Wall, Salt River Canyon

| Common Name | Species Name | Family | Spring '02 Census | | | | | | | Fall '02 Census | | | | | | |
|------------------------|---|---------------|-------------------|-------------|-------|-------|------|-----|-----|-----------------|-------------|-------|-------|------|-----|-----|
| | | | % Freq (n=6) | No. of Fish | | | Size | | | % Freq (n=6) | No. of Fish | | | Size | | |
| | | | | Total | Avg. | StDev | Avg | Min | Max | | Total | Avg. | StDev | Avg | Min | Max |
| slippery dick | <i>Halichoeres bivittatus</i> | Labridae | 16.7 | 2 | 0.3 | 0.8 | 2.5 | 2 | 3 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| yellowhead wrasse | <i>Halichoeres garnoti</i> | Labridae | 33.3 | 2 | 0.3 | 0.5 | 10.0 | 10 | 10 | 83.3 | 14 | 2.3 | 1.2 | 9.4 | 5 | 15 |
| bluehead wrasse | <i>Thalassoma bifasciatum</i> | Labridae | 100.0 | 1333 | 222.2 | 41.8 | 3.0 | 1.5 | 10 | 100.0 | 1235 | 205.8 | 130.2 | 4.6 | 2 | 10 |
| schoolmaster | <i>Lutjanus apodus</i> | Lutjanidae | 33.3 | 3 | 0.5 | 0.8 | 22.0 | 18 | 25 | 33.3 | 2 | 0.3 | 0.5 | 30.0 | 30 | 30 |
| yellowtail snapper | <i>Ocyurus chrysurus</i> | Lutjanidae | 33.3 | 20 | 3.3 | 5.2 | 25.0 | 20 | 30 | 50.0 | 15 | 2.5 | 2.8 | 26.0 | 21 | 30 |
| scrawled filefish | <i>Ahuterus scripta</i> | Monacanthidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 35.0 | 35 | 35 |
| orangespotted filefish | <i>Cantherhines pullus</i> | Monacanthidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 50.0 | 3 | 0.5 | 0.5 | 15.7 | 15 | 16 |
| yellow goatfish | <i>Mulloidichthys martinicus</i> | Mullidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 4 | 0.7 | 1.6 | 17.0 | 16 | 19 |
| spotted goatfish | <i>Psuedupeneus maculatus</i> | Mullidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 20.0 | 20 | 20 |
| rock beauty | <i>Holacanthus tricolor</i> | Pomacanthidae | 16.7 | 1 | 0.2 | 0.4 | 17.0 | 17 | 17 | 16.7 | 1 | 0.2 | 0.4 | 6.0 | 6 | 6 |
| sergeant major | <i>Abudefduf saxatilis</i> | Pomacentridae | 50.0 | 20 | 3.3 | 5.9 | 11.3 | 10 | 14 | 50.0 | 20 | 3.3 | 4.3 | 12.8 | 10 | 14 |
| blue chromis | <i>Chromis cyanea</i> | Pomacentridae | 16.7 | 3 | 0.5 | 1.2 | 4.0 | 3 | 5 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| brown chromis | <i>Chromis multilineata</i> | Pomacentridae | 33.3 | 41 | 6.8 | 14.0 | 5.3 | 4 | 7 | 33.3 | 90 | 15.0 | 23.5 | 6.0 | 4 | 7 |
| yellowtail damselfish | <i>Microspathodon chrysurus</i> | Pomacentridae | 100.0 | 14 | 2.3 | 0.8 | 10.0 | 5 | 12 | 100.0 | 17 | 2.8 | 1.2 | 12.1 | 2.5 | 15 |
| dusky damselfish | <i>Stegastes adustus</i> (<i>S. fuscus</i>) | Pomacentridae | 100.0 | 39 | 6.5 | 10.1 | 7.9 | 3 | 12 | 83.3 | 15 | 2.5 | 1.8 | 8.9 | 7 | 11 |
| beaugregory | <i>Stegastes leucostictus</i> | Pomacentridae | 16.7 | 1 | 0.2 | 0.4 | 9.0 | 9 | 9 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |

Appendix 2D (continued). Fish census data from West Wall, Salt River Canyon

| Common Name | Species Name | Family | Spring '02 Census | | | | | | | Fall '02 Census | | | | | | | | |
|-----------------------|---|----------------|-------------------|-------------|------|-------|------|-----|-----|-----------------|-------------|---------|-------|------|-----|-----|--|--|
| | | | %Freq | No. of Fish | | | Size | | | %Freq | No. of Fish | | | Size | | | | |
| | | | (n=6) | Total | Avg. | StDev | Avg | Min | Max | (n=6) | Total | Avg. | StDev | Avg | Min | Max | | |
| bicolor damselfish | <i>Stegastes partitus</i> | Pomacentridae | 100.0 | 52 | 8.7 | 8.7 | 5.3 | 1 | 12 | 100.0 | 220 | 36.7 | 10.8 | 4.4 | 2 | 8 | | |
| striped parrotfish | <i>Scarus iserti</i> (<i>S. croicensis</i>) | Scaridae | 33.3 | 2 | 0.3 | 0.5 | 24.5 | 24 | 25 | 50.0 | 6 | 1.0 | 1.3 | 20.2 | 15 | 23 | | |
| princess parrotfish | <i>Scarus taeniopterus</i> | Scaridae | 66.7 | 16 | 2.7 | 2.9 | 17.8 | 10 | 22 | 100.0 | 16 | 2.7 | 1.6 | 20.9 | 14 | 30 | | |
| queen parrotfish | <i>Scarus vetula</i> | Scaridae | 16.7 | 3 | 0.5 | 1.2 | 24.0 | 18 | 31 | 16.7 | 1 | 0.2 | 0.4 | 40.0 | 40 | 40 | | |
| redband parrotfish | <i>Sparisoma aurofrenatum</i> | Scaridae | 100.0 | 18 | 3.0 | 1.9 | 15.5 | 6 | 20 | 83.3 | 20 | 3.3 | 2.8 | 15.6 | 5 | 23 | | |
| redtail parrotfish | <i>Sparisoma chrysopterus</i> | Scaridae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 23.0 | 23 | 23 | | |
| yellowtail parrotfish | <i>Sparisoma rubripinne</i> | Scaridae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 25.0 | 25 | 25 | | |
| stoplight parrotfish | <i>Sparisoma viride</i> | Scaridae | 16.7 | 2 | 0.3 | 0.8 | 15.0 | 10 | 20 | 50.0 | 5 | 0.8 | 1.0 | 26.0 | 13 | 32 | | |
| cero mackerel | <i>Scomberomorus regalis</i> | Scombridae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 30.0 | 30 | 30 | | |
| coney | <i>Cephalopholis fulvus</i> | Serranidae | 100.0 | 19 | 3.2 | 1.5 | 11.5 | 5 | 18 | 100.0 | 20 | 3.3 | 1.0 | 18.7 | 12 | 28 | | |
| yellowtail hamlet | <i>Hypoplectrus chlorurus</i> | Serranidae | 16.7 | 2 | 0.3 | 0.8 | 12.5 | 10 | 15 | 0.0 | 0 | 0.0 | 0.0 | - | - | - | | |
| shy hamlet | <i>Hypoplectrus guttavarius</i> | Serranidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 11.0 | 11 | 11 | | |
| barred hamlet | <i>Hypoplectrus puella</i> | Serranidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 10.0 | 10 | 10 | | |
| harlequin bass | <i>Serranus tigrinus</i> | Serranidae | 66.7 | 4 | 0.7 | 0.5 | 8.0 | 6 | 9 | 66.7 | 10 | 1.7 | 1.6 | 8.9 | 7 | 10 | | |
| sand diver | <i>Synodus intermedius</i> | Synodontidae | 16.7 | 1 | 0.2 | 0.4 | 22.0 | 22 | 22 | 0.0 | 0 | 0.0 | 0.0 | - | - | - | | |
| sharpnose puffer | <i>Canthigaster rostrata</i> | Tetraodontidae | 16.7 | 1 | 0.2 | 0.4 | 4.0 | 4 | 4 | 50.0 | 3 | 0.5 | 0.5 | 7.7 | 4 | 13 | | |
| Total = | | | | 1922 | | | | | | | | Total = | 2221 | | | | | |

Appendix 2E. Fish census data from Eagle Ray

| Common Name | Species | Family | Spring '02 Census | | | | | | | Fall '02 Census | | | | | | |
|------------------------|-------------------------------|----------------|-------------------|-------------|------|-------|--------|-----|-----|-----------------|-------------|------|-------|------|-----|-----|
| | | | % Freq | No. of Fish | | | Size | | | % Freq | No. of Fish | | | Size | | |
| | | | (n=6) | Total | Avg. | StDev | Avg | Min | Max | (n=6) | Total | Avg. | StDev | Avg | Min | Max |
| ocean surgeonfish | <i>Acanthurus bahianus</i> | Acanthuridae | 100.0 | 54 | 9.0 | 1.8 | 15 | 12 | 18 | 100.0 | 37 | 6.2 | 2.6 | 12.8 | 7 | 16 |
| doctorfish | <i>Acanthurus chirurgus</i> | Acanthuridae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 3 | 0.5 | 1.2 | 18.0 | 16 | 20 |
| blue tang | <i>Acanthurus coeruleus</i> | Acanthuridae | 66.7 | 6 | 1.0 | 0.9 | 15.625 | 12 | 18 | 83.3 | 14 | 2.3 | 1.8 | 13.7 | 8 | 18 |
| trumpetfish | <i>Aulostomus maculatus</i> | Aulostomidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 33.3 | 2 | 0.3 | 0.5 | 41.0 | 32 | 50 |
| black durgon | <i>Melichthys niger</i> | Balistidae | 16.7 | 2 | 0.3 | 0.8 | 23 | 22 | 24 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| bar jack | <i>Caranx ruber</i> | Carangidae | 0.0 | 0 | 0.00 | 0.0 | - | - | - | 33.3 | 7 | 1.2 | 2.0 | 22.5 | 20 | 30 |
| four-eye butterflyfish | <i>Chaetodon capistratus</i> | Chaetodontidae | 83.3 | 11 | 1.8 | 1.0 | 9.2 | 7 | 12 | 83.3 | 13 | 2.2 | 1.2 | 10.1 | 9 | 12 |
| banded butterflyfish | <i>Chaetodon striatus</i> | Chaetodontidae | 0.0 | 0 | 0.00 | 0.0 | - | - | - | 66.7 | 8 | 1.3 | 1.0 | 11.5 | 10 | 15 |
| caesar grunt | <i>Haemulon carbonarium</i> | Haemulidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 30.0 | 30 | 30 |
| french grunt | <i>Haemulon flavolineatum</i> | Haemulidae | 83.3 | 10 | 1.7 | 1.8 | 15.8 | 15 | 17 | 100.0 | 13 | 2.2 | 1.0 | 14.8 | 12 | 18 |
| white grunt | <i>Haemulon plumieri</i> | Haemulidae | 33.3 | 2 | 0.3 | 0.5 | 23 | 21 | 25 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| squirrelfish | <i>Holocentrus acensionis</i> | Holocentridae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 50.0 | 6 | 1.0 | 1.3 | 19.0 | 17 | 31 |
| longspine squirrelfish | <i>Holocentrus rufus</i> | Holocentridae | 50.0 | 4 | 0.7 | 0.8 | 18.167 | 17 | 20 | 33.3 | 2 | 0.3 | 0.5 | 16.0 | 16 | 16 |
| blackbar soldierfish | <i>Myripristis jacobus</i> | Holocentridae | 50.0 | 19 | 3.2 | 4.8 | 15.167 | 12 | 17 | 83.3 | 18 | 3.0 | 3.6 | 13.8 | 10 | 18 |
| spanish hogfish | <i>Bodianus rufus</i> | Labridae | 16.7 | 1 | 0.2 | 0.4 | 20 | 20 | 20 | 66.7 | 5 | 0.8 | 0.8 | 11.4 | 5 | 25 |
| creole wrasse | <i>Clepticus parrae</i> | Labridae | 66.7 | 96 | 16.0 | 12.9 | 7 | 3 | 12 | 50.0 | 260 | 43.3 | 70.9 | 6.3 | 2 | 15 |

Appendix 2E (continued). Fish census data from Eagle Ray

| Common Name | Species | Family | Spring '02 Census | | | | | | | Fall '02 Census | | | | | | |
|-----------------------|----------------------------------|---------------|-------------------|-------------|-------|-------|--------|-----|-----|-----------------|-------------|-------|-------|------|-----|-----|
| | | | % Freq | No. of Fish | | | Size | | | % Freq | No. of Fish | | | Size | | |
| | | | (n=6) | Total | Avg. | StDev | Avg | Min | Max | (n=6) | Total | Avg. | StDev | Avg | Min | Max |
| yellowcheek wrasse | <i>Halichoeres cyanocephalus</i> | Labridae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 15.0 | 15 | 15 |
| yellowhead wrasse | <i>Halichoeres garnoti</i> | Labridae | 100.0 | 18 | 3.0 | 1.4 | 8.3333 | 3 | 15 | 83.3 | 29 | 4.8 | 3.3 | 8.6 | 4 | 16 |
| bluehead wrasse | <i>Thalassoma bifasciatum</i> | Labridae | 100.0 | 1460 | 243.3 | 36.1 | 4.25 | 2 | 10 | 100.0 | 805 | 134.2 | 75.7 | 4.6 | 2 | 12 |
| mutton snapper | <i>Lutjanus analis</i> | Lutjanidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 33.3 | 2 | 0.3 | 0.5 | 35.0 | 35 | 35 |
| schoolmaster | <i>Lutjanus apodus</i> | Lutjanidae | 16.7 | 2 | 0.3 | 0.8 | 17 | 17 | 17 | 16.7 | 1 | 0.2 | 0.4 | 17.0 | 17 | 17 |
| yellowtail snapper | <i>Ocyurus chrysurus</i> | Lutjanidae | 83.3 | 10 | 1.7 | 1.8 | 20.1 | 15 | 25 | 50.0 | 5 | 0.8 | 1.2 | 20.7 | 20 | 25 |
| yellow goatfish | <i>Mulloidichthys martinicus</i> | Mullidae | 33.3 | 2 | 0.3 | 0.5 | 23.5 | 22 | 25 | 16.7 | 7 | 1.2 | 2.9 | 17.0 | 0 | 19 |
| spotted goatfish | <i>Pseudupeneus maculatus</i> | Mullidae | 50.0 | 3 | 0.5 | 0.5 | 16.333 | 14 | 19 | 83.3 | 6 | 1.0 | 0.6 | 17.2 | 15 | 20 |
| honeycomb cowfish | <i>Acanthostracion ploygonia</i> | Ostraciidae | 16.7 | 1 | 0.2 | 0.4 | 24 | 24 | 24 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| spotted trunkfish | <i>Lactophrys bicaudalis</i> | Ostraciidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 13.0 | 13 | 13 |
| smooth trunkfish | <i>Lactophrys triqueter</i> | Ostraciidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 14.0 | 14 | 14 |
| rock beauty | <i>Holacanthus tricolor</i> | Pomacanthidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 17.0 | 17 | 17 |
| sergeant major | <i>Abudefduf saxatilis</i> | Pomacentridae | 33.3 | 12 | 2.0 | 3.1 | 11.5 | 10 | 14 | 83.3 | 57 | 9.5 | 7.4 | 11.7 | 8 | 14 |
| blue chromis | <i>Chromis cyanea</i> | Pomacentridae | 100.0 | 104 | 17.3 | 12.9 | 4.6667 | 2 | 8 | 100.0 | 200 | 33.3 | 11.1 | 5.0 | 3 | 8 |
| brown chromis | <i>Chromis multilineata</i> | Pomacentridae | 16.7 | 10 | 1.7 | 4.1 | 3 | 3 | 4 | 66.7 | 260 | 43.3 | 68.4 | 3.6 | 3 | 6 |
| yellowtail damselfish | <i>Microspathodon chrysurus</i> | Pomacentridae | 66.7 | 7 | 1.2 | 1.5 | 11.25 | 6 | 12 | 33.3 | 4 | 0.7 | 1.2 | 11.0 | 8 | 12 |

Appendix 2E (continued). Fish census data from Eagle Ray

| Common Name | Species | Family | Spring '02 Census | | | | | | | Fall '02 Census | | | | | | |
|--------------------------------|---|---------------|-------------------|-------------|------|-------|--------|-----|-----|-----------------|-------------|------|-------|------|-----|-----|
| | | | % Freq (n=6) | No. of Fish | | | Size | | | % Freq (n=6) | No. of Fish | | | Size | | |
| | | | | Total | Avg. | StDev | Avg | Min | Max | | Total | Avg. | StDev | Avg | Min | Max |
| dusky damselfish | <i>Stegastes adustus</i> (<i>S. fuscus</i>) | Pomacentridae | 83.3 | 31 | 5.2 | 4.0 | 7.2 | 5 | 10 | 83.3 | 22 | 3.7 | 2.2 | 8.2 | 6 | 12 |
| beaugregory bicolor damselfish | <i>Stegastes leucostictus</i> | Pomacentridae | 66.7 | 9 | 1.5 | 1.5 | 6.5 | 4 | 9 | 33.3 | 7 | 1.2 | 1.8 | 9.5 | 10 | 12 |
| cocoa damselfish | <i>Stegastes partitus</i> | Pomacentridae | 100.0 | 192 | 32.0 | 8.1 | 4.3333 | 2 | 7 | 100.0 | 251 | 41.8 | 19.6 | 3.8 | 2 | 6 |
| striped parrotfish | <i>Stegastes variabilis</i> | Pomacentridae | 16.7 | 5 | 0.8 | 2.0 | 6 | 5 | 8 | 33.3 | 6 | 1.0 | 1.7 | 9.0 | 5 | 12 |
| princess parrotfish | <i>Scarus iserti</i> (<i>S. croicensis</i>) | Scaridae | 50.0 | 8 | 1.3 | 2.0 | 16.833 | 10 | 21 | 50.0 | 20 | 3.3 | 5.6 | 14.0 | 7 | 21 |
| queen parrotfish | <i>Scarus taeniopterus</i> | Scaridae | 83.3 | 15 | 2.5 | 1.9 | 17.7 | 10 | 26 | 66.7 | 13 | 2.2 | 2.2 | 15.5 | 10 | 22 |
| redband parrotfish | <i>Scarus vetula</i> | Scaridae | 16.7 | 3 | 0.5 | 1.2 | 30 | 25 | 35 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| redtail parrotfish | <i>Sparisoma aurofrenatum</i> | Scaridae | 83.3 | 21 | 3.5 | 2.7 | 16.9 | 8 | 24 | 83.3 | 26 | 4.3 | 3.1 | 16.9 | 10 | 22 |
| stoplight parrotfish | <i>Sparisoma chrysopteron</i> | Scaridae | 50.0 | 4 | 0.7 | 0.8 | 26.5 | 20 | 32 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| graysby | <i>Sparisoma viride</i> | Scaridae | 66.7 | 5 | 0.8 | 0.8 | 23 | 5 | 35 | 83.3 | 9 | 1.5 | 1.0 | 25.9 | 7 | 40 |
| coney | <i>Cephalopholis cruentatus</i> | Serranidae | 66.7 | 9 | 1.5 | 1.6 | 16.5 | 14 | 25 | 50.0 | 4 | 0.7 | 0.8 | 18.7 | 17 | 20 |
| red hind | <i>Cephalopholis fulvus</i> | Serranidae | 50.0 | 7 | 1.2 | 1.3 | 14.667 | 10 | 17 | 66.7 | 11 | 1.8 | 1.6 | 17.6 | 12 | 21 |
| yellowtail hamlet | <i>Epinephelus guttatus</i> | Serranidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 18.0 | 18 | 18 |
| black hamlet | <i>Hypoplectrus chlorurus</i> | Serranidae | 16.7 | 1 | 0.2 | 0.4 | 12 | 12 | 12 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| butter hamlet | <i>Hypoplectrus nigricans</i> | Serranidae | 16.7 | 1 | 0.2 | 0.4 | 13 | 13 | 13 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| | <i>Hypoplectrus unicolor</i> | Serranidae | 16.7 | 1 | 0.2 | 0.4 | 11 | 11 | 11 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |

Appendix 2E (continued). Fish census data from Eagle Ray

| Common Name | Species | Family | Spring '02 Census | | | | | | | Fall '02 Census | | | | | | |
|------------------|------------------------------|----------------|-------------------|-------------|------|-------|------|-----|-----|-----------------|-------------|------|-------|------|-----|-----|
| | | | % Freq (n=6) | No. of Fish | | | Size | | | % Freq (n=6) | No. of Fish | | | Size | | |
| | | | | Total | Avg. | StDev | Avg | Min | Max | | Total | Avg. | StDev | Avg | Min | Max |
| tobaccofish | <i>Serranus tabacarius</i> | Serranidae | 16.7 | 2 | 0.3 | 0.8 | 4 | 4 | 4 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| harlequin bass | <i>Serranus tigrinus</i> | Serranidae | 16.7 | 3 | 0.5 | 1.2 | 8 | 7 | 9 | 50.0 | 6 | 1.0 | 1.3 | 6.3 | 5 | 8 |
| jolthead pogy | <i>Calamus bajonado</i> | Sparidae | 33.3 | 2 | 0.3 | 0.5 | 25.5 | 25 | 26 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| sharpnose puffer | <i>Canthigaster rostrata</i> | Tetraodontidae | 33.3 | 3 | 0.5 | 0.8 | 5.75 | 5 | 6 | 33.3 | 3 | 0.5 | 0.8 | 5.5 | 7 | 7 |
| Total = | | | | 2156 | | | | | | | 2148 | | | | | |

Appendix 2F. Fish census data from Cane Bay

| Common Name | Species | Family | Spring '02 Census | | | | | | | Fall '02 Census | | | | | | |
|------------------------|-------------------------------|----------------|-------------------|-------------|------|-------|------|-----|-----|-----------------|-------------|----------|-------|------|-----|-----|
| | | | % Freq (n=6) | No. of Fish | | | Size | | | % Freq (n=6) | No. of Fish | | | Size | | |
| | | | | Total | Avg. | StDev | Avg | Min | Max | | Total No. | Avg. No. | StDev | Avg | Min | Max |
| ocean surgeonfish | <i>Acanthurus bahianus</i> | Acanthuridae | 100.0 | 24 | 4.0 | 1.1 | 14.7 | 10 | 20 | 100.0 | 29 | 4.8 | 2.2 | 16.0 | 11 | 20 |
| doctorfish | <i>Acanthurus chirurgus</i> | Acanthuridae | 16.7 | 1 | 0.2 | 0.4 | 15.0 | 15 | 15 | 50.0 | 5 | 0.8 | 1.0 | 17.0 | 15 | 20 |
| blue tang | <i>Acanthurus coeruleus</i> | Acanthuridae | 100.0 | 13 | 2.2 | 0.4 | 15.8 | 10 | 20 | 66.7 | 10 | 1.7 | 1.6 | 15.3 | 11 | 18 |
| trumpetfish | <i>Aulostomus maculatus</i> | Aulostomidae | 16.7 | 1 | 0.2 | 0.4 | 35.0 | 35 | 35 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| orangespotted filefish | <i>Cantherhines pullus</i> | Monacanthidae | 33.3 | 3 | 0.5 | 0.8 | 12.8 | 12 | 14 | 33.3 | 3 | 0.5 | 0.9 | 15.5 | 14 | 16 |
| black durgon | <i>Melichthys niger</i> | Balistidae | 100.0 | 39 | 6.5 | 6.4 | 22.0 | 12 | 28 | 100.0 | 66 | 11.0 | 6.7 | 22.7 | 13 | 30 |
| bar jack | <i>Caranx ruber</i> | Carangidae | 16.7 | 1 | 0.2 | 0.4 | 17.0 | 17 | 17 | 83.3 | 11 | 1.8 | 1.5 | 22.0 | 12 | 30 |
| four-eye butterflyfish | <i>Chaetodon capistratus</i> | Chaetodontidae | 66.7 | 8 | 1.3 | 1.0 | 11.3 | 8 | 15 | 50.0 | 5 | 0.8 | 1.0 | 9.7 | 8 | 10 |
| fairy basslet | <i>Gramma loreto</i> | Grammatidae | 50.0 | 10 | 1.7 | 2.1 | 2.7 | 1 | 3 | 66.7 | 32 | 5.3 | 4.6 | 4.0 | 2 | 5 |
| caesar grunt | <i>Haemulon carbonarium</i> | Haemulidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 33.3 | 2 | 0.3 | 0.5 | 21.5 | 18 | 25 |
| smallmouth grunt | <i>Haemulon chrysargyreum</i> | Haemulidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 2 | 0.3 | 0.9 | 14.5 | 14 | 15 |
| french grunt | <i>Haemulon flavolineatum</i> | Haemulidae | 50.0 | 20 | 3.3 | 6.7 | 13.5 | 12 | 15 | 66.7 | 13 | 2.2 | 2.2 | 16.0 | 12 | 20 |
| white grunt | <i>Haemulon plumieri</i> | Haemulidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 18.0 | 18 | 18 |
| bluestriped grunt | <i>Haemulon sciurus</i> | Haemulidae | 16.7 | 3 | 0.5 | 1.2 | 22.0 | 20 | 25 | 16.7 | 4 | 0.7 | 1.6 | 24.0 | 20 | 25 |
| squirrelfish | <i>Holocentrus adensionis</i> | Holocentridae | 16.7 | 1 | 0.2 | 0.4 | 13.0 | 13 | 13 | 16.7 | 3 | 0.5 | 1.3 | 18.0 | 16 | 20 |
| longspine squirrelfish | <i>Holocentrus rufus</i> | Holocentridae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 15.0 | 15 | 15 |
| blackbar soldierfish | <i>Myripristis jacobus</i> | Holocentridae | 16.7 | 2 | 0.3 | 0.8 | 15.0 | 15 | 15 | 33.3 | 5 | 0.8 | 1.7 | 17.0 | 15 | 20 |

Appendix 2F (continued). Fish census data from Cane Bay

| Common Name | Species | Family | Spring '02 Census | | | | | | | Fall '02 Census | | | | | | |
|--------------------------------|--------------------------------------|---------------|-------------------|-------------|-------|-------|------|-----|-----|-----------------|-------------|----------|-------|------|-----|-----|
| | | | % Freq (n=6) | No. of Fish | | | Size | | | % Freq (n=6) | No. of Fish | | | Size | | |
| | | | | Total | Avg. | StDev | Avg | Min | Max | | Total No. | Avg. No. | StDev | Avg | Min | Max |
| longjaw squirrelfish | <i>Neoniphon marianus</i> | Holocentridae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 13.0 | 13 | 13 |
| spanish hogfish | <i>Bodianus rufus</i> | Labridae | 50.0 | 3 | 0.5 | 0.5 | 19.7 | 17 | 22 | 33.3 | 2 | 0.3 | 0.5 | 25.0 | 20 | 30 |
| creole wrasse | <i>Clepticus parrae</i> | Labridae | 66.7 | 311 | 51.8 | 84.9 | 7.0 | 1 | 16 | 50.0 | 2332 | 388.7 | 463.4 | 4.0 | 2 | 18 |
| yellowhead wrasse | <i>Halichoeres garnoti</i> | Labridae | 66.7 | 10 | 1.7 | 1.6 | 8.0 | 5 | 12 | 66.7 | 24 | 4.0 | 4.3 | 6.1 | 4 | 15 |
| bluehead wrasse | <i>Thalassoma bifasciatum</i> | Labridae | 100.0 | 722 | 120.3 | 43.5 | 2.3 | 1 | 6 | 100.0 | 600 | 100.0 | 91.0 | 4.4 | 2 | 13 |
| schoolmaster mahogany snapper | <i>Lutjanus apodus</i> | Lutjanidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 50.0 | 5 | 0.8 | 1.0 | 36.2 | 30 | 40 |
| yellow snapper | <i>Lutjanus mahogoni</i> | Lutjanidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 33.3 | 4 | 0.7 | 1.3 | 19.5 | 18 | 22 |
| yellow goatfish | <i>Mulloidichthys martinicus</i> | Mullidae | 33.3 | 5 | 0.8 | 1.3 | 16.8 | 12 | 22 | 50.0 | 12 | 2.0 | 3.4 | 17.0 | 12 | 19 |
| spotted trunkfish | <i>Lactophrys bicaudalis</i> | Ostraciidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 18.0 | 18 | 18 |
| sergeant major | <i>Abudefduf saxatilis</i> | Pomacentridae | 83.3 | 10 | 1.7 | 1.0 | 11.8 | 10 | 15 | 83.3 | 13 | 2.2 | 2.1 | 13.0 | 10 | 20 |
| blue chromis | <i>Chromis cyanea</i> | Pomacentridae | 100.0 | 682 | 113.7 | 79.0 | 3.7 | 1 | 10 | 100.0 | 642 | 107.0 | 62.3 | 5.0 | 2 | 10 |
| brown chromis | <i>Chromis multilineata</i> | Pomacentridae | 100.0 | 901 | 150.2 | 101.7 | 4.3 | 1 | 8 | 83.3 | 1170 | 195.0 | 126.2 | 5.2 | 2 | 10 |
| yellowtail damselfish | <i>Microspathodon chrysurus</i> | Pomacentridae | 83.3 | 12 | 2.0 | 1.1 | 9.9 | 4 | 14 | 83.3 | 13 | 2.2 | 1.2 | 13.3 | 4 | 20 |
| dusky damselfish | <i>Stegastes adustus</i> (S. fuscus) | Pomacentridae | 100.0 | 48 | 8.0 | 5.3 | 8.2 | 5 | 12 | 100.0 | 36 | 6.0 | 2.8 | 9.8 | 5 | 15 |
| beaugregory bicolor damselfish | <i>Stegastes leucostictus</i> | Pomacentridae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 7.0 | 7 | 7 |
| | <i>Stegastes partitus</i> | Pomacentridae | 100.0 | 227 | 37.8 | 14.0 | 3.7 | 1 | 7 | 100.0 | 285 | 47.5 | 26.0 | 4.5 | 2 | 8 |

Appendix 2F (continued). Fish census data from Cane Bay

| Common Name | Species | Family | Spring '02 Census | | | | | | | Fall '02 Census | | | | | | | | |
|----------------------|---|----------------|-------------------|-------------|------|-------|------|-----|-----|-----------------|-------------|----------|-------|------|-----|-----|--|--|
| | | | % Freq (n=6) | No. of Fish | | | Size | | | % Freq (n=6) | No. of Fish | | | Size | | | | |
| | | | | Total | Avg. | StDev | Avg | Min | Max | | Total No. | Avg. No. | StDev | Avg | Min | Max | | |
| threespot damselfish | <i>Stegastes planifrons</i> | Pomacentridae | 50.0 | 10 | 1.7 | 2.0 | 5.7 | 3 | 10 | 66.7 | 24 | 4.0 | 5.4 | 7.3 | 3 | 11 | | |
| striped parrotfish | <i>Scarus iserti</i> (<i>S. croicensis</i>) | Scaridae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 50.0 | 11 | 1.8 | 2.4 | 8.7 | 5 | 25 | | |
| princess parrotfish | <i>Scarus taeniopterus</i> | Scaridae | 100.0 | 18 | 3.0 | 1.1 | 16.9 | 10 | 22 | 83.3 | 20 | 3.3 | 3.2 | 19.3 | 5 | 30 | | |
| queen parrotfish | <i>Scarus vetula</i> | Scaridae | 16.7 | 1 | 0.2 | 0.4 | 32.0 | 32 | 32 | 0.0 | 0 | 0.0 | 0.0 | - | - | - | | |
| redband parrotfish | <i>Sparisoma aurofrenatum</i> | Scaridae | 83.3 | 20 | 3.3 | 2.3 | 17.4 | 7 | 24 | 66.7 | 14 | 2.3 | 1.6 | 16.8 | 10 | 30 | | |
| stoplight parrotfish | <i>Sparisoma viride</i> | Scaridae | 50.0 | 5 | 0.8 | 1.2 | 20.7 | 6 | 35 | 83.3 | 17 | 2.8 | 2.3 | 27.0 | 5 | 35 | | |
| graysby | <i>Cephalopholis cruentatus</i> | Serranidae | 33.3 | 2 | 0.3 | 0.5 | 20.0 | 18 | 22 | 83.3 | 11 | 1.8 | 0.8 | 18.3 | 15 | 23 | | |
| coney | <i>Cephalopholis fulvus</i> | Serranidae | 50.0 | 6 | 1.0 | 1.1 | 14.8 | 12 | 18 | 50.0 | 5 | 0.8 | 1.2 | 17.0 | 15 | 20 | | |
| black hamlet | <i>Hypoplectrus nigricans</i> | Serranidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 50.0 | 3 | 0.5 | 0.5 | 9.0 | 6 | 12 | | |
| harlequin bass | <i>Serranus tigrinus</i> | Serranidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 6.0 | 6 | 6 | | |
| sharpnose puffer | <i>Canthigaster rostrata</i> | Tetraodontidae | 16.7 | 1 | 0.2 | 0.4 | 6.0 | 6 | 6 | 33.3 | 4 | 0.7 | 1.1 | 5.5 | 4 | 7 | | |
| Total = | | | | 3120 | | | | | | | | Total = | 5443 | | | | | |

Appendix 2G. Fish census data from Sprat Hole

| Common Name | Species Name | Family | Spring '02 Census | | | | | | | Fall '02 Census | | | | | | |
|------------------------|--------------------------------|----------------|-------------------|-------------|-------|-------|------|-----|-----|-----------------|-------------|-------|-------|------|-----|-----|
| | | | % Freq (n=6) | No. of Fish | | | Size | | | % Freq (n=6) | No. of Fish | | | Size | | |
| | | | | Total | Avg. | StDev | Avg | Min | Max | | Total | Avg. | StDev | Avg | Min | Max |
| ocean surgeonfish | <i>Acanthurus bahianus</i> | Acanthuridae | 83.3 | 16 | 2.7 | 2.1 | 15.1 | 12 | 18 | 83.3 | 14 | 2.3 | 2.3 | 15.1 | 12 | 20 |
| blue tang | <i>Acanthurus coeruleus</i> | Acanthuridae | 83.3 | 15 | 2.5 | 1.5 | 14.4 | 10 | 18 | 83.3 | 9 | 1.5 | 1.0 | 16.8 | 15 | 20 |
| trumpetfish | <i>Aulostomus maculatus</i> | Aulostomidae | 16.7 | 1 | 0.2 | 0.4 | 50.0 | 50 | 50 | 50.0 | 3 | 0.5 | 0.5 | 24.7 | 15 | 34 |
| blue runner | <i>Caranx crysos</i> | Carangidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 30.0 | 30 | 30 |
| bar jack | <i>Caranx ruber</i> | Carangidae | 33.3 | 5 | 0.8 | 1.6 | 28.0 | 26 | 30 | 16.7 | 2 | 0.3 | 0.8 | 21.0 | 20 | 22 |
| four-eye butterflyfish | <i>Chaetodon capistratus</i> | Chaetodontidae | 33.3 | 4 | 0.7 | 1.0 | 9.0 | 7 | 11 | 83.3 | 10 | 1.7 | 0.8 | 9.6 | 7 | 12 |
| yellowfin mojarra | <i>Gerres cinereus</i> | Gerreidae | 16.7 | 2 | 0.3 | 0.8 | 18.5 | 17 | 20 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| fairy basslet | <i>Gramma loreto</i> | Grammatidae | 16.7 | 1 | 0.2 | 0.4 | 2.0 | 2 | 2 | 50.0 | 5 | 0.8 | 1.0 | 4.0 | 3 | 5 |
| caesar grunt | <i>Haemulon carbonarium</i> | Haemulidae | 16.7 | 2 | 0.3 | 0.8 | 17.5 | 15 | 20 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| french grunt | <i>Haemulon flavolineatum</i> | Haemulidae | 66.7 | 7 | 1.2 | 1.2 | 15.5 | 14 | 17 | 66.7 | 5 | 0.8 | 0.8 | 17.0 | 15 | 20 |
| squirrelfish | <i>Holocentrus adensionis</i> | Holocentridae | 16.7 | 1 | 0.2 | 0.4 | 17.0 | 17 | 17 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| blackbar soldierfish | <i>Myripristis jacobus</i> | Holocentridae | 83.3 | 16 | 2.7 | 2.9 | 14.2 | 10 | 17 | 50.0 | 12 | 2.0 | 2.3 | 15.7 | 14 | 21 |
| spanish hogfish | <i>Bodianus rufus</i> | Labridae | 16.7 | 1 | 0.2 | 0.4 | 22.0 | 22 | 22 | 50.0 | 6 | 1.0 | 1.3 | 15.2 | 5 | 25 |
| creole wrasse | <i>Clepticus parrae</i> | Labridae | 100.0 | 1467 | 244.5 | 252.7 | 5.5 | 1 | 14 | 66.7 | 685 | 114.2 | 240.1 | 9.0 | 2 | 16 |
| yellowhead wrasse | <i>Halichoeres garnoti</i> | Labridae | 83.3 | 100 | 16.7 | 31.3 | 6.3 | 2 | 14 | 66.7 | 19 | 3.2 | 3.0 | 7.6 | 4 | 12 |
| clown wrasse | <i>Halichoeres maculipinna</i> | Labridae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 11.0 | 11 | 11 |
| bluehead wrasse | <i>Thalassoma bifasciatum</i> | Labridae | 83.3 | 650 | 108.3 | 79.6 | 2.7 | 1 | 10 | 83.3 | 325 | 54.2 | 31.8 | 4.0 | 2 | 10 |

Appendix 2G (continued). Fish census data from Sprat Hole

| Common Name | Species Name | Family | Spring '02 Census | | | | | | | Fall '02 Census | | | | | | |
|-------------------------------|--------------------------------------|---------------|-------------------|-------------|-------|-------|------|-----|-----|-----------------|-------------|-------|-------|------|-----|-----|
| | | | % Freq (n=6) | No. of Fish | | | Size | | | % Freq (n=6) | No. of Fish | | | Size | | |
| | | | | Total | Avg. | StDev | Avg | Min | Max | | Total | Avg. | StDev | Avg | Min | Max |
| mutton snapper | <i>Lutjanus analis</i> | Lutjanidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 38.0 | 38 | 38 |
| | <i>Lutjanus</i> | | | | | | | | | | | | | | | |
| schoolmaster mahogany snapper | <i>apodus</i> | Lutjanidae | 16.7 | 2 | 0.3 | 0.8 | 47.0 | 40 | 55 | 16.7 | 1 | 0.2 | 0.4 | 40.0 | 40 | 40 |
| | <i>Lutjanus mahogoni</i> | Lutjanidae | 66.7 | 20 | 3.3 | 5.0 | 16.3 | 10 | 20 | 100.0 | 20 | 3.3 | 1.6 | 20.3 | 15 | 25 |
| | <i>Malacanthus plumieri</i> | Malacanthidae | 16.7 | 1 | 0.2 | 0.4 | 40.0 | 40 | 40 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| sand tilefish | <i>Mulloidichthys martinicus</i> | Mullidae | 33.3 | 6 | 1.0 | 2.0 | 14.0 | 12 | 16 | 50.0 | 127 | 21.2 | 31.2 | 14.3 | 10 | 24 |
| yellow goatfish | <i>Gymnothorax moraya</i> | Muraenidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 60.0 | 60 | 60 |
| honeycomb cowfish | <i>Acanthostracion ploygonia</i> | Ostraciidae | 16.7 | 1 | 0.2 | 0.4 | 30.0 | 30 | 30 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| smooth trunkfish | <i>Lactophrys triqueter</i> | Ostraciidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 10.0 | 10 | 10 |
| | <i>Holacanthus tricolor</i> | Pomacanthidae | 33.3 | 2 | 0.3 | 0.5 | 13.5 | 12 | 15 | 50.0 | 4 | 0.7 | 0.8 | 18.7 | 15 | 22 |
| rock beauty sergeant major | <i>Abudefduf saxatilis</i> | Pomacentridae | 16.7 | 2 | 0.3 | 0.8 | 13.0 | 13 | 13 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| blue chromis | <i>Chromis cyanea</i> | Pomacentridae | 100.0 | 800 | 133.3 | 48.9 | 2.8 | 1 | 7 | 100.0 | 789 | 131.5 | 61.9 | 4.7 | 2 | 9 |
| brown chromis | <i>Chromis multilineata</i> | Pomacentridae | 100.0 | 701 | 116.8 | 32.0 | 4.5 | 1 | 7 | 83.3 | 650 | 108.3 | 70.3 | 4.8 | 2 | 8 |
| yellowtail damselfish | <i>Microspathodon chrysurus</i> | Pomacentridae | 16.7 | 2 | 0.3 | 0.8 | 10.0 | 10 | 10 | 0.0 | 0 | 0.0 | 0.0 | - | - | - |
| | <i>Stegastes adustus</i> (S. fuscus) | Pomacentridae | 83.3 | 29 | 4.8 | 3.8 | 6.4 | 4 | 10 | 66.7 | 26 | 4.3 | 4.4 | 10.3 | 7 | 15 |
| dusky damselfish | <i>Stegastes partitus</i> | Pomacentridae | 100.0 | 235 | 39.2 | 13.6 | 4.1 | 1 | 6 | 100.0 | 397 | 66.2 | 29.6 | 4.5 | 2 | 10 |
| bicolor damselfish | <i>Stegastes planifrons</i> | Pomacentridae | 100.0 | 101 | 16.8 | 5.7 | 5.5 | 3 | 8 | 83.3 | 70 | 11.7 | 9.2 | 8.2 | 4 | 15 |

Appendix 2G (continued). Fish census data from Sprat Hole

| Common Name | Species Name | Family | Spring '02 Census | | | | | | | Fall '02 Census | | | | | | | | |
|----------------------|--------------------------------------|----------------|-------------------|-------------|------|-------|------|-----|-----|-----------------|-------------|------|-------|------|-----|-----|--|--|
| | | | % Freq (n=6) | No. of Fish | | | Size | | | % Freq (n=6) | No. of Fish | | | Size | | | | |
| | | | | Total | Avg. | StDev | Avg | Min | Max | | Total | Avg. | StDev | Avg | Min | Max | | |
| cocoa damselfish | <i>Stegastes variabilis</i> | Pomacentridae | 16.7 | 1 | 0.2 | 0.4 | 5.0 | 5 | 5 | 0.0 | 0 | 0.0 | 0.0 | - | - | - | | |
| striped parrotfish | <i>Scarus iserti (S. croicensis)</i> | Scaridae | 16.7 | 4 | 0.7 | 1.6 | 5.0 | 4 | 6 | 83.3 | 26 | 4.3 | 2.7 | 11.6 | 4 | 20 | | |
| princess parrotfish | <i>Scarus taeniopterus</i> | Scaridae | 100.0 | 46 | 7.7 | 5.0 | 17.2 | 6 | 25 | 100.0 | 35 | 5.8 | 4.0 | 18.6 | 5 | 30 | | |
| queen parrotfish | <i>Scarus vetula</i> | Scaridae | 66.7 | 10 | 1.7 | 1.5 | 26.8 | 15 | 45 | 33.3 | 2 | 0.3 | 0.5 | 27.5 | 25 | 30 | | |
| redband parrotfish | <i>Sparisoma aurofrenatum</i> | Scaridae | 100.0 | 41 | 6.8 | 1.9 | 16.2 | 10 | 25 | 83.3 | 25 | 4.2 | 2.7 | 16.0 | 3 | 25 | | |
| stoplight parrotfish | <i>Sparisoma viride</i> | Scaridae | 100.0 | 13 | 2.2 | 0.8 | 21.6 | 6 | 40 | 66.7 | 14 | 2.3 | 2.6 | 13.5 | 4 | 25 | | |
| graysby | <i>Cephalopholis cruentatus</i> | Serranidae | 66.7 | 9 | 1.5 | 1.4 | 14.9 | 12 | 19 | 66.7 | 7 | 1.2 | 1.0 | 16.5 | 10 | 22 | | |
| coney | <i>Cephalopholis fulvus</i> | Serranidae | 16.7 | 4 | 0.7 | 1.6 | 17.0 | 15 | 20 | 66.7 | 9 | 1.5 | 1.5 | 16.1 | 8 | 22 | | |
| yellowtail hamlet | <i>Hypoplectrus chlorurus</i> | Serranidae | 0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 2 | 0.3 | 0.8 | 9.0 | 8 | 10 | | |
| shy hamlet | <i>Hypoplectrus guttavarius</i> | Serranidae | 16.7 | 1 | 0.2 | 0.4 | 8.0 | 8 | 8 | 0.0 | 0 | 0.0 | 0.0 | - | - | - | | |
| black hamlet | <i>Hypoplectrus nigricans</i> | Serranidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 50.0 | 3 | 0.5 | 0.5 | 10.0 | 9 | 11 | | |
| barred hamlet | <i>Hypoplectrus puella</i> | Serranidae | 33.3 | 3 | 0.5 | 0.8 | 13.5 | 10 | 15 | 16.7 | 1 | 0.2 | 0.4 | 10.0 | 10 | 10 | | |
| harlequin bass | <i>Serranus tigrinus</i> | Serranidae | 16.7 | 1 | 0.2 | 0.4 | 10.0 | 10 | 10 | 33.3 | 2 | 0.3 | 0.5 | 7.0 | 6 | 8 | | |
| sand diver | <i>Synodus intermedius</i> | Synodontidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 1 | 0.2 | 0.4 | 32.0 | 32 | 32 | | |
| sharpnose puffer | <i>Canthigaster rostrata</i> | Tetraodontidae | 33.3 | 3 | 0.5 | 0.8 | 4.2 | 3 | 5 | 50.0 | 6 | 1.0 | 1.3 | 4.5 | 4 | 6 | | |
| porcupinefish | <i>Diodon hystrix</i> | Tetraodontidae | 0.0 | 0 | 0.0 | 0.0 | - | - | - | 16.7 | 2 | 0.3 | 0.8 | 35.0 | 35 | 35 | | |
| | | | Total= | 4326 | | | | | | | Total= | 3319 | | | | | | |