

The Wind-Field Structure of the Tropical Cyclone Boundary-Layer

Thesis submitted for the degree of

Doctor of Philosophy

by

Jeffrey David Kepert

BSc (Honours), University of Western Australia

MSc, Monash University

Department of Mathematics and Statistics

Monash University

November 2002

This thesis contains no material which has been accepted for the award of any other degree or diploma in any university or other institution, and, to the best of my knowledge, contains no material previously published or written by another person, except where due reference is made in the text.

The work described in chapter 2 (with the exception of section 2.5) was published as:

Kepert, J.D., 2001: The dynamics of boundary layer jets within the tropical cyclone core. Part I: Linear theory. *J. Atmos. Sci.*, **58**, 2469-2484.

The work described in chapter 3 was published as:

Kepert, J.D. and Y. Wang, 2001: The dynamics of boundary layer jets within the tropical cyclone core. Part II: Nonlinear enhancement. *J. Atmos. Sci.*, **58**, 2485-2501.

Dr. Wang's contribution to that paper was in formulating and coding the dynamics portion of the model, writing that section of the paper, and in some discussions during the analysis of the results. The remainder of the model was coded and the analysis performed by myself.

Jeff Kepert

November 29, 2002.

Acknowledgements

I am grateful to my supervisors, Dr Greg Holland and Dr Dale Hess, for suggesting the research topic, and for their encouragement and many helpful suggestions along the way, and for their guidance in refining the write-up. I also acknowledge the efforts by Dr Mike Montgomery and an anonymous reviewer in reviewing the published version of Chapters 2 and 3. Their suggestions resulted in significant improvements in the work. My employer, the Bureau of Meteorology, allowed me paid leave for a part of this work, and to extensively use their computer facilities for it. I am particularly grateful to my immediate bosses, Dr Mike Manton and Dr Bill Bourke, for their encouragement and understanding.

The analyses of Hurricanes Georges and Mitch in Chapter 4 would have been impossible without access to the high quality data obtained by the Hurricane Research Division (HRD) of the U.S. National Oceanic and Atmospheric Administration. I am full of admiration for the hard work and dedication of the flight and scientific crews in obtaining these data, and gratefully acknowledge the generosity of the HRD in making it so freely available. The Bureau of Meteorology Research Centre observing station at North West Cape, which took the observations in Severe Tropical Cyclone Vance, is partly supported by the U.S. Office of Naval Research.

I was fortunate to attend several of the American Meteorology Society's Conferences on Hurricanes and Tropical Meteorology, with support from the U.S. Office of Naval Research and the Bureau of Meteorology. The experience and interactions helped significantly to advance the research.

I would also like to thank various colleagues for helpful comments,

encouragement and advice. In particular, Dr James Franklin of the U.S. National Hurricane Center advised me on several technical aspects of the GPS dropsonde, and read and commented on an early draft of the analysis of Hurricane Mitch. Dr Mark Powell of HRD helped in understanding the intricacies of his vertical wind profile model, and with positive comments on other aspects of this work. Dr Hugh Willoughby of HRD provided pre-publication access to and code for his parametric radial wind profile, and advice on its use. Dr Yuqing Wang of the University of Hawaii wrote the dynamics portion of the numerical model in Chapter 3, and provided advice on a range of numerical issues. Barry Hanstrum of the Bureau of Meteorology's Western Australian Regional Forecasting Centre advised on the characteristics of sheared tropical cyclones.

Finally, I would like to express my deepest gratitude to my wife Karen, and children Amanda, Emily and Simon, for their patience with and understanding for the many nights and weekends I have spent working on this thesis. I also thank them for the occasional but timely reminders that there are other things in life. Completing this work as a part-time student would not have been possible without their support, and in acknowledgment of their contribution, I dedicate this thesis to them.

Abstract

The recent massive increase in observations within the tropical cyclone boundary-layer has emphasised that a low-level wind speed maximum, or jet, commonly occurs there. Two complementary tools are developed here to diagnose the boundary-layer winds in a translating tropical cyclone: a linear analytical model, and a high-resolution numerical model with sophisticated parameterisations of the sea-air transfers and turbulence.

The solution to the linear model has three components: a symmetric one due to the cyclone, and two asymmetric ones which result from the interaction of the moving cyclone with the earth's surface. The asymmetric components are shown to be frictionally stalled inertia waves, while all three components are modified Ekman spirals. It is argued that this simple Ekman-type model may be appropriate in tropical cyclones since diurnal effects are weak or absent, turbulence is predominantly shear-generated, and baroclinicity is weak.

The jet is similar to the supergradient flow found at the top of the classical Ekman spiral. It is only a few percent supergradient in the linear model, because the neglect of vertical advection there substantially reduces the strength. The jet height scales as $(2K/I)^{1/2}$, where K is the turbulent diffusivity and I the inertial stability, is typically several hundreds of metres in the cyclone core, and increases with radius. In a moving storm, the jet is most supergradient – several times stronger than in a stationary storm – at the eyewall to the left (right) and front of the storm in the Northern (Southern) Hemisphere, and extends into a significant area around to the left (right) of the storm. It is, however, much less marked to the right (left), where the strongest near-surface winds are found. The ratio of near-surface to gradient-wind speed is shown to have a substantial spatial

variability. Larger values are found near the eye, and there is a marked increase from right to left (left to right) of the track of a moving storm.

The second tool used is a high resolution, full primitive equations, dry, hydrostatic, numerical model. It relaxes the constraint of linearity and includes sophisticated physical parameterisations. Strong inwards advection of angular momentum produces a strong jet, typically from 10% to 25% supergradient near the radius of maximum winds (RMW). The inflow is maintained against the outwards acceleration resulting from gradient-wind imbalance in a stationary vortex mainly by vertical diffusion and vertical advection. In a moving cyclone, horizontal advection also becomes important. The jet height, motion-induced asymmetries, and spatial variability of the surface-wind factor compare well in the two models.

Predictions from the modelling work are tested against observations in five tropical cyclones. Hurricane Mitch (1998) had strongly azimuthal-mean supergradient flow near the RMW from 400-m to 2-km height, while the flow in Hurricane Georges (1998) was apparently close to balanced. This difference may be because Georges was commencing an eyewall-replacement cycle. The asymmetric part of the near-eyewall flow in both these storms is shown to have a similar structure to the frictionally stalled inertia wave and numerical simulations, although the frictional asymmetry which forces this is provided in Mitch by proximity to land rather than by motion. Severe Tropical Cyclone Vance (1999) also showed a large area of supergradient flow at about 1.5 km height ahead of and to the right of the storm, the strength and height of which was consistent with numerical simulations.

The observed ratio of near-surface wind speed to the wind aloft is also shown to

be in good agreement with the models. In particular Hurricanes Hugo (1989), Andrew (1992) and Georges displayed an increase of this ratio towards the centre of the storm, and higher values on the left of the track than on the right. Hurricane Mitch had an increase towards the centre, but different asymmetry. The observed ratio was lower than modelled in the outer core of Tropical Cyclone Vance. The agreement could be improved by increasing the ocean-surface roughness in the model to account for the shallow ocean and limited fetch in this case.

Contents

1. Introduction	1
1.1. Opening Remarks	1
1.2 Observed Boundary Layer Wind Structure	6
1.2.1 Horizontal wind structure	6
1.2.2 Observational Comparisons of Boundary Layer and Surface Winds	26
1.2.3 Observed Vertical Profiles of Boundary-Layer Wind	32
1.2.4 Coherent structures in the tropical cyclone boundary layer ..	38
1.2.5 Final remarks on observational studies	39
1.3 Theoretical and Modelling Studies of Tropical Cyclone Boundary Layer Wind Structure	42
1.3.1 The Ekman Spiral	43
1.3.2 Axisymmetric models of tropical cyclone boundary-layer structure	46
1.3.3 The horizontal structure	52
1.3.4 The Deardorff-Moss-Rosenthal-Powell Wind Profile Model .	55
1.3.5 Final remarks on theoretical and modelling studies	61
1.4 Hypothesis and Scope of this Study	63
2. Linear Theory	69
2.1 Introduction	69
2.2 Derivation of the Linear Model	71
2.3 The boundary layer of a stationary vortex	80
2.3.1 Horizontal Flow	80

2.3.2 Vertical velocity forcing in a stationary vortex	85
2.3.3 Surface wind reduction	86
2.3.4 Application to a typical cyclone	87
2.3.5 The location of the maximum updraft	90
2.4 The boundary layer in a moving cyclone	96
2.4.1 Physical interpretation of the asymmetries	100
2.5 The impact of vertical advection on the jet strength	104
2.6 Concluding Remarks	112
 3. Numerical Modelling	 117
3.1 Introduction	117
3.2 Model Formulation	120
3.2.1 Governing equations	121
3.2.2 Physical parameterisations	122
3.2.3 Numerical Method	124
3.2.4 Initial and Boundary Conditions	126
3.3 The Boundary Layer of a Symmetric Storm	130
3.3.1 The Inertially Stable, Moderately Intense Case	130
3.3.2 Height scales in Storm I	142
3.3.3 Comparison to the nocturnal jet	144
3.3.4 An Inertially Stable, Intense Storm	146
3.3.5 The Inertially Neutral Case	148
3.4 The Effect of Cyclone Movement	156
3.4.1 Boundary Layer Flow in a Translating Cyclone	156
3.5 Summary and Conclusions	169

4. Comparison with Observations	173
4.1. Introduction	173
4.2 Hurricane Georges	177
4.2.1 Introduction and Data Coverage	177
4.2.2 Cyclone Track	186
4.2.3 The Wind Field	194
4.2.3.1 The Winds Near the Radius of Maximum Winds	194
4.2.3.2 The Winds in the Eye	200
4.2.3.3 The Winds in the Outer Core	200
4.2.3.4 Surface Wind Factor	206
4.2.4 Analysis of Balance	209
4.2.4.1 Gradient Wind Equation I: Pressure Analysis	216
4.2.4.2 Gradient Wind Equation II: Wind Analysis	225
4.2.5 Model Simulations	226
4.2.6 Discussion	232
4.3 Hurricane Mitch	236
4.3.1 Introduction and Data Coverage	236
4.3.2 Cyclone Track	243
4.3.3 The Wind Field	244
4.3.3.1 Inner Core	246
4.3.3.2 The Winds in the Eye	253
4.3.3.3 The Winds in the Outer Core	253
4.3.3.4 Surface-Wind Factor	253
4.3.4 Analysis of Balance	256
4.3.4.1 Hydrostatic Integrations	256
4.3.4.1 Gradient Wind Equation I: Pressure Analysis	260

4.3.4.2 Gradient Wind Equation I: Wind Analysis	264
4.3.5 Model Simulations	268
4.3.6 Discussion	272
4.4 Hurricane Andrew	276
4.5 Hurricane Hugo	282
4.6 Severe Tropical Cyclone Vance	285
4.6.1 Introduction	285
4.6.3 Mean Wind Data	288
4.6.4 Gradient Balance Analysis	290
4.6.5 Surface Wind Reduction	292
4.6.6 Modelling	294
4.6.6 Discussion	294
4.7 Concluding Remarks	300
Appendix 4.A: Some Mathematical Details	304
4.A1 The Holland Parametric Profile	304
4.A2 The Willoughby Parametric Profile	306
4.A3 Some Notes on the Levenberg-Marquadt Fitting Procedure	314
5 Summary and Conclusions	319
5.1 The Hypothesis Revisited	319
5.2 Theory, Modelling and Observations	323
5.3 Closing Remarks	335
6 Bibliography	337