

1 **Is there an optimal ENSO pattern that increases U.S. tornado activity?**

2
3
4
5
6 Sang-Ki Lee^{1,2}, David Enfield^{1,2}, Hailong Liu^{1,2}, Chunzai Wang², Robert Atlas², and Brian
7 Mapes³

8 ¹Cooperative Institute for Marine and Atmospheric Studies, University of Miami, Miami,
9 Florida, USA

10 ²Atlantic Oceanographic and Meteorological Laboratory, NOAA, Miami Florida, USA
11 USA

12 ³Rosenstiel School of Marine and Atmospheric Science, University of Miami, Miami, Florida,
13 USA

14
15
16
17 June 2011

18
19
20
21
22 Corresponding author address: Dr. Sang-Ki Lee, NOAA/AOML, 4301 Rickenbacker Causeway,
23 Miami, FL 33149, USA. E-mail: Sang-Ki.Lee@noaa.gov.

1 **Abstract**

2 Observations and modeling experiments are used to show that a positive phase of Trans-
3 Niño, characterized by cooling in the central tropical Pacific and warming in the eastern tropical
4 Pacific, is linked to an increased number of intense U.S. tornadoes in spring. The warming in the
5 eastern tropical Pacific increases convection locally, but also contributes to suppressing
6 convection in the central tropical Pacific. This in turn works constructively with cooling in the
7 central tropical Pacific to force a strong and persistent negative phase Pacific – North American
8 (PNA)-like teleconnection pattern. The anomalous winds that are associated with this
9 teleconnection pattern bring more cold and dry upper-level air from the high-latitudes and more
10 warm and moist lower-level air from the Gulf of Mexico converging into the central U.S., and
11 thus provide a favorable condition for increased U.S. tornado activity. A distinctive feature in the
12 2011 Trans-Niño event is warming in the western tropical Pacific that further aided to suppress
13 convection in the central tropical Pacific and thus contributed to strengthening the teleconnection
14 response in the central U.S. in favor of increased U.S. tornado activity.

15
16
17
18
19
20
21
22
23

1 **1. Introduction**

2 In April and May of 2011, a record breaking 1,243 tornadoes were reported in the United
3 States. This resulted in 517 tornado related fatalities, making 2011 one of the deadliest tornado
4 years in U.S. history [<http://www.spc.noaa.gov/climo/torn/fataltorn.html>]. Questions were raised
5 almost immediately as to whether the series of extreme tornado outbreaks in 2011 could be
6 linked to long-term climate changes. The severe weather database (SWD) from the National
7 Oceanic and Atmospheric Administration indicates that the number of total U.S. tornadoes (i.e.,
8 from F0 to F5 in the Fujita-Pearson scale) during the most active tornado months of April and
9 May (AM) has been steadily increasing since 1950 (Figure S1a). However, due to the
10 improvements in tornado detection technology with time, one must be cautious in attributing this
11 secular increase in the number of U.S. tornadoes to a specific long-term climate signal [Brooks
12 and Doswell, 2001]. Since intense and long-lived tornadoes are much more likely to be detected
13 and reported even before a national network of Doppler radar was build in the 1990s, the number
14 of intense U.S. tornadoes (i.e., from F3 to F5 in the Fujita-Pearson scale) in AM during 1950-
15 2010 is obtained from the SWD (Figure S1b) and used, after detrending, as the primary
16 diagnostic index in this study (Figure S1c).

17 In the central U.S. east of the Rocky Mountains, cold and dry upper-level air from the high
18 latitudes often converges with warm and moist lower-level air coming from the Gulf of Mexico
19 (GoM). Due to this so-called large-scale differential advection (i.e., two or more different air
20 masses converging at different heights), a conditionally unstable atmosphere with high
21 convective available potential energy is formed that causes frequent and intense thunderstorms.
22 With the addition of a triggering mechanism, such as the horizontal spinning effect provided by
23 the lower-level wind shear (i.e., wind speed increasing or wind direction changing with height),

1 these thunderstorms can spawn intense tornadoes. Consistently, the moisture transport from the
2 GoM to the central U.S. is significantly correlated with the number of intense U.S. tornadoes in
3 AM (see Table 1). See Brooks et al. [2003] for various meteorological indices used to estimate
4 the occurrence of tornadoes.

5 The Pacific – North American (PNA) pattern in boreal winter and spring is linked to the
6 large-scale differential advection in the central U.S. as discussed in earlier studies [e.g., Munoz
7 and Enfield, 2011]. During a negative phase of the PNA, an anomalous cyclone is formed over
8 North America that bring more cold and dry upper-level air from the high latitudes to the central
9 U.S., and an anomalous anticyclone is formed over the southeastern seaboard that increases the
10 southwesterly wind from the GoM to the central U.S., thus enhancing the Gulf-to-U.S. moisture
11 transport. Although the PNA is a naturally occurring atmospheric phenomenon driven by
12 intrinsic variability of the atmosphere, a La Niña in the tropical Pacific can project onto a
13 negative phase PNA pattern [e.g., Straus and Shukla 2002]. In addition, since the Gulf-to-U.S.
14 moisture transport can be enhanced with a warmer GoM, the sea surface temperature (SST)
15 anomaly in the GoM can also affect U.S. tornado activity. During the decay phase of La Niña in
16 spring, the GoM is typically warmer than usual [e.g., Alexander and Scott 2002]. Therefore, the
17 Gulf-to-U.S. moisture transport could be increased during the decay phase of La Niña in spring
18 due to the increased SSTs in the GoM and the strengthening of the southwesterly wind from the
19 GoM to the U.S. Nevertheless, none of these (i.e., PNA, GoM SST, and La Nina) are
20 significantly correlated with the number of intense tornadoes in AM (see Table 1) as discussed in
21 earlier studies [e.g., Cook and Schaefer, 2008]. Currently, seasonal forecast skill for intense U.S.
22 tornado outbreaks, such as occurred in 2011, has not been demonstrated.

1 Among the long-term climate patterns considered in Table 1, only the Trans-Niño (TNI) is
2 significantly correlated ($r = 0.33$) with the number of intense U.S. tornadoes in AM. The TNI,
3 which is defined as the difference in normalized SST anomalies between the Niño-1+2 (10° -
4 0° ; 90° W - 80° W) and Niño-4 (5° N - 5° S; 160° E - 150° W) regions, represents the evolution of
5 the El Niño-Southern Oscillation (ENSO) in the months leading up to the event and the
6 subsequent evolution with opposite sign after the event [Trenberth and Stepaniak, 2001]. Given
7 that AM is typically characterized with the development or decay phase of ENSO events, it is
8 more likely that the tropical Pacific SST anomalies in AM are better represented by the TNI
9 index than the conventional ENSO indices such as Niño-3.4 (5° N - 5° S; 170° W - 120° W) or
10 Niño-3 (5° N - 5° S; 150° W- 90° W). Nevertheless, it is not at all clear why the number of intense
11 U.S. tornadoes in AM is significantly correlated with the TNI index, but not with other ENSO
12 indices. This is the central question that we explore in the following sections by using both
13 observations and an atmospheric general circulation model (AGCM).

14

15 **2. Observations**

16 To better understand the potential link between the TNI and U.S. tornado activity, we ranked
17 the years from 1950 to 2010 (61 years in total) based on the number of intense U.S. tornadoes in
18 AM. The top ten years are characterized by increased Gulf-to-U.S. moisture transport in the
19 lower-level (Figure 1a) and an anomalous upper-level cyclone over North America that advects
20 more cold and dry air to the central U.S. (Figure 1c), whereas the bottom ten years are associated
21 with decreased Gulf-to-U.S. moisture transport (Figure 1b) and an anomalous upper-level
22 anticyclone over North America (Figure 1d). This is consistent with the notion that springtime
23 U.S. tornado activity is modulated by the large-scale differential advection in the central U.S.

1 Among the top ten years (Table S1), seven years including the top three are identified with a
2 positive phase (i.e., above $\frac{1}{4}$ quantile) TNI index (i.e., normalized SST anomalies are larger in
3 the Niño-1+2 than in Niño-4 region). Five out of those seven years are characterized by a La
4 Niña transitioning to a different phase or persisting beyond AM (1957, 1965, 1974, 1999, and
5 2008) and the other two with an El Niño transitioning to either a La Nina or neutral phase (1983
6 and 1998). Figure 2a shows the composite SST anomalies for those five positive phase TNI years
7 transitioning from a La Niña.

8 On the other hand, among the bottom ten years (Table S2), only one year is identified with a
9 positive phase TNI, and the other nine years are with a neutral phase (i.e., between $\frac{1}{4}$ and $\frac{3}{4}$
10 quantile) TNI, suggesting that a negative phase of the TNI does not either decrease or increase
11 the number of intense U.S. tornadoes in AM. Interestingly, four years among the bottom ten
12 years are identified with a La Niña transitioning to a different phase or persisting beyond AM
13 (1950, 1951, 1955 and 2001), and four are with an El Niño transitioning to a different phase or
14 persisting beyond AM (1958, 1987, 1988 and 1992). As shown in Figure 2b, the composite SST
15 anomaly pattern for the four years of the bottom ten years with a La Niña transitioning is that of
16 a typical La Niña with the SST anomalies in the Niño-4 and Niño-1+2 being both strongly
17 negative (i.e., neutral phase TNI). Similarly, as shown in Figure 2c, the composite SST anomaly
18 pattern for the four years in the bottom ten years with an El Niño transitioning is that of a typical
19 El Niño with the SST anomalies in the Niño-4 and Niño-1+2 being both strongly positive (i.e.,
20 neutral phase TNI).

21 In summary, observations indicate that a positive phase TNI (i.e., normalized SST anomalies
22 are larger in the Niño-1+2 than in Niño-4 region) is linked to an increased number of intense
23 U.S. tornadoes in AM, whereas both La Niñas and El Niños with a neutral phase TNI (i.e., the

1 SST anomalies in the Niño-1+2 region are as strong and the same sign as the SST anomalies in
2 the Niño-4) are linked to a decreased number of intense U.S. tornadoes in AM.

3

4 **3. Model Experiments**

5 To explore the potential link between the three tropical Pacific SST anomaly patterns (see
6 Figure 2), identified in the previous section, and the number of intense U.S. tornadoes in AM, a
7 series of AGCM experiments are performed by using version 3.1 of the NCAR community
8 atmospheric model coupled to a slab mixed layer ocean model (CAM3). The model is a global
9 spectral model with a triangular spectral truncation of the spherical harmonics at zonal wave
10 number 42 (T42). It is vertically divided into 26 hybrid sigma-pressure layers. Model
11 experiments are performed by prescribing various composite evolutions of SSTs in the tropical
12 Pacific region (15°S–15°N; 120°E-coast of the Americas) while predicting the SSTs outside the
13 tropical Pacific using the slab ocean model. To prevent discontinuity of SST around the edges of
14 the forcing region, the model SSTs of three grid points centered at the boundary are determined
15 by combining the simulated and prescribed SSTs. Each ensemble consists of ten model
16 integrations that are initialized with slightly different conditions to represent intrinsic
17 atmospheric variability. The same methodology was previously used in Lee et al. [2008] for
18 studying ENSO teleconnection to the tropical North Atlantic region.

19 Four sets of ensemble runs are performed (Table S3). In the first experiment (EXP_CLM),
20 the SSTs in the tropical Pacific region are prescribed with climatological SSTs. In the second
21 experiment (EXP_TNI), the composite SSTs of the positive phase TNI years identified among
22 the ten most active U.S. tornado years are prescribed in the tropical Pacific region. Note that only
23 the five positive TNI years transitioning from a La Niña (1957, 1965, 1974, 1999, and 2008) are

1 considered here because the other two positive TNI years are transitioning from an El Niño
2 (1983 and 1998) and thus tend to cancel the tropical SST anomalies of the other five. In the next
3 two experiments, the SSTs in the tropical Pacific region are prescribed with the composite SSTs
4 of the four years in the bottom ten years with a La Niña transitioning (1950, 1951, 1955 and
5 2001) for EXP_LAN, and the four years in the bottom ten years with an El Niño transitioning
6 (1958, 1987, 1988 and 1992) for EXP_ELN.

7

8 **4. Model Results**

9 Figure 3 and S2 show the difference in moisture transport, and geopotential height and wind
10 at 500 hPa in AM obtained by subtracting those fields in EXP_CLM from the three model
11 experiments. In EXP_TNI (Figure 3), the Gulf-to-U.S. moisture transport is greatly increased
12 and an anomalous upper-level cyclone is formed over North America that brings more cold and
13 dry air to the central U.S., all indicating enhanced large-scale differential advection in the central
14 U.S. In EXP_LAN and EXP_ELN (Figure S2), on the other hand, the Gulf-to-U.S. moisture
15 transport is neither increased nor decreased. A relatively weak anomalous upper-level cyclone is
16 formed in EXP_LAN, but it is positioned over the eastern seaboard. In EXP_ELN, a weak
17 anomalous upper-level anticyclone is formed and also positioned over the eastern seaboard.

18 Therefore, these model results fully support the hypothesis that a positive phase of the TNI
19 with cooling in the central tropical Pacific (CP) and warming in the eastern tropical Pacific (EP)
20 enhances the large-scale differential advection in the central U.S. advecting more cold and dry
21 upper-level air from the high-latitudes and more warm and moist lower-level from the GoM, and
22 thus increasing U.S. tornado activity in spring. However, the model results do not show any
23 increase or decrease in the large-scale differential advection in the central U.S. under La Niña

1 and El Niño conditions as long as the SST anomalies in EP are as strong and the same sign as the
2 SST anomalies in CP.

3

4 **5. CP- versus EP-forced Teleconnection**

5 The model results strongly suggest that cooling in CP and warming in EP may have a
6 constructive influence on the teleconnection pattern that strengthens the large-scale differential
7 advection over the central U.S. To better understand how the real atmosphere with moist diabatic
8 processes responds to CP cooling and EP warming, two sets of additional model experiments
9 (EXP_CPC and EXP_EPW) are performed (Table S3). These two experiments are basically
10 identical to EXP_TNI except that the composite SSTs of the positive phase TNI years are
11 prescribed only in the western and central tropical Pacific region (15°S–15°N; 120°E - 110°W)
12 for EXP_CPC and only in the eastern tropical Pacific region (15°S–15°N; 110°W-coast of the
13 Americas) for EXP_EPW.

14 Figure 4 show the difference in moisture transport, and geopotential height and wind at 500
15 hPa in AM obtained by subtracting those fields in EXP_CLM from the two additional model
16 experiments. In EXP_CPC, the teleconnection pattern emanating from the tropical Pacific
17 consists of an anticyclone over the Aleutian Low in the North Pacific, a cyclone over North
18 America, and an anticyclone over the southeastern U.S. and central Americas, consistent with a
19 typical negative phase PNA pattern (Figure 4c). As expected from the anomalous anticyclone
20 over the southeastern U.S. and central Americas, the Gulf-to-U.S. moisture transport is increased
21 in EXP_CPC (Figure 4b).

22 Surprisingly, the Rossby wave train forced by warming in EP (EXP_EPW) is very similar to
23 that in EXP_CPC (Figure 4f). Consistently, the Gulf-to-U.S. moisture transport is also increased

1 in EXP_EPW as in EXP_CPC and EXP_TNI (Figure 4e). A question arises as to why the
2 teleconnection pattern forced by warming in EP is virtually the same as that forced by cooling in
3 CP (see Figure 4c and 4f). It appears that the Rossby wave train in EXP_EPW is not directly
4 forced from EP. In EXP_EPW, convection is increased locally in EP, but it is decreased in CP as
5 in EXP_CPC (Figure S3c). This suggests that increased convection in EP associated with the
6 increased local SSTs suppresses convection in CP that in turn forces a negative phase PNA-like
7 pattern. Therefore, these model results confirm that cooling in CP and warming in EP do have
8 constructive influence on the teleconnection pattern that strengthens the large-scale differential
9 advection over the central U.S. The model results also suggest that cooling in CP with neutral
10 SST anomalies in EP or warming in EP with neutral SST anomalies in CP can strengthen the
11 large-scale differential advection over the central U.S.

12 An apparently important question is why warming in EP does not directly excite a Rossby
13 wave train to the high-latitudes. As shown in earlier theoretical studies, the vertical background
14 wind shear is one of the two critical factors, which are required for tropical heating to radiate
15 barotropic teleconnections to the high-latitudes [e.g., Lee et al. 2009]. In both observations and
16 EXP_CLM, the background vertical wind shear between 200 and 850 hPa in AM is largest in the
17 central tropical North Pacific and smallest in EP and the western tropical Pacific (WP), providing
18 an explanation as to why the Rossby wave train in EXP_EPW is not directly forced in EP
19 (Figure S4). See Lee et al. [2009] and references therein for more discussions on this issue.

20

21 **6. Effect of Intrinsic Variability**

22 The conclusion so far is that a positive phase TNI with cooling in CP and warming in EP can
23 increase U.S. tornado activity in spring because CP cooling and EP warming have constructive

1 influence on the teleconnection pattern that strengthens the large-scale differential advection
2 over the central U.S. However, the correlation between the TNI and the number of intense U.S.
3 tornadoes in AM is only 0.33 indicating that the TNI can explain only about 10% of the total
4 variance. This suggests that intrinsic variability in the atmosphere may overwhelm the TNI-
5 teleconnection pattern over North America as discussed in earlier studies for El Niño-
6 teleconnection patterns in the Pacific–North American region [e.g., Hoerling and Kumar, 1997].
7 In order to better illustrate this point, we select neutral TNI years during 1950-2010 and divide
8 them into one group with an increased Gulf-to-U.S. moisture transport in AM and the other
9 group with a decreased Gulf-to-U.S. moisture transport. Figure S5b shows the difference in
10 moisture transport between these two groups. Similarly, we divide ten ensemble experiments for
11 EXP_CLM into two groups, one with an increased Gulf-to-U.S. moisture transport in AM and
12 the other group with a decreased Gulf-to-U.S. moisture transport. Figure S5c shows the
13 difference in moisture transport between these two groups.

14 As expected, the typical amplitude of Gulf-to-U.S. moisture transport induced by intrinsic
15 variability in the atmosphere such as the PNA during neutral phase TNI years is larger than the
16 anomalous Gulf-to-U.S. moisture transport for positive phase TNI years (Figure S5a). Similarly,
17 internal variability in EXP_CLM can generate anomalous Gulf-to-U.S. moisture transport with
18 comparable amplitude to that simulated in EXP_TNI. Therefore, we can conclude that the
19 strengthening of the large-scale differential advection over the central U.S. during a positive
20 phase of the TNI can be greatly reduced or enhanced by internal variability in the atmosphere. In
21 other words, although seven of the ten most active tornado years during 1950-2010 including the
22 top three years are characterized with a strongly positive phase of the TNI (Table S1), the

1 associated predictability of U.S. tornado activity, which can be defined as a ratio of the climate
2 signal (the TNI index in this case) relative to the climate noise, is low.

3

4 **7. U.S. Tornado Outbreaks in 2011**

5 A positive phase of the TNI prevailed during AM of 2011 with cooling in CP and warming
6 EP (Figure S6a). An important question is whether the series of extreme U.S. tornado outbreaks
7 during AM of 2011 can be attributed to this positive of the TNI. During AM of 2011, an
8 anomalous upper-level anticyclone was formed over North America (Figure S6c), and the Gulf-
9 to-US moisture was greatly increased (Figure S6b), both indicating the coherent teleconnection
10 response to a positive phase of the TNI. To confirm this, a set of model experiments (EXP_011)
11 is performed by prescribing the SSTs for 2010 - 2011 in the tropical Pacific region while
12 predicting the SSTs outside the tropical Pacific using the slab ocean model (Table S3). As
13 summarized in Figure S7, the model results are consistent with the observations, although the
14 anomalous Gulf-to-US moisture is weaker in the model experiment.

15 A distinctive feature in the 2011 TNI event is warming in WP (Figure S6a). Further
16 experiments (Table S3) suggest that the warming in WP indirectly suppresses convection in CP,
17 and thus works constructively with the cooling in CP to force a strong and persistent negative
18 phase PNA-like pattern (Figure S8 and S9). Thus, despite the low signal-to-noise ratio in the
19 TNI-teleconnection response in the central U.S., it is highly likely that the 2011 positive phase
20 TNI event did contribute to the increased number of intense U.S. tornadoes in AM of 2011 by
21 enhancing the differential advection in the central U.S.

22

23 **8. Discussion**

1 The record-breaking U.S. tornado outbreak in the spring of 2011 prompts the need to identify
2 and understand long-term climate signals that provide seasonal predictability for intense tornado
3 outbreaks. Here, we use both observations and modeling experiments to demonstrate that a
4 positive phase of the TNI, characterized by cooling in CP and warming in EP, is one such
5 climate pattern that can strengthen the large-scale differential advection in the central U.S. and
6 thus increase U.S. tornado activity. However, although seven of the ten most active tornado years
7 during 1950 – 2010 including the top three are characterized with a strongly positive phase of the
8 TNI, the TNI explains only 10% of the total variance in the number of intense U.S. tornadoes in
9 AM. Therefore, further study is needed to refine the predictive skill provided by the TNI and to
10 explore other long-term climate signals that can provide additional predictability in seasonal and
11 longer time scales. It is important to point out some limitations in this study. In particular, the
12 atmospheric model used as the main tool in this study (CAM3) has a relatively low-resolution
13 (T42), which may be subject to large model bias. Obviously, more rigorous multi-model
14 experiments with various horizontal resolutions are required to test if the major conclusions of
15 this study are not model-dependent.

16

17 **Acknowledgments.** This study was motivated and benefited from interactions with scientists at
18 NOAA ESRL, GFDL, CPC, and NCDC. In particular, we wish to thank Wayne Higgins, Tom
19 Karl, Marty Hoerling, and Harold Brooks for initiating and leading discussions that motivated
20 this study. This work was supported by grants from the National Oceanic and Atmospheric
21 Administration’s Climate Program Office and by grants from the National Science Foundation.

22

23

References

- 1 Alexander, M., and J. Scott (2002), The influence of ENSO on air-sea interaction in the Atlantic.
2 *Geophys. Res. Lett.*, **29**, 1701. doi:10.1029/2001GL014347.
- 3 Brooks, H. E., C. A. Doswell III (2001), Some aspects of the international climatology of
4 tornadoes by damage classification, *Atmos. Res.*, **56**, 191– 201.
- 5 Brooks, H. E., J. W. Lee, and J. P. Cravenc (2003), The spatial distribution of severe
6 thunderstorm and tornado environments from global reanalysis data, *Atmos. Res.*, **67-68**, 73-
7 94.
- 8 Cook, A. R., and J. T. Schaefer (2008), The relation of El Niño–Southern Oscillation (ENSO) to
9 winter tornado outbreaks, *Mon. Wea. Rev.*, **136**, 3121–3137.
- 10 Hoerling, M. P., and A. Kumar (1997), Why do North American climate anomalies differ from
11 one El Niño event to another?, *Geophys. Res. Lett.*, **24**, 1059-1062.
- 12 Lee, S.-K., D. B. Enfield and C. Wang (2008), Why do some El Ninos have no impact on
13 tropical North Atlantic SST? *Geophys. Res. Lett.*, **35**, L16705, doi:10.1029/2008GL034734.
- 14 Lee, S.-K., C. Wang and B. E. Mapes (2009), A simple atmospheric model of the local and
15 teleconnection responses to tropical heating anomalies. *J. Clim.*, **22**, 272-284.
- 16 Munoz, E., and D. Enfield (2010), The boreal spring variability of the Intra-Americas low-level
17 jet and its relation with precipitation and tornadoes in the eastern United States, *Clim. Dyn.*
18 **36**, 247–259.
- 19 Straus, D. M., and J. Shukla (2002), Does ENSO force the PNA?, *J. Clim.*, **15**, 2340–2358.
- 20 Trenberth, K. E., and D. P. Stepaniak (2001), Indices of El Niño evolution, *J. Clim.*, **14**, 1697–
21 1701.

22


23

1 **Table 1.** Correlation coefficients of various long-term climate patterns with the number of
 2 intense (F3 - F5) tornados in April and May (AM) during 1950-2010. All indices including the
 3 tornado index are detrended. The SWD, ERSST3, and NCEP-NCAR reanalysis are used to
 4 obtain the long-term climate indices used in this table. Any correlation value with above the 95%
 5 significance is in bold^a

Index	DJF	FMA	AM
Gulf-to-U.S. moisture transport	0.08	0.20	0.40
GoM SST	0.15	0.21	0.20
Niño-4	-0.22	-0.20	-0.19
Niño-3.4	-0.13	-0.13	-0.11
Niño-1+2	0.02	0.11	0.15
TNI	0.28	0.29	0.33
PNA	-0.05	-0.10	-0.20
PDO	-0.12	-0.10	-0.14
NAO	-0.01	-0.10	-0.18

6 ^aThe Gulf-to-U.S. meridional moisture transport is obtained by averaging the vertically
 7 integrated moisture transport in the region of 25°N - 35°N and 100°W - 90°W. The North
 8 Atlantic Oscillation (NAO) index and the Pacific - North American (PNA) pattern are defined as
 9 the first and second leading modes of Rotated Empirical Orthogonal Function (REOF) analysis
 10 of monthly mean geopotential height at 500 hPa, respectively. The Pacific Decadal Oscillation
 11 (PDO) is the leading principal component of monthly SST anomalies in the North Pacific Ocean
 12 north of 20°N.

NCEP-NCAR Reanalysis: Moisture Flux and GPOT Height at 500hPa (APR-MAY)



1
2 **Figure 1.** Anomalous moisture transport for (a) the ten most active U.S. tornado years and (b)
3 the ten least active U.S. tornado years in AM during 1950-2010 obtained from NCEP-NCAR
4 reanalysis. Anomalous geopotential height and wind at 500 hPa for (c) the ten most active U.S.
5 tornado years and (d) the ten least active U.S. tornado years in AM during 1950-2010 obtained
6 from NCEP-NCAR reanalysis. The unit is $\text{kg m}^{-1}\text{sec}^{-1}$ for moisture transport, m for geopotential
7 height, and m s^{-1} for wind.

8

9


10

11

12

13

ERSST3: SST Anomalies (APR-MAY)




1

2 **Figure 2.** Composite SST anomalies in AM, obtained from ERSST3, for (a) the five positive
3 TNI years transitioning from a La Niña identified among the ten most active U.S. tornado years
4 in AM during 1950-2010, and for (b) the four years with a La Niña transitioning and (c) the four
5 years with an El Niño transitioning identified among the ten least active U.S. tornado years in
6 AM during 1950-2010. Thick black lines in (a) indicate the Niño-4 (5°N - 5°S; 160°E - 150°W)
7 and Niño-1+2 (10S° - 0°; 90°W - 80°W) regions.

8


CAM3: EXP_TNI - EXP_CLM (APR-MAY)



1
2 **Figure 3.** Simulated anomalous (a) SST, (b) moisture transport, and (c) geopotential height and
3 wind at 500 hPa in AM obtained from EXP_TNI - EXP_CLM. The unit is $^{\circ}\text{C}$ for SST, kg m^{-1}
4 sec^{-1} for moisture transport, m for geopotential height, m s^{-1} for wind. Thick black lines in (a)
5 indicate the tropical Pacific region where the model SSTs are prescribed.

6

7



1
 2 **Figure 4.** Simulated anomalous (a) SST, (b) moisture transport, and (c) geopotential height and
 3 wind at 500 hPa in AM obtained from EXP_CPC - EXP_CLM. Simulated anomalous (d) SST,
 4 (e) moisture transport, and (f) geopotential height and wind at 500 hPa in AM obtained from
 5 EXP_EPW - EXP_CLM. The unit is $^{\circ}\text{C}$ for SST, $\text{kg m}^{-1} \text{sec}^{-1}$ for moisture transport, m for
 6 geopotential height, and m s^{-1} for wind. Thick black lines in (a) and (d) indicate the regions
 7 where the model SSTs are prescribed.

Table S1. The total of 61 years from 1950 to 2010 are ranked based on the detrended number of intense U.S. tornadoes in AM. The top ten most active U.S. tornado years are listed with ENSO phase in spring and TNI index in AM for each year. Strongly positive (above $\frac{1}{4}$ quantile) and negative (below $\frac{3}{4}$ quantile) TNI index values are in bold and italic, respectively.

Ranking	Year	ENSO phase in spring	TNI index (detrended)
1	1974	La Niña persists	1.30 (1.48)
2	1965	La Niña transitions to El Niño	1.39 (1.54)
3	1957	La Niña transitions to El Niño	0.57 (0.69)
4	1982	El Niño develops	<i>-1.11 (-0.89)</i>
5	1973	El Niño transitions to La Niña	-0.42 (-0.24)
6	1999	La Niña persists	0.47 (0.75)
7	1983	El Niño decays	1.86 (2.08)
8	2003	El Niño decays	<i>-1.24 (-0.94)</i>
9	2008	La Niña decays	1.41 (1.73)
10	1998	El Niño transitions to La Niña	1.69 (1.97)

Table S2. The total of 61 years from 1950 to 2010 are ranked based on the detrended number of intense U.S. tornadoes in AM. The bottom ten years are listed with ENSO phase in spring and TNI index in AM for each year. Strongly positive (above $\frac{1}{4}$ quantile) and negative (below $\frac{3}{4}$ quantile) TNI index values are in bold and italic, respectively.

Ranking	Year	ENSO phase in spring	TNI index (detrended)
52	1958	El Niño decays	-0.61 (-0.49)
53	1955	La Niña persists	-0.27 (-0.16)
54	2001	La Niña decays	0.21 (0.50)
55	1986	El Niño develops	-0.39 (-0.16)
56	1988	El Niño transitions to La Niña	-0.37 (-0.13)
57	1987	El Niño persists	0.10 (0.34)
58	1992	El Niño decays	0.21 (0.47)
59	1952	Neutral	-0.67 (-0.57)
60	1951	La Niña transitions to El Niño	-0.31 (-0.22)
61	1950	La Niña persists	0.77 (0.86)

Table S3. Prescribed SSTs in the tropical Pacific region for each model experiment. All model experiments are initiated from April of the prior year to December of the modeling year. For instance, in EXP_TNI, the model is integrated for 21 months starting in April using the composite April SSTs of 1956, 1964, 1973, 1998, and 2007.

Experiments	Prescribed SSTs in the tropical Pacific region
EXP_CLM	Climatological SSTs are prescribed in the tropical Pacific region (15°S–15°N; 120°E-coast of the Americas).
EXP_TNI	Composite SSTs of the five positive phase TNI years transiting from a La Niña identified among the ten most active U.S. tornado years (1957, 1965, 1974, 1999, and 2008) are prescribed in the tropical Pacific region.
EXP_LAN	Composite SSTs of the four years with a La Niña transitioning (1950, 1951, 1955 and 2001) identified among the ten least active U.S. tornado years are prescribed in the tropical Pacific region.
EXP_ELN	Composite SSTs of the four years with an El Niño transitioning (1958, 1987, 1988 and 1992) identified among the ten least active U.S. tornado years are prescribed in the tropical Pacific region
EXP_CPC	Same as EXP_TNI except that during that during the modeling year the composite SSTs are prescribed only in the western and central tropical Pacific region (15°S–15°N; 120°E - 110°W).
EXP_EPW	Same as EXP_TNI except that during the modeling year composite SSTs are prescribed only in the eastern tropical Pacific region (15°S–15°N; 110°W-coast of the Americas).
EXP_011	SSTs for 2010-2011 are prescribed in the tropical Pacific region.
EXP_WPW	Same as EXP_011 except that during that during the modeling year SSTs for 2010-2011 are prescribed only in the western Pacific region (15°S–15°N; 120°E - 180°).




Figure S1. The number of (a) total (F0 – F5) and (b) intense (F3 – F5) US tornadoes for the most active tornado months of April and May (AM) during 1950-2010 obtained from SWD. The detrended number of intense U.S. tornadoes in AM, which is the primary diagnostic index used in this study, is shown in (c).




Figure S2. Simulated anomalous (a) SST, (b) moisture transport, and (c) geopotential height and wind at 500 hPa in AM obtained from EXP_LAN – EXP_CLM. Simulated anomalous (d) SST, (e) moisture transport, and (f) geopotential height and wind at 500 hPa in AM obtained from EXP_ELN – EXP_CLM. The unit is $^{\circ}\text{C}$ for SST, $\text{kg m}^{-1} \text{sec}^{-1}$ for moisture transport, m for geopotential height, and m s^{-1} for wind. Thick black lines in (a) and (d) indicate the tropical Pacific region where the model SSTs are prescribed.

CAM3: Convective Precipitation (APR–MAY)




Figure S3. Simulated anomalous convective precipitation rate in AM obtained from (a) EXP_TNI - EXP_CLM, (b) EXP_CPC - EXP_CLM, and (c) EXP_EPW - EXP_CLM. The unit is mm day^{-1} .




Figure S4. Background (climatological) vertical wind shear between 200 and 850 hPa in AM obtained from (a) NCEP-NCAR reanalysis, and (b) EXP_CLM. The unit is m sec^{-1} .

Moisture Flux (APR–MAY)




Figure S5. (a) Anomalous composite moisture transport for positive (above $\frac{1}{4}$ quantile) TNI years during 1950–2010. Neutral TNI years during 1950–2010 are divided into one group with increased Gulf-to-US moisture transport in AM and the other group with decreased Gulf-to-US moisture transport. (b) The difference in moisture transport between these two groups obtained from NCEP-NCAR reanalysis. Similarly, ten ensemble experiments for EXP_CLM are divided into two groups, one with increased Gulf-to-US moisture transport in AM and the other group with decreased Gulf-to-US moisture transport. (c) The difference in moisture transport between these two groups derived from EXP_CLM. The unit is $\text{kg m}^{-1} \text{sec}^{-1}$.




Figure S6. Anomalous (a) SST, (b) moisture transport, and (c) geopotential height and wind at 500 hPa in AM of 2011. The SST is obtained from ERSST3. The moisture transport, geopotential height, and wind are obtained from NCEP-NCAR reanalysis. The unit is °C for SST, kg m⁻¹ sec⁻¹ for moisture transport, m for geopotential height, and m s⁻¹ for wind.

CAM3: EXP_011 - EXP_CLM (APR-MAY)




Figure S7. Simulated anomalous (a) SST, (b) moisture transport, and (c) geopotential height and wind at 500 hPa in AM obtained from EXP_011 - EXP_CLM. The unit is $^{\circ}\text{C}$ for SST, $\text{kg m}^{-1} \text{sec}^{-1}$ for moisture transport, m for geopotential height, m s^{-1} for wind. Thick black lines in (a) indicate the tropical Pacific region where the model SSTs are prescribed.

CAM3: EXP_WPW - EXP_CLM (APR-MAY)




Figure S8. Simulated anomalous (a) SST, (b) moisture transport, and (c) geopotential height and wind at 500 hPa in AM obtained from EXP_WPW - EXP_CLM. The unit is °C for SST, kg m⁻¹ sec⁻¹ for moisture transport, m for geopotential height, m s⁻¹ for wind. Thick black lines in (a) indicate the western Pacific region where the model SSTs are prescribed.

CAM3: Convective Precipitation (APR-MAY)




Figure S9. Simulated anomalous convective precipitation rate in AM obtained from (a) EXP_011 - EXP_CLM, and (b) EXP_WPW - EXP_CLM. The unit is mm day^{-1} .