

**THE RELATIONSHIP BETWEEN ENVIRONMENTAL FACTORS  
AND CORAL BLEACHING AT LEE STOCKING  
ISLAND, BAHAMAS IN 1990**

GEORGE D. DENNIS  
and  
ROBERT I. WICKLUND

[Converted to electronic format by Damon J. Gomez (NOAA/RSMAS) in 2003. Copy available at the NOAA Miami Regional Library. Minor editorial changes were made.]

# THE RELATIONSHIP BETWEEN ENVIRONMENTAL FACTORS AND CORAL BLEACHING AT LEE STOCKING ISLAND, BAHAMAS IN 1990

George D. Dennis and Robert I. Wicklund

Caribbean Marine Research Center  
805 East 46th Place  
Vero Beach, Florida 32963

## ABSTRACT

Following the Caribbean-wide bleaching event of 1987 we instituted monitoring of environmental factors that might contribute to stress on coral reefs at our marine laboratory in the Bahamas. Once again in 1990 extensive bleaching occurred from shallow water to 33 m. Elevated water temperatures were closely associated with the start of this bleaching. Periods of calm wind with little rain and high insolation in the summer preceded these elevated water temperatures. The extensive nature of the bleaching in 1990 may be attributable to the long duration of elevated temperatures, atypical for this site. Density cascading of warm hypersaline water on to the shelf in the summer may be an additional stress on deeper reefs at this location.

## INTRODUCTION

The apparent decline of coral reefs worldwide has prompted the Caribbean Marine Research Center (CMRC) to establish a long-term study of reefs in the Exuma Cays, Bahamas. During the fall of 1987 scientists observed bleached hard and soft corals, sponges, and anemones from shallow water to 95 m (Lang et al., 1988). This Caribbean-wide bleaching event of 1987 was believed to be associated with elevated water temperatures but we could find no long-term in-situ seawater temperature data set anywhere in the Caribbean to address this question. At that time CMRC started monitoring more closely environmental conditions on reefs near our marine laboratory at Lee Stocking Island (LSI), Bahamas. The site is particularly interesting because: 1) coral reefs are well monitored, thus the timing of bleaching can be well documented, 2) there is little anthropogenic contribution to environmental stress (e.g., sewer discharge, heavy metal, or pesticide pollution), 3) turbidity stress is very low due to the lack of river runoff, and 4) a range of depths and habitats are in close proximity to the laboratory facilitating the study and comparison of reef systems. Other unique physical characteristics of the site include: 1) extreme temperature range experienced by shallow-water coral reefs in the Bahamas (18-32°C), and 2) reefs are situated on the edge of a large shallow-water bank that rapidly responds to local meteorological conditions and generates density currents (masses of warm saline water) that might affect deeper reefs on the shelf.

Lang et al. (1988) hypothesized that elevated water temperature in 1987 resulted in coral bleaching and that the cause of this warm water was local calm wind conditions. Other contributing factors proposed include density cascading of warm saline water off the bank and clear skies resulting in synergistic light-related effects. Following the 1987 coral bleaching event, CMRC instituted continuous water temperature monitoring and measurement of meteorological parameters might that influence water temperature. Elevated water temperatures at our site could be due to a regional rise in temperature (i.e., an ENSO event) or simply localized conditions such as calm winds and clear skies or a combination of the two. Past events in the Florida Keys (Jaap, 1979; 1985) and locally (Lang et al., 1988) have suggested that doldrum periods contribute to warm water events resulting in bleaching. Duration of the thermal stress also may be a factor in determining the magnitude of a bleaching event. Here we address the hypothesis that elevated water temperature resulted in the coral bleaching event of 1990 in the Bahamas and identify the potential cause of thermal-related stress on corals.

Visual observations on coral reefs were made by divers on a weekly basis and more frequently after the start of bleaching. Three locations were monitored: Rainbow Garden (on the bank), Perry Reef (shallow fringing reef, 20 m), and Adderley Dropoff (shelf-edge reef, 33 m) (Figure 1). Thermographs, recording water temperature every 30 min, were deployed at several locations on the bank, as well as, at Perry reef and the shelf edge off Adderley Cut (Figure 1)(see Wicklund et al., 1993 for other locations). A General Oceanic current meter with conductivity probe was used to determine salinity in Adderley Cut (1990) and at Adderley Dropoff (1992). A Campbell Scientific meteorological station summarized hourly measurements of air temperature, relative humidity, insolation (pyranometer, cal/cm<sup>2</sup>), rainfall, wind speed and direction from 1990-1992.




Figure 1. Map of Lee Stocking Island and Sampling Locations.

## RESULTS

### Bleaching Observations

Following the massive bleaching event of 1987, little bleaching was observed in 1988 and 1989 and cannot be attributed to any specific environmental cause. Widespread bleaching was again observed in mid-August 1990 at 20-25 m on the shelf-edge fringing reef and involved 1-2% of the reef. By mid-September bleaching was observed in a depth range of 2-46 m. Bleaching at 20 m consisted of about 25% of the reef. Bleached organisms included stony coral, gorgonians, and sponges (see Table 1 for list of species). Bleaching continued to increase in number of colonies affected through the first week of October. Several colonies of *A. cervicornis* and *P. porites* that bleached and survived in 1987 died in 1990. One loggerhead sponge (*Sphaciospongia vesparia*) at 35 m, reduced in size through tissue loss in 1987, bleached again in 1990. No further bleaching was observed in 1991 or 1992.

Table 1. List of species bleached in 1990 at Lee Stocking Island.

|                                |                             |                              |
|--------------------------------|-----------------------------|------------------------------|
| Porifera (sponge)              | Scleractinia (hard coral)   | Gorgonacea (soft coral)      |
| <i>Aplysina</i> spp.           | <i>Acropora cervicornis</i> | several unidentified species |
| <i>Sphaciospongia vesparia</i> | <i>Agaricia agaricites</i>  |                              |
| <i>Xestospongia muta</i>       | <i>Diploria strigosa</i>    |                              |
|                                | <i>Montastrea annularis</i> |                              |
| Hydrozoa (fire coral)          | <i>Montastrea cavernosa</i> |                              |
| <i>Millepora alcicornis</i>    | <i>Porites porites</i>      |                              |

## Environmental Observations

### Hydrographic.— Bank (3 m)

Coral bleaching occurred at Lee Stocking Island following a short period (18 days) of daily average water temperatures above  $30.5^{\circ}\text{C}$  on the bank in July-August 1990 followed by another temperature peak in mid-September (Figure 2). A maximum temperature of  $32.5$  was reached on 23 September 1990. The July monthly average for 1990 did not differ from surrounding years, but August and September were significantly warmer in 1990 (Table 2). The 5-yr record at the LSI dock distinctly shows a longer-term maxima in 1990 (Figure 3); this trend needs to be substantiated.

We examined two trends in water temperature in detail. One hypothesis was that a more rapid rate of warming in the spring might result in a pool of warm water upon which singular events in the summer could react. Spring water temperature increase is relatively linear from April through June, so we compared the rate of spring temperature rise for 1988-1992 (Table 3). In 1990 the rate of spring water temperature increase was not exceptional, in fact in most years the temperature rose more rapidly than in 1990.

Typically by the end of August water temperature starts its annual decline. Examination of plots of the yearly data suggest that the warm period was prolonged in 1990 (Figure 2). To test this hypothesis we examined the temperature trends from July through September. Our hypothesis of prolonged warming would be confirmed if the slope of the trend was significantly less than in other years. For bank waters 1990 was the only year with a significant positive slope for this period confirming the prolonged warming period in 1990 (Table 3).

Great Bahama Bank salinities are typically 38 ppt and can reach 43 ppt (Pitts and Smith, 1993), thus adding a synergistic stress factor to the local environment (Marcus and Thorhaug, 1981). During July and August 1990 pulses of warm (to  $31.2^{\circ}\text{C}$ ) saline (to 40 ppt) water exited through island passes onto the shelf. Bank patch reefs previously studied by Lang et al. (1988) are bathed in these water masses without apparent detrimental effects.




Figure 2. Five-year record of water temperature at the LSI Dock.




Figure 3. Annual water temperature record for 1989-1991 at the LSI Dock.

*Hydrographic.*— Fringing Reef (20 m)

Perry reef also exhibited two temperature peaks in July/August and at the end of September in 1990 with the second being higher. In addition the longer-term maxima in 1990 is evident. Warming by month paralleled that observed on the bank with September being significantly warmer than other years (Table 2). Increasing temperature through the summer also is evident at this location (Table 3). No salinity data is available from this area, though Perry Reef is upstream from the pass at the north end of the island. Given the prevailing currents, hypersaline waters would not be expected to reach this reef on a regular basis, though they are known to occur there.

*Hydrographic.*— Shelf-edge Reef (33 m)

The deepwater basin of Exuma Sound acts as a buffer to water temperature change, being warm in the winter and cooler in the summer than shallow bank waters. Instrument failure resulted in missing most of July and August 1990 for this location. Early July was no warmer than other years but by the last week in September temperature was elevated and comparable to the fringing reef (Table 2). Though our data are more fragmented at the shelf-edge reef, the longer-term trend is still evident and indicates that in 1990 water temperatures at 33 m were the highest of the 5-yr record; even at 98 m this warming was evident.

There was no conductivity recorder on the shelf in 1990, but data from a near-bottom recorder in 1992 showed that high salinity pulses can reach the bottom at this site, a mechanism for carrying warm water to the deep reefs.

Table 2. Comparison of average monthly water temperature for summer months by year. Means compared by ANOVA followed by Tukey's HSD test. Underlined years are not significantly different.

|                               | Tukey's HSD Test for Differences Among Years |             |             |             |             | Monthly Average Temperature (°C) |       |       |       |       |
|-------------------------------|----------------------------------------------|-------------|-------------|-------------|-------------|----------------------------------|-------|-------|-------|-------|
|                               | LOW<----->HIGH                               |             |             |             |             | 1988                             | 1989  | 1990  | 1991  | 1992  |
| July                          |                                              |             |             |             |             |                                  |       |       |       |       |
| Dock                          | <u>1992</u>                                  | <u>1989</u> | <u>1990</u> | <u>1988</u> | <u>1991</u> | 29.70                            | 29.52 | 29.61 | 29.74 | 29.16 |
| Perry Reef                    | <u>1989</u>                                  | <u>1990</u> | <u>1988</u> | <u>1992</u> |             | 28.23                            | 27.86 | 28.17 | ---   | 28.78 |
| Adderley<br>(2-21 Jul)        | <u>1990</u>                                  | <u>1988</u> | <u>1989</u> | <u>1991</u> | <u>1992</u> | 26.97                            | 27.02 | 26.78 | 27.39 | 28.15 |
| August                        |                                              |             |             |             |             |                                  |       |       |       |       |
| Dock                          | <u>1992</u>                                  | <u>1988</u> | <u>1991</u> | <u>1989</u> | <u>1990</u> | 29.34                            | 30.10 | 30.49 | 29.96 | 29.27 |
| Perry Reef<br>(21-31 Aug)     | <u>1988</u>                                  | <u>1989</u> | <u>1992</u> | <u>1990</u> | <u>1991</u> | 28.44                            | 28.68 | 29.53 | 29.53 | 28.80 |
| September                     |                                              |             |             |             |             |                                  |       |       |       |       |
| Dock                          | <u>1992</u>                                  | <u>1988</u> | <u>1991</u> | <u>1989</u> | <u>1990</u> | 29.56                            | 29.74 | 30.59 | 29.62 | 28.48 |
| Perry Reef                    | <u>1992</u>                                  | <u>1988</u> | <u>1989</u> | <u>1991</u> | <u>1990</u> | 28.57                            | 28.94 | 29.79 | 29.45 | 28.44 |
| Adderley<br>(28 Sep - 10 Oct) | <u>1992</u>                                  | <u>1989</u> | <u>1988</u> | <u>1991</u> | <u>1990</u> | 28.72                            | 28.71 | 29.65 | 28.99 | 28.65 |

Table 3. Comparison of water temperature trends for spring and summer at two locations. Slopes compared by ANCOVA followed by Tukey's HSD test. Underlined years are not significantly different.

Tukey's HSD Test for Differences Among Years  
Trend Slope  
LOW<----->HIGH

|                         |             |             |             |             |             |
|-------------------------|-------------|-------------|-------------|-------------|-------------|
| Spring (April-June)     |             |             |             |             |             |
| Dock                    | <u>1989</u> | <u>1990</u> | <u>1988</u> | <u>1991</u> | <u>1992</u> |
| Perry Reef              | <u>1991</u> | <u>1990</u> | <u>1989</u> | <u>1992</u> | <u>1988</u> |
| Summer (July-September) |             |             |             |             |             |
| Dock                    | <u>1992</u> | <u>1988</u> | <u>1991</u> | <u>1989</u> | <u>1990</u> |
| Perry Reef              | <u>1991</u> | <u>1992</u> | <u>1988</u> | <u>1989</u> | <u>1990</u> |

*Meteorological.*—

Wind data at Lee Stocking Island are available for 1990-1992; typically July-October is the calmest period of the year. There is no significant difference in number of days of low winds ( $\leq 3$  m/s) between 1990 and 1991, but 1992 was significantly more windy (Table 4). In addition 1992 had significantly fewer days with above average insolation (Table 4). There was no difference in the number of summer days with rainfall between 1990 and 1991, though 1990 had only about 60% of the total summer rainfall of 1991.

When examined in detail the prolonged warm water period from 24 July to 3 October 1990 was the result of three rises in temperature closely related to declines in wind speed (Figure 4). Over a 14-day period in late July water temperature increased from a daily average of 28.6° to 31.7°C, concurrent with a period of low winds, little insolation. The second small rise is related to a short period of reduced winds (28-31 Aug) and no increase in insolation (Figure 4). The final temperature rise is associated with a 12-day calm period (14-30 Sept) with an increase in insolation followed by rain and declining water and air temperature.




Figure 4. Daily average bank water temperature, wind speed, and insolation for the summer of 1990.

Table 4. Frequency of calm days ( $\leq 3$  m/s), days with rain, and calm days with above average insolation in the summer at Lee Stocking Island. Arrows indicate significant differences based on chi-square test.

| Number of days with calm winds |     |     |     |                | Number of Days with Rain |     |     |       | Total Summer Rainfall (mm) |
|--------------------------------|-----|-----|-----|----------------|--------------------------|-----|-----|-------|----------------------------|
| Year                           | JUL | AUG | SEP | Total          | JUL                      | AUG | SEP | Total |                            |
| 1990                           | 8   | 9   | 13  | 30             | 8                        | 12  | 8   | 28    | 63.85                      |
| 1991                           | 6   | 9   | 13  | 28             | 11                       | 10  | 9   | 30    | 106.83 $\uparrow$          |
| 1992                           | 0   | 4   | 5   | 9 $\downarrow$ | 5                        | 10  | 15  | 30    | 83.92                      |

Number of calm days with above average insolation

| Year | JUL | AUG | SEP | Total |
|------|-----|-----|-----|-------|
| 1990 | 6   | 6   | 6   | 18    |
| 1991 | 4   | 3   | 6   | 13    |
| 1992 | 0   | 1   | 2   | 3     |

## DISCUSSION

Corals thrive near their upper thermal tolerance level in summer prompting claims that global warming might cause more frequent thermal stress (Williams and Bunkley-Williams, 1990). There were no sea-surface-temperature (SST) anomalies in 1990 (Atwood et al., 1992, but see Strong and McCormick, 1990), though a trend in increasing SST through the 1980's for the Caribbean has been suggested (Goreau et al., 1992). Certainly no irrefutable evidence of SST anomalies in the Caribbean exists such as found in the 1982/83 ENSO event for the eastern Pacific (Glynn, 1990). Our whole range of thermographs (3-98 m) indicates that 1990 was warmer than the surrounding years. The 1990 bleaching event at Lee Stocking Island is correlated with elevated water temperatures, possibly a result of favorable weather conditions on top of a longer-term maxima in temperature. Bleaching might have been negligible but for a late September calm period that prolonged the period of elevated water temperatures. Rainfall appears to have a mitigating influence on the effect of solar heating on bank waters as there was little rise in water temperature when a calm period was associated with substantial rainfall.

Rather than deeper waters staying cooler in the summer, shelf-edge water temperature rose close to levels at shallow-water fringing reefs. Deeper-water reefs must be adapted to lower temperatures based on the lower maximum temperature causing bleaching as has been proposed for Bermuda corals (Cook et al., 1990).

Bank waters are typically hypersaline, ranging from 38-44 ppt and corals must be adapted to these conditions as well developed patch reefs exist in the Bahamas (Newell et al., 1959). The intrusion of these warm hypersaline conditions on shelf habitats is not as well documented. Basin waters have typical oceanic salinities (36-37 ppt) (US Navy, 1967; Schroeder, 1977; this study), thus shelf-edge reefs should not be acclimated to hypersaline conditions. Periodic density cascading of warm hypersaline water off the bank may be an additional stress on shelf reefs especially around passes. The role of salinity stress cannot be accurately evaluated without further information on the frequency, duration, and areal extent of density cascading.

## CONCLUSIONS

Elevated water temperature around Lee Stocking Island was closely related to a significant coral bleaching event in 1990. Bleaching was not the result of a single upper level temperature threshold, as bleaching occurred over a range of depths and temperatures suggesting a large pool of water with elevated temperature. Elevated temperatures in 1990 were related to local climatological phenomena, including calm wind conditions, high insolation, and low rainfall that occurred on top of a longer-term maxima in water

temperature. Rainfall, not previously considered an important factor, can have a substantial influence on water temperature. Exposure to elevated water temperature (>31°C on the bank) in 1990 appeared to take more than a couple of days, but less than three weeks, to initiate bleaching. Prolonged elevated temperatures may be the primary contributing factor to the magnitude of the bleaching event in 1990.

#### ACKNOWLEDGEMENTS

J. Lang and R. Dill provided additional information on bleaching organisms and N. Smith provided salinity data. K. Mueller maintained the water temperature database and provided quality control. Review by G. Wenz helped improve the paper. The NOAA National Undersea Research Program provided support for this project.

#### LITERATURE CITED

- Atwood, D.K., J.C. Hendee and A. Mendez. 1992. An assessment of global warming stress on Caribbean coral reef ecosystems. *Bull. Mar. Sci.* 51: 118-130.
- Cook, C.B., A. Logan, J. Ward, B. Luckhurst and C.J. Berg. 1990. Elevated temperatures on a high latitude coral reef: the 1988 Bermuda event. *Coral Reefs* 9: 45-49.
- Glynn, P.W. 1990. Coral mortality and disturbances to coral reefs in the Tropical Eastern Pacific. Pages 55-126 *In* P.W. Glynn, ed. *Global Ecological Consequences of the 1982-83 El Niño-Southern Oscillation*, Elsevier Oceanogr. Ser.
- Goreau, T.J., R.L. Hayes, J.W. Clark, D.J. Basta and C.N. Robertson. 1992. Elevated satellite sea surface temperatures correlated with Caribbean coral bleaching. *In* R.A. Geyer, Ed. *Geophysical/Geochemical Aspects of Global Warming*. CRC Press
- Jaap, W.C. 1979. Observations on zooxanthellae expulsion at Middle Sambo Reef, Florida Keys. *Bull. Mar. Sci.* 29: 414-422.
- Jaap, W.C. 1985. An epidemic zooxanthellae expulsion during 1983 in the lower Florida Keys coral reefs: hyperthermic etiology. *Proc. 5th Inter. Coral Reef Congr. Tahiti* 6: 143-148.
- Jokiel, P.L. and S.L. Coles. 1990. Response of Hawaiian and other Indo-Pacific reef corals to elevated temperature. *Coral Reefs* 8: 155-162.
- Lang, J.C., R.I. Wicklund and R.F. Dill. 1988. Depth- and habitat-related bleaching of zooxanthellate reef organisms near Lee Stocking Island, Exuma Cays, Bahamas. *Proc. 6th Inter. Coral Reef Symp. Townsville, Aust.* 3: 269-274.
- Marcus, J. and A. Thorhaug, 1981. Pacific versus Atlantic responses of the subtropical hermatypic coral *Porites* spp. to temperature and salinity effects. *Proc. 4th Inter. Coral Reef Symp. Manila* 2: 15-20.
- Newell, N.D., J. Imbrie, E.G. Purdy and D.L. Thurber. 1959. Organism communities and bottom facies, Great Bahama Bank. *Bull. Amer. Mus. Nat. Hist.* 117(4): 177-228.
- Pitts, P.A. and N.P. Smith. 1993. Annotated summary of temperature and salinity data from the vicinity of Lee Stocking Island, Exuma Cays, Bahamas. CMRC Tech. Rept. No. 93-3, Vero Beach, FL
- Schroeder, W.W. 1977. Current and hydrographic characterization of the south central insular shelf of Grand Bahama Island. *Proc. 3rd Inter. Coral Reef Symp., Miami* 2: 517-523.
- Strong, A.E. and R.C. McCormick. 1991. Coral bleaching and sea surface temperatures. *Proc. Oceanogr. Soc. Conf. St. Petersburg, FL.*
- U.S. Navy. 1967. Environmental Atlas of the Tongue of the Ocean Bahamas. U.S. Naval Oceanographic Office, Spec. Publ. SP-94, 74 p.
- Wicklund, R.I., G.D. Dennis and K.W. Mueller. 1993. Summary of data from the Water Temperature Monitoring Network at Lee Stocking Island, Bahamas, 1988-1991. CMRC Tech Rept. No. 93-1, Vero Beach, FL
- Williams, E.H. and L. Bunkley-Williams. 1990. The world-wide coral reef bleaching cycle and related sources of coral mortality. *Atoll Res. Bull.* No. 335, 71 p.