

EARLY STAGE EXPERIMENT

*Science Description*

---

**Experiment/Module:** Convective Burst Structure and Evolution Module (CBM)

**Investigator(s):** Rob Rogers (PI), Jon Zawislak, Trey Alvey, Michael Bell (CSU), Anthony Didlake (PSU), Josh Wadler (UM/RSMAS)

**Requirements:** TD, TS, Category 1

**Plain Language Description:** This module samples the vertical motion and reflectivity structure of strong thunderstorm complexes at a high frequency, e.g., every 15-20 minutes, over a 1-2 h period to observe how the structure of these systems change over time and as they move around the TC center, along with observing how those changes affect the structure and intensity of TCs.

**Early Stage Science Objective(s) Addressed:**

1. Obtain a quantitative description of the kinematic and thermodynamic structure and evolution of intense convective systems (convective bursts) and the nearby environment to examine their role in TC intensity change [*IFEX Goals 1, 3*].

**Motivation:** The objectives are to obtain a quantitative description of the kinematic and thermodynamic structure and evolution of intense convective systems (convective bursts) and the nearby environment to examine their role in TC intensity change.

**Background:** It has long been known that deep convection is an integral component of TC structure. What has received greater attention in recent years is the potential role that deep convection, termed here “convective bursts”, or CBs, representing the peak updrafts and highest echo tops, plays in TC structure and intensity evolution. Various hypotheses attribute their contribution to TC intensification by vortex gradient adjustment to the imposed diabatic heating in the high-inertial stability region inside the radius of maximum wind (RMW) (e.g., Shapiro and Willoughby 1982, Schubert and Hack 1982, Hack and Schubert 1986, Nolan and Grasso 2003, Nolan et al. 2007, Vigh and Schubert 2009, Pendergrass and Willoughby 2009, Rogers et al. 2013, 2015, 2016), convergence of angular momentum surfaces in the lower troposphere and boundary layer (Smith and Montgomery 2016), upper-level subsidence warming around the CB periphery (e.g., Heymsfield et al. 2001, Guimond et al. 2010, Rogers 2010, Zhang and Chen 2012, Chen and Zhang 2013, Chen and Gopal 2015), stretching and axisymmetrization in vortical hot towers (Hendricks et al. 2004, Montgomery et al. 2006, Reasor et al. 2009), and vortex alignment/downshear reformation (Reasor et al. 2009, Molinari and Vollaro 2010, Nguyen and Molinari 2012, Reasor and Eastin 2012, Stevenson et al. 2014, Rogers et al. 2015, 2020, Nguyen and Molinari 2015). While these studies have emphasized the role of deep convection in TC intensification, other studies have focused on the role of shallow to moderate convection, and even stratiform precipitation, in initiating TC intensification (Kieper and Jiang 2012, Zagrodnik and Jiang 2014, Tao and Jiang 2015, Tao et al. 2017, Nguyen et al. 2017, Rogers et al. 2020). Common to these and other (e.g., Miyamoto and Takemi 2015) studies, though, is that TC intensification is favored when precipitation, including CBs, are preferentially located inside the RMW with a maximum azimuthal distribution.

EARLY STAGE EXPERIMENT

*Science Description*

---

Vertical shear is one factor that has been shown to be important in organizing precipitation, including CBs, azimuthally around the TC vortex. This has generally been attributed to the fact that vertical shear tilts the vortex, leading to preferred regions of vortex-scale low-level convergence and upward motion downshear and low-level divergence and subsidence upshear (Jones 1995, Bender 1997, Frank and Ritchie 2001, Black et al. 2002, Corbosiero and Molinari 2003, Rogers et al. 2003, Braun et al. 2006, Wu et al. 2006, Reasor et al. 2009, Reasor and Eastin 2012, Reasor et al. 2013, Dolling and Barnes 2014, DeHart et al. 2014). Recent composite studies of vortices in shear using airborne Doppler radar have shown that the shear-induced circulations are maximized downshear right (DSR) (low-level convergence/upward motion) and upshear left (USL) (low-level divergence/downward motion) (Reasor et al. 2013, DeHart et al. 2014). A similar composite methodology was performed in a CB-relative coordinate system (Wadler et al. 2018). This study found that the peak updraft magnitude and altitude for CBs was minimized DSR, consistent with the notion that this is the quadrant where CBs are initiated. Peak updraft magnitude and altitude increase in the DSL quadrant, as the CBs mature, and they reach their highest and strongest values USL. A similar shear-relative azimuthal relationship was found for echo top height. Significantly, when stratifying TCs by intensity change, it was found that the most significant differences in CB structure between intensifying and non-intensifying TCs were located in the USL quadrant. Intensifying TCs have CBs with stronger peak updrafts, at a higher altitude, with higher echo tops in the USL quadrant than non-intensifying TCs. This relationship suggests that the structure and evolution of CBs, which are to some extent a function of the local environment from which they initiate downshear and mature upshear -- including convective available potential energy, midlevel humidity, and subsidence upshear (Zawislak et al. 2016, Rogers et al. 2016, Nguyen et al. 2017, 2019) -- is an important factor to consider in assessing the potential for a TC to intensify.

It should be noted that the above descriptions presume that CBs do translate downwind, i.e., upshear. However, in some situations, mostly revealed from modeling studies (Munsell et al. 2017, Chen et al. 2017), CBs can remain “trapped” on the downshear side. In fact, cases where the CBs remain downshear were more likely to be associated with non-intensifying periods of TC evolution. This is consistent with the notion of greater azimuthal symmetry of diabatic heating being associated with TC intensification. CBs propagating into the upshear quadrants may also be related to a greater likelihood of vortex alignment, as revealed in the observational analysis of Hurricane Earl (2010; Rogers et al. 2015) and a WRF-ARW ensemble forecast of Edouard (2014; Munsell et al. 2017).

As a CB matures it transitions to predominantly stratiform precipitation characterized by different microphysical processes than during the active phase of convection (Houze 1997). Ice processes such as vapor deposition and aggregation take on a more dominant role as vertical motions become weaker, and evaporation below the melting level also becomes important. Dynamically, the shift from active convection with strong low-level convergence and vortex stretching to stratiform precipitation with mid-level convergence and low-level divergence has an impact on the vortex intensification (Bell and Montgomery 2019), and may also play an important role in secondary eyewall formation (Didlake et al. 2018). In an early stage TC, the transition from a CB to stratiform may occur in roughly the same location, but as the rotational wind speeds intensify the transition

**EARLY STAGE EXPERIMENT**

*Science Description*

---

can occur cyclonically downwind (Didlake and Houze 2013, Foerster et al. 2014). The microphysical processes involved in this transition must be parameterized in high-resolution models but not well-understood in the TC environment (Feng and Bell 2018).

The results described above are valid for composites of many different CBs from many different TCs. They therefore lack the temporal continuity needed to measure the structure of specific individual (or groups of) CBs, and how they evolve in a shear-relative sense. The purpose of this module is to repeatedly sample individual (or groups of) CBs to provide this temporal continuity.

**Goal(s):** Obtain a quantitative description of the kinematic and thermodynamic structure and evolution of intense convective systems (convective bursts) and the nearby environment to examine their role in TC intensity change.

**Hypotheses:** The following hypotheses will guide the sampling strategies for CBs:

1. CBs are preferentially initiated in the DSR quadrant; as such, the updraft maxima are likely to be weaker and at a lower altitude in this quadrant;
2. Traveling downwind into the DSL quadrant, peak updrafts will strengthen and be located at a higher altitude;
3. The strength of the CB in the USL quadrant (as measured by strength and height of peak updraft and echo top height relative to the DSL quadrant) will vary depending on the local, vortex-scale environment of the convection. This environment includes midlevel humidity, strength of subsidence upshear, and sea surface temperature (and CAPE) on the downshear side of the TC;
4. If the CB strength, as measured by peak updraft magnitude and altitude and echo top height, is higher USL than DSL, then the CB will persist on the upshear side with a mixture of convective and stratiform precipitation, and the TC will be more likely to intensify.
5. If CBs are primarily maximized DSL, the upshear will be more dominated by stratiform precipitation with downdrafts that are potentially detrimental to intensification.

**Objectives:**

1. The objectives of this module are to repeatedly sample the kinematic and reflectivity structure and evolution of individual convective bursts and their associated precipitation structures using airborne Doppler radar
2. Sampling will follow CBs as they translate azimuthally downwind in a shear-relative framework
3. Spiral ascents in stratiform regions of CB systems at radii outside the RMW will provide microphysical measurements from in situ probes, include how the microphysics vary in a shear-relative framework
4. Optimal additional measurements include:
  - a. Deep-layer measurements of temperature and humidity in the local environment of the CBs from high-altitude aircraft will provide the thermodynamic context within the mid-and lower troposphere

**EARLY STAGE EXPERIMENT**  
*Science Description*

---

- b. Measurements of sea-surface temperature and subsurface temperature profiles from ocean probes and/or IR dropsondes will provide context on the surface boundary

**Aircraft Pattern/Module Descriptions (see *Flight Pattern* document for more detailed information):** This is a stand-alone module that takes 1-2 h to complete. Execution is dependent on system attributes, aircraft fuel and weight restrictions, and proximity to operations base. It can be flown separately within a mission designed to study local areas of convection or at the end of one of the survey patterns. Once a local area of intense convection is identified, the P-3 will transit at altitude (10-12 kft) to the nearest point just outside of the convective cores and sample the convective area. The sampling pattern will be a series of inbound/outbound radial penetrations or bowtie patterns (when sampling a CB near the radius of maximum wind of a tropical storm or hurricane). If the CB is at or near the RMW, repeated sampling can allow for a following of the burst around the storm. This is especially useful to sample the structural evolution of the burst as it moves around the storm. If the CB remains confined to the downshear side of the TC rather than translating upshear, the pattern should still be flown. If the CB has transitioned to stratiform convection and is located at a larger radius, a spiral ascent and descent can be made to obtain direct hydrometeor measurements with the P-3 cloud and precipitation probes. This module can also be flown during the mature stage, in conjunction with the rainband module.

**Links to Other Early Stage Experiments/Modules:** CBM can be flown in conjunction with the following *Early Stage* experiments: AIPEX, TDR Experiment, Synoptic Flow Experiment, and NESDIS JPSS Satellite Validation Experiment.

**Analysis Strategy:** Radar analyses will be performed for each radial pass through the CB, preferentially with a temporal spacing of 30 minutes or less. These analyses will provide high-frequency observations of the structure of the CB, as measured by the peak updraft magnitude and altitude and echo top heights. Additionally, the full spectrum of vertical velocity associated with each radar analysis will be evaluated using contoured frequency by altitude diagrams (CFADs; Yuter and Houze 1995) to obtain a more complete picture of the updraft and downdraft structure and evolution of the CB. Ideally a CB will be flown beginning with its initiation (likely to be downshear) and then followed around the storm as it travels through the downwind quadrants and into the upshear quadrants (or continuously sampled on the downshear side if it remains confined there). If the CB transitions to stratiform then cloud and precipitation probe data will document the hydrometeor characteristics in the stratiform transition of the CB, along with in situ measurements of vertical velocity and thermodynamic information. Dropsondes released at the starting and ending points of each radial leg will document the thermodynamic structure of the boundary layer radially bracketing the CB. A higher altitude dropsonde release at the apex of any spiral ascents will provide a deeper profile of thermodynamic structure.

Optimally, the G-IV will be flying in the storm to provide deep-layer humidity profiles around the storm in addition to the P-3 dropsondes. If spiral ascents are performed at larger radii in range of the G-IV radar then the in situ and radar data can be synthesized to provide a more complete picture of the stratiform cloud dynamics. If the G-IV is not available, the module could still be flown to examine the evolution using the Doppler radar and boundary layer thermodynamics from the P-3

EARLY STAGE EXPERIMENT  
*Science Description*

---

dropsondes. Additionally, when multiple P-3 aircraft are available and sampling simultaneously, one aircraft can perform the radial penetrations as outlined above, while the other aircraft circumnavigates the CB outside the precipitation shield at the maximum allowable altitude (preferably  $\geq 18$  kft). Dropsondes should be released from the higher altitude aircraft at locations upwind and downwind from the CB.

In addition to the observational analysis described above, the high-resolution data collected in this module is planned to be embedded within the typical Hurricane Ensemble Data Assimilation System (HEDAS; e.g., Aksoy et al. 2013) framework to carry out storm-scale data assimilation that focuses specifically on the high-resolution analysis of the identified intense convective region. With current technology, a smaller domain with 1-km grid spacing will be nested within the HEDAS 3-km analysis domain, where the data will be assimilated for the duration of its collection (1-2 hours, at 5-10 min intervals). This is a typical setup that has been traditionally used in continental storm-scale radar data assimilation applications and has been shown to be effective to obtain realistic storm structures in analyses and short-range forecasts. With such high-resolution analyses, we hope to be able to obtain fully three-dimensional model representations of the observed convective regions for more detailed investigation, as well as investigate their short-range predictability. In an observing system experiment (OSE) mode, various assimilation experiments can also be devised to investigate hypothetical scenarios for how an observed convective region could interact with the surrounding vortex and impact its evolution. Dropsonde, superobbed Doppler, flight-level, and SFMR data will be transmitted over the GTS and assimilated in real time; full Doppler analyses and lower fuselage imagery will be available post-flight.

**References:**

- Aksoy, A., S. D. Aberson, T. Vukicevic, K. J. Sellwood, S. Lorsolo, and X. Zhang (2013), Assimilation of high-resolution tropical cyclone observations with an ensemble Kalman filter using NOAA/AOML/HRD's HEDAS: Evaluation of the 2008–11 vortex-scale analyses, *Mon. Wea. Rev.*, **141**, 1842–1865, doi: <http://dx.doi.org/10.1175/MWR-D-12-00194.1>.
- Bell, M. M., and M. T. Montgomery, 2019: Mesoscale Processes During the Genesis of Hurricane Karl (2010). *J. Atmos. Sci.*, *in press*.
- Bender, M. A., 1997: The effect of relative flow on the asymmetric structure in the interior of hurricanes. *J. Atmos. Sci.*, **54**, 703–724, doi:10.1175/1520-0469(1997)054<0703:TEORFO>2.0.CO;2.
- Black, M. L., J. F. Gamache, F. D. Marks, C. E. Samsury, and H. E. Willoughby, 2002: Eastern Pacific Hurricanes Jimena of 1991 and Olivia of 1994: The effect of vertical shear on structure and intensity. *Mon. Wea. Rev.*, **130**, 2291–2312, doi:10.1175/1520-0493(2002)130<2291:EPHJOA>2.0.CO;2.
- Braun, S. A., M. T. Montgomery, and Z. Pu, 2006: High-resolution simulation of Hurricane Bonnie (1998). Part I: The organization of eyewall vertical motion. *J. Atmos. Sci.*, **63**, 19–42, doi:10.1175/JAS3598.1.
- Chen, H., and S. G. Gopalakrishnan, 2015: A study on the asymmetric rapid intensification of Hurricane Earl (2010) using the HWRF system. *J. Atmos. Sci.*, **72**, 531–550, doi:10.1175/JAS-D-14-0097.1.
- Chen, H., and D. L. Zhang, 2013: On the rapid intensification of Hurricane Wilma (2005). Part II: Convective bursts and the upper-level warm core. *J. Atmos. Sci.*, **70**, 146–162, doi:10.1175/JAS-D-12-062.1.
- Chen, H., S. Gopalakrishnan, J.A. Zhang, R.F. Rogers, Z. Zhang, and V. Tallapragada, 2017: Use of HWRF ensembles for providing improved understanding of hurricane RI problem: Case study of Hurricane Edouard (2014). *J. Atmos. Sci.*, *in review*.

## EARLY STAGE EXPERIMENT

*Science Description*

- 
- Corbosiero, K. L., and J. Molinari, 2003: The relationship between storm motion, vertical wind shear, and convective asymmetries in tropical cyclones. *J. Atmos. Sci.*, **60**, 366–376, doi:10.1175/1520-0469(2003)060<0366:TRBSMV>2.0.CO;2.
- DeHart, J. C., R. A. Houze Jr., and R. F. Rogers, 2014: Quadrant distribution of tropical cyclone inner-core kinematics in relation to environmental shear. *J. Atmos. Sci.*, **71**, 2713–2732, doi:10.1175/JAS-D-13-0298.1.
- Didlake, A. C., and R. A. Houze, 2013: Dynamics of the stratiform sector of a tropical cyclone rainband. *J. Atmos. Sci.*, **70**, 1891–1911.
- Didlake, A.C., P.D. Reasor, R.F. Rogers, and W. Lee, 2018: Dynamics of the Transition from Spiral Rainbands to a Secondary Eyewall in Hurricane Earl (2010). *J. Atmos. Sci.*, **75**, 2909–2929, <https://doi.org/10.1175/JAS-D-17-0348.1>
- Dolling, K., and G. M. Barnes, 2014: The evolution of Hurricane Humberto (2001). *J. Atmos. Sci.*, **71**, 1276–1291, doi:10.1175/JAS-D-13-0164.1.
- Feng, Y.-C., and Bell, M. M, 2018: “Microphysical characteristics of an asymmetric eyewall in major Hurricane Harvey (2017)”. *Geophys. Res. Lett.*, doi: 10.1029/2018gl080770.
- Frank, W. M., and E. A. Ritchie, 2001: Effects of vertical wind shear on the intensity and structure of numerically simulated hurricanes. *Mon. Wea. Rev.*, **129**, 2249–2269, doi:10.1175/1520-0493(2001)129<2249:EOVWSO>2.0.CO;2.
- Foerster, A. M., M. M. Bell, P. A. Harr, and S. C. Jones, 2014: Observations of the eyewall structure of Typhoon Sinlaku (2008) during the transformation stage of extratropical transition. *Mon. Wea. Rev.*, **142**, 3372–3392.
- Guimond, S. R., G. M. Heymsfield, and F. J. Turk, 2010: Multiscale observations of Hurricane Dennis (2005): The effects of hot towers on rapid intensification. *J. Atmos. Sci.*, **67**, 633–654, doi:10.1175/2009JAS3119.1.
- Hack, J. J., and W. H. Schubert, 1986: Nonlinear response of atmospheric vortices to heating by organized cumulus convection. *J. Atmos. Sci.*, **43**, 1559–1573, doi:10.1175/1520-0469(1986)043<1559:NROAVT>2.0.CO;2.
- Hendricks, E. A., M. T. Montgomery, and C. A. Davis, 2004: The role of “vortical” hot towers in the formation of Tropical Cyclone Diana (1984). *J. Atmos. Sci.*, **61**, 1209–1232, doi:10.1175/1520-0469(2004)061<1209:TROVHT>2.0.CO;2.
- Heymsfield, G. M., J. B. Halverson, J. Simpson, L. Tian, and T. P. Bui, 2001: ER-2 Doppler radar investigations of the eyewall of Hurricane Bonnie during the Convection and Moisture Experiment-3. *J. Appl. Meteor.*, **40**, 1310–1330, doi:10.1175/1520-0450(2001)040<1310:EDRIOT>2.0.CO;2.
- Houze, R. A., 1997: Stratiform precipitation in regions of convection: A meteorological paradox? *Bull. Amer. Meteor. Soc.*, **78**, 2179–2196.
- Jones, S. C., 1995: The evolution of vortices in vertical shear. I: Initially barotropic vortices. *Quart. J. Roy. Meteor. Soc.*, **121**, 821–851, doi:10.1002/qj.49712152406.
- Kieper, M. E., and H. Jiang, 2012: Predicting tropical cyclone rapid intensification using the 37 GHz ring pattern identified from passive microwave measurements. *Geophys. Res. Lett.*, **39**, L13804, doi:10.1029/2012GL052115.
- Miyamoto, Y., and T. Takemi, 2015: A triggering mechanism for rapid intensification of tropical cyclones. *J. Atmos. Sci.*, **72**, 2666–2681, doi:10.1175/JAS-D-14-0193.1.
- Molinari, J., and D. Vollaro, 2010: Rapid intensification of a sheared tropical storm. *Mon. Wea. Rev.*, **138**, 3869–3885, doi:10.1175/2010MWR3378.1.
- Montgomery, M. T., M. Nicholls, T. Cram, and A. Saunders, 2006: A “vortical” hot tower route to tropical cyclogenesis. *J. Atmos. Sci.*, **63**, 355–386, doi:10.1175/JAS3604.1.

## EARLY STAGE EXPERIMENT

*Science Description*

- 
- Munsell, E.B., F. Zhang, J.A. Sippel, S.A. Braun, and Y. Weng, 2017: Dynamics and Predictability of the Intensification of Hurricane Edouard (2014). *J. Atmos. Sci.*, **74**, 573–595, doi: 10.1175/JAS-D-16-0018.1.
- Nguyen, L. T., and J. Molinari, 2012: Rapid intensification of a sheared, fast-moving hurricane over the Gulf Stream. *Mon. Wea. Rev.*, **140**, 3361–3378, doi:10.1175/MWR-D-11-00293.1.
- Nguyen, L. T., and J. Molinari, 2015: Simulation of the downshear reformation of a tropical cyclone. *J. Atmos. Sci.*, **72**, 4529–4551, doi:10.1175/JAS-D-15-0036.1.
- Nguyen, L.T., R.F. Rogers, and P.D. Reasor, 2017: Thermodynamic and kinematic influences on precipitation symmetry in sheared tropical cyclones: Bertha and Cristobal (2014). *Mon. Wea. Rev.*, in review.
- Nolan, D. S., and L. D. Grasso, 2003: Nonhydrostatic, three-dimensional perturbations to balanced, hurricane-like vortices. Part II: Symmetric response and nonlinear simulations. *J. Atmos. Sci.*, **60**, 2717–2745, doi:10.1175/1520-0469(2003)060<2717:NTPTBH>2.0.CO;2.
- Nolan, D. S., Y. Moon, and D. P. Stern, 2007: Tropical cyclone intensification from asymmetric convection: Energetics and efficiency. *J. Atmos. Sci.*, **64**, 3377–3405, doi:10.1175/JAS3988.1.
- Pendergrass, A. G., and H. E. Willoughby, 2009: Diabatically induced secondary flows in tropical cyclones. Part I: Quasi-steady forcing. *Mon. Wea. Rev.*, **137**, 805–821, doi:10.1175/2008MWR2657.1.
- Reasor, P. D., and M. D. Eastin, 2012: Rapidly intensifying Hurricane Guillermo (1997). Part II: Resilience in shear. *Mon. Wea. Rev.*, **140**, 425–444, doi:10.1175/MWR-D-11-00080.1.
- Reasor, P. D., M. D. Eastin, and J. F. Gamache, 2009: Rapidly intensifying Hurricane Guillermo (1997). Part I: Low-wavenumber structure and evolution. *Mon. Wea. Rev.*, **137**, 603–631, doi:10.1175/2008MWR2487.1.
- Reasor, P. D., R. F. Rogers, and S. Lorsolo, 2013: Environmental flow impacts on tropical cyclone structure diagnosed from airborne Doppler radar composites. *Mon. Wea. Rev.*, **141**, 2949–2969, doi:10.1175/MWR-D-12-00334.1.
- Rogers, R.F., 2010: Convective-scale structure and evolution during a high-resolution simulation of tropical cyclone rapid intensification. *J. Atmos. Sci.*, **67**, 44–70, doi:10.1175/2009JAS3122.1.
- Rogers, R., S. S. Chen, J. E. Tenerelli, and H. E. Willoughby, 2003: A numerical study of the impact of vertical shear on the distribution of rainfall in Hurricane Bonnie (1998). *Mon. Wea. Rev.*, **131**, 1577–1599, doi:10.1175//2546.1.
- Rogers, R., P. D. Reasor, and S. Lorsolo, 2013: Airborne Doppler observations of the inner-core structural differences between intensifying and steady-state tropical cyclones. *Mon. Wea. Rev.*, **141**, 2970–2991, doi:10.1175/MWR-D-12-00357.1.
- Rogers, R., P. D. Reasor, and J. A. Zhang, 2015: Multiscale structure and evolution of Hurricane Earl (2010) during rapid intensification. *Mon. Wea. Rev.*, **143**, 536–562, doi:10.1175/MWR-D-14-00175.1.
- Rogers, R.F., J.A. Zhang, J. Zawislak, H. Jiang, G.R. Alvey III, E.J. Zipser, and S.N. Stevenson, 2016: Observations of the structure and evolution of Hurricane Edouard (2014) during intensity change. Part II: Kinematic structure and the distribution of deep convection. *Mon. Wea. Rev.*, **144**, 3355–3376.
- Schubert, W. H., and J. J. Hack, 1982: Inertial stability and tropical cyclone development. *J. Atmos. Sci.*, **39**, 1687–1697, doi:10.1175/1520-0469(1982)039<1687:ISATCD>2.0.CO;2.
- Shapiro, L. J., and H. E. Willoughby, 1982: The response of balanced hurricanes to local sources of heat and momentum. *J. Atmos. Sci.*, **39**, 378–394, doi:10.1175/1520-0469(1982)039<0378:TROBHT>2.0.CO;2.
- Smith, R. K., and M. T. Montgomery, 2016: The efficiency of diabatic heating and tropical cyclone intensification. *Quart. J. Roy. Meteor. Soc.*, **142**, 2081–2086, doi:10.1002/qj.2804.
- Stevenson, S. N., K. L. Corbosiero, and J. Molinari, 2014: The convective evolution and rapid intensification of Hurricane Earl (2010). *Mon. Wea. Rev.*, **142**, 4364–4380, doi:10.1175/MWR-D-14-00078.1.

EARLY STAGE EXPERIMENT  
*Science Description*

---

- Tao, C., and H. Jiang, 2015: Distributions of shallow to very deep precipitation–convection in rapidly intensifying tropical cyclones. *J. Climate*, **28**, 8791–8824, doi:10.1175/JCLI-D-14-00448.1.
- Tao, C., H. Jiang, and J. Zawislak, 2017: The Relative Importance of Stratiform and Convective Rainfall in Rapidly Intensifying Tropical Cyclones. *Mon. Wea. Rev.*, **145**, 795–809, doi: 10.1175/MWR-D-16-0316.1.
- Vigh, J. L., and W. H. Schubert, 2009: Rapid development of the tropical cyclone warm core. *J. Atmos. Sci.*, **66**, 3335–3350, doi:10.1175/2009JAS3092.1.
- Wadler, J., R.F. Rogers, and P.D. Reasor, 2017: Radial and azimuthal variations in convective burst structure in tropical cyclones from airborne Doppler observations. *Mon. Wea. Rev.*, manuscript in review.
- Wu, L., S. A. Braun, J. Halverson, and G. Heymsfield, 2006: A numerical study of Hurricane Erin (2001). Part I: Model verification and storm evolution. *J. Atmos. Sci.*, **63**, 65–86, doi:10.1175/JAS3597.1.
- Zagrodnik, J. P., and H. Jiang, 2014: Rainfall, convection, and latent heating distributions in rapidly intensifying tropical cyclones. *J. Atmos. Sci.*, **71**, 2789–2809, doi:10.1175/JAS-D-13-0314.1.
- Zawislak, J., H. Jiang, G. R. Alvey III, E. J. Zipser, R. F. Rogers, J. A. Zhang, and S. N. Stevenson, 2016: Observations of the structure and evolution of Hurricane Edouard (2014) during intensity change. Part I: Relationship between the thermodynamic structure and precipitation. *Mon. Wea. Rev.*, **144**, 3333–3354, doi:10.1175/MWR-D-16-0018.1.
- Zhang, D.-L., and H. Chen, 2012: Importance of the upper level warm core in the rapid intensification of a tropical cyclone. *Geophys. Res. Lett.*, **39**, L02806, doi:10.1029/2012GL052355.
- Yuter, S. E., and R. A. Houze, 1995: Three-dimensional kinematic and microphysical evolution of Florida cumulonimbus. Part II: Frequency distributions of vertical velocity, reflectivity, and differential reflectivity. *Mon. Wea. Rev.*, **123**, 1941–1963, doi:10.1175/1520-0493(1995)123<1941:TDKAME>2.0.CO;2.