

GENESIS STAGE EXPERIMENT
Science Description

Experiment/Module: Favorable Air Mass (FAM)

Investigator(s): Ghassan Alaka, Jason Dunion, Rob Rogers, Sharan Majumdar (Univ. of Miami/RSMAS), Alexis Wilson (Univ. of Miami/RSMAS), Quinton Lawton (Univ. of Miami/RSMAS), Alan Brammer (CSU/CIRA), Chris Thorncroft (SUNY Albany)

Requirements: Pre-genesis disturbances (pre-tropical cyclones; pre-TCs), including NHC-designated “Invests”

Plain Language Description: Although the ingredients for tropical cyclone formation have been well-documented for decades, it is still difficult to predict which disturbances will develop and which ones will not. A big factor in this uncertainty is the favorability of the air mass ahead of, and interacting with, the disturbance. This experiment proposes to collect observations of mid-level humidity and winds to assess the favorability of the disturbance’s environment for tropical cyclogenesis. These aircraft observations may also provide helpful guidance for the expanded use of satellite observations in the absence of aircraft observations.

Genesis Stage Science Objective(s) Addressed:

The overarching objective is to investigate the physical processes that determine if a pre-genesis disturbance will mature into a tropical cyclone (TC), including the organization of convection and the development of a closed low-level circulation.

- 1) To investigate the favorability in both dynamics (e.g., vertical wind shear) and thermodynamics (e.g., moisture, stability) for tropical cyclogenesis in the environment near a pre-TC, especially the downstream environment [*APHEX Goal 3*].
- 2) Test new (or improved) technologies with the potential to fill gaps, both spatially and temporally, in the existing suite of airborne measurements in tropical disturbances that are in the pre-genesis or genesis stage. These measurements include improved three-dimensional representation of the tropical disturbance/TC wind field, more spatially dense thermodynamic sampling of the boundary layer, and more accurate measurements of ocean surface winds [*APHEX Goal 2*]

Motivation: The environment near a pre-TC is critical to the favorability for tropical cyclogenesis to occur. The probability of cyclogenesis for a given pre-TC, such as an African easterly wave, is dependent upon thermodynamics (e.g., moisture, relative humidity) and dynamics (e.g., vertical wind shear) in the adjacent air mass(es). Increased observations of lower- and mid-tropospheric humidity in the near-disturbance environment would shed light upon critical moisture thresholds important (or necessary) for tropical cyclogenesis and would help correct moisture biases in numerical weather prediction models. The downstream environment is most important for cyclogenesis predictions because that is the environment that a pre-TC moves into.

GENESIS STAGE EXPERIMENT

Science Description

Background: As early as the 1930s, westward propagating disturbances in the lower troposphere were identified as seed circulations for most TCs in the North Atlantic Ocean (Dunn 1940). The origins of these pre-genesis disturbances, or pre-tropical cyclones (pre-TCs), were traced back to North Africa and are now known as African easterly waves (AEWs; Riehl 1945). About 70% of all TCs and, more impressively, 85% of major hurricanes in the North Atlantic Ocean have been found to initiate from AEWs (Landsea 1993). On average, sixty AEWs exit the West African coast each year. However, determining which of these AEWs will develop into TCs has proven to be a forecasting challenge. For example, over 50% of TC genesis events in the Atlantic main development region predicted by the Global Forecast System (GFS) from 2004–2011 were false alarms (Halperin et al. 2013).

Recent research has shed some light on the relationship between AEWs and TC genesis in the North Atlantic Ocean. The AEW-relative flow around an incipient disturbance has been hypothesized to be an important factor in protecting the disturbance from environmental intrusions, and thus creating or maintaining a favorable environment for TC genesis to occur (Dunkerton et al. 2009). Brammer and Thorncroft (2015) have shown that, as AEWs leave West Africa, the troughs are sensitive to the low-level environment to their west and northwest. Although the vortex at 700 hPa typically has a closed circulation in the wave-relative reference frame, the AEW troughs are still cold-core in the lower troposphere and, therefore, there is relative westerly flow under the vortex and through the lower levels of the trough. In a composite analysis, significant differences in the moisture of the low-level environment to the northwest of the troughs were found between developing and non-developing waves. Favorable developing waves had significantly higher moisture content in the lower troposphere to the northwest of the trough as they exited the West African coast compared to favorable non-developing waves. Trajectory analysis for all the waves revealed that as the AEWs transition over the West African coast the troughs are typically open to the environment ahead and to the northwest of the trough. For developing waves this means that moist air (e.g., moist tropical sounding, Dunion 2011) is ingested into the lower levels of the system, while for non-developing waves dry air (e.g., the Saharan Air Layer (SAL) or mid-latitude dry air intrusion soundings) is ingested. At this stage in the AEW life cycle, moisture differences may be fundamental in determining whether a favorable wave will develop or not.

The depth and the integrity of the closed circulation around the pre-genesis disturbance is an important consideration for providing a convectively favorable environment for TC genesis. Freismuth et al. (2016) argue that the vortex of ex-Gaston (2010) was susceptible to dry air above the vortex maxima, which hindered deep convection and led to a weakening of the vortex. In addition, non-developing disturbance AL90 (2014) encountered lower tropospheric dry air to its west and northwest, which was ingested by the disturbance and was likely a major contributor in the failed genesis. Brammer and Thorncroft (2017) showed that as AEWs leave the West African coast there is an increase in air parcel trajectories reaching the AEW circulation from low-levels to northwest of the trough circulation. Although the AEW troughs typically possess closed circulations at 700–600 hPa as they leave the West African coast, the circulation can remain open to the environment both above and below the 700–600-hPa layer. As AEWs propagate across the North Atlantic, the troughs are more likely to exhibit closed circulations at low-levels due to either increased vorticity within the trough or the changing background shear profile over the central

GENESIS STAGE EXPERIMENT
Science Description

Atlantic. It was therefore hypothesized that AEWs are especially sensitive to the low-level environment to the west and northwest of the trough during the first three days after leaving the West African coast. Since AEWs typically propagate at 7.5 m s^{-1} over the Atlantic (Kiladis et al. 2006), these waves are typically located near 35°W after three days.

Goal(s): To investigate the favorability in both dynamics (e.g., vertical wind shear) and thermodynamics (e.g., moisture, relative humidity) for tropical cyclogenesis in the environment near a pre-tropical depression, especially the downstream environment.

Hypotheses:

1. Environmental air downstream from a pre-TC (or AEW trough axis) is ingested before the low-level circulation is closed.
2. High environmental relative humidity to the West or Northwest and low environmental vertical wind shear in the vicinity of a pre-TC are critical for a disturbance to develop into a tropical cyclone.
3. Dry air, often associated with but not limited to the Saharan Air Layer (SAL), inhibits or delays genesis of pre-TCs.

Objectives:

1. Collect aircraft observations of thermodynamics and dynamics in the air mass surrounding and ahead of a pre-TC.
2. Consecutive missions are recommended to observe the evolution of the observations over time and how that pertains to the (non-)development of a pre-TC.

Aircraft Pattern/Module Descriptions (see *Flight Pattern* document for more detailed information):

G-IV Pattern #1: Sample the environment to the west of a pre-tropical cyclone (pre-TC), especially if dry air is detected in that region. Sample when the pre-TC is forecast to develop in reliable computer models or is showing signs of development in observations. Standard Lawnmower pattern should be used to set up a grid of observations and dropsondes, with drops every 150 n mi (280 km). The most likely orientation of the lawnmower pattern will be to the West or Northwest of the pre-TC. To maximize the usefulness of the data, a minimum of two lawnmower legs should be flown. In some situations, the same box could be flown twice to maximize data coverage in a more specific region.

P-3 Pattern #1: G-IV Pattern #1 (described above) can be modified to accommodate the P-3. Flight level should be 20-25 kft to maximize the altitude of dropsonde data. P-3 missions will likely start later than G-IV missions due to the greater range of the G-IV, especially if the disturbance is in the Atlantic Main Development Region.

GENESIS STAGE EXPERIMENT

Science Description

Links to Other Genesis Stage Experiments/Modules: This experiment is ideally suited to include sampling of the disturbance itself, including the vorticity maximum and precipitation properties identified from the TDR, and thus the PREFORM experiment (also part of the Genesis Stage). Of course, special consideration must be given to the length of the flight and the distance of the disturbance from the takeoff/landing airport(s). It may be especially fruitful to evaluate the relative humidity in the environment ahead of a disturbance and consequently investigate the precipitation properties within the disturbance itself. The FAM may be modified to support the following coincident experiments/field campaigns. The scientific descriptions of these collaborative efforts are detailed in this document insofar as they pertain to the FAM experiment and its potential modifications. The full scientific documentation for these collaborative efforts is available upon request.

- Impact of Targeted Observations on Forecasts (ITOFS)
- ONR's Moisture and Aerosol Gradients / Physics of Inversion Evolution (MAGPIE)
- Organized Convection and Earthcare Studies over the Tropical Atlantic (ORCESTRA) international field campaign

GENESIS STAGE EXPERIMENT
Science Description

Links to ITOFS

The overarching objective of ITOFS is to fly the NOAA G-IV to collect rare measurements of tropical disturbances and tropical cyclones in their very early stages of development, farther east than any NOAA hurricane reconnaissance aircraft has ever flown. Operating from Cabo Verde, G-IV research missions will provide an opportunity to sample an African easterly wave (AEW) as it emerges from the West African coast. AEWs (as a pre-cursor to a TC) accounts for about 60% of the tropical storms and non-major hurricanes and 85% of the major hurricanes in the Atlantic and is therefore a high priority target for ITOFS. The three goals of ITOFS are:

1. Examining the operational impact of in-situ observations in the normally data sparse eastern and central North Atlantic through real-time monitoring of AEWs and the early stages of TCs, and assessing how those observations may improve model and National Hurricane Center (NHC) short- and long-term forecasts of TC formation;
2. Understanding the structural aspects of AEWs, such as the SAL and African easterly jet (AEJ), that make them favorable or unfavorable for possible development in the western North Atlantic;
3. Evaluating the accuracy of experimental (e.g., NASA TROPICS) and operational (e.g., from GOES, NOAA-20) satellite product guidance in the region against collected G-IV observations (collaboration with the 2024 HFP Experiments: *Evaluation of Tropical Cyclone Environment Using Satellite Soundings Experiment, TROPICS Satellite Validation Module* and *Synthetic Aperture Radar Wind Inspection with NOAA-P3 Data (SARWIND)*).

In addition to the research opportunities ITOFS offers, NOAA NHC is interested in:

- Accessing GPS dropsonde and tail Doppler radar observations in the central and eastern North Atlantic to capture the early lifecycle stages of tropical disturbances and tropical cyclones, especially those that pose a threat to land areas in the central and western part of the Atlantic basin, including the CONUS, Puerto Rico, and the U.S. Virgin Islands.
- Testing strategies for sampling tropical disturbances and tropical cyclones farther east and earlier in their lifecycles. These efforts will help address strategies for future requirements and capabilities, including NOAA NHC 7-day tropical cyclone forecasts and the enhanced range capabilities of the anticipated G550 aircraft.

GENESIS STAGE EXPERIMENT
Science Description

Links to ONR MAGPIE

FAM could also be flown in coordination with MAGPIE, particularly when that team is sampling the marine boundary layer in the vicinity of Barbados (ideally between 15 June - 15 July). In partnership with MAGPIE, we propose to investigate the marine boundary layer structure, evolution and its interaction with the mesoscale thermodynamic and kinematic environments associated with pre-genesis disturbances, differentiating between developing and non-developing systems. To accomplish this, we will target the structure of AEWs, including moist tropical environments and embedded mesoscale convective systems (MCSs), the environments with which they are interacting with such as, e.g., SAL outbreaks (Dunion et al. 2004; Hopsch et al. 2010; Brammer and Thorncroft 2015), and the boundaries between AEWs and the SAL.

Emphasis will be placed on TC genesis in the Atlantic main development region (Goldenberg and Shapiro 1996) and the physical processes controlling genesis:

- Marine boundary layer (MBL) processes (e.g., air-sea interaction and turbulent mixing) and evolution in environments of AEWs, the SAL, and AEW-SAL boundaries
- Static stability, saturation fraction, and relationships with cloud structure and evolution in AEWs, the SAL, and AEW-SAL boundaries

Many of the efforts described above are similar to the objectives outlined in FAM. In addition to conducting these process studies, we will also be testing new technologies for sampling the MBL in these unique environments, such as Altius 600, BlackSwift, and Dragoon sUASs.

Three MAGPIE modules have been identified that could be executed within or in place of the default FAM flight patterns to maximize collaborations. Details of these modules and how they fit into the FAM Experiment are described in the *Flight Patterns* document and include:

1. MAGPIE “Large-Scale” P-3 Module
2. MAGPIE “Convective-Scale” P-3 Module
3. MAGPIE “Island Wake” P-3 Module

Analysis Strategy: Dropsonde profiles will be evaluated to determine the horizontal gradients and advection of environmental relative humidity. In addition, special attention may be paid to isolated convective events to better understand their maintenance and lifecycle. Characteristics of the dry air mass will be scrutinized, including the minimum relative humidity, the height/depth of the dry air, and the horizontal extent of the dry air. Wind analyses from dropsondes and TDR will be evaluated to determine the impact of environmental vertical wind shear on the pre-genesis disturbance. This analysis will go beyond the traditional deep layer vertical wind shear metric, taking into account the hodograph to evaluate vertical wind shear through a number of different levels. The observations collected in this experiment will be crucial to evaluation of dynamics/thermodynamics and the diagnosis of genesis false alarms in numerical weather prediction models (e.g., GFS, HAFS, HWRF). In particular, moisture biases in numerical weather prediction models have long been hypothesized to play a crucial role in false alarm genesis forecasts and, therefore, will be evaluated using observations collected by this module. Data will be made available through in real-time transmission of GPS dropsondes to the GTS.

GENESIS STAGE EXPERIMENT
Science Description

Links to ORCESTRA

The ORCESTRA module is envisioned as a component of the *Favorable Air Mass (FAM) Genesis Stage* experiment, though it could also be flown in conjunction with *PREFORM*. ORCESTRA Pattern #1 is flown in coordination with the location of the German RV Meteor as part of ORCESTRA/BOWTIE and ORCESTRA/PICCOLO. If the RV Meteor is not in a location conducive to coordination, ORCESTRA Pattern #1 could also be flown in coordination with ORCESTRA/PERCUSION HALO G550 flights out of Barbados. ORCESTRA Pattern #2 is ideally flown in coordination with ORCESTRA/PERCUSION HALO G550 flights out of Barbados. Find more information about these flight patterns in the brief descriptions below and in the more detailed descriptions in the *Flight Pattern* document.

Analysis Strategy: Data collected by the ORCESTRA module of the NOAA/AOML HRD Hurricane Field Program will be combined with coincident measurements taken on RV Meteor, the German HALO G550 aircraft, and at the Barbados Cloud Observatory. We will diagnose profiles of mesoscale divergence and vertical mass flux (Bony and Stevens 2019; George et al. 2021) from dropsondes densely deployed by the P-3/G-IV in circles centered on the RV Meteor, which will complement similar coordinated sampling between HALO and the RV Meteor. We will also combine TDR retrievals from the P-3/G-IV with SEA-POL retrievals from the RV Meteor, to characterize the 3D precipitation and wind field.

The RV Meteor will be equipped with the SEA-POL radar, a vertically pointing W-band cloud radar, a Raman (water vapor) lidar, wind lidar, a ceilometer, microwave radiometers, ocean gliders, as well as measurements of aerosol composition, water-stable isotopes, eddy covariance, and skin-temperature. There will also be standard surface meteorological instrumentation, including in-situ measurements of temperature, relative humidity, and downwelling long- and short-wave broadband irradiance. Radiosonde launches are planned for every 3 hours, though we also have the ability to launch additional radiosondes in coordination with NOAA aircraft missions. The Raman lidar is *not* eye safe within the troposphere but there is a safety radar that shuts a latch if objects are detected above.

Goal(s): To characterize the mesoscale thermodynamic environment and circulation and sample precipitation features around the RV Meteor or in coordination with the German G550 HALO in a convective environment within/near a tropical disturbance and/or the Northern edge of the ITCZ.

Hypotheses: The hypotheses of ORCESTRA/PICCOLO are:

- 1) Increasing stratiform area fraction and column moisture lead to greater volumetric rainfall in association with convective upscale development.

GENESIS STAGE EXPERIMENT
Science Description

- 2) Cloud-radiative feedbacks are essential to driving mesoscale organization under conditions of weak vertical wind shear.
- 3) The time scale of cloud-circulation coupling increases with convective upscale growth, resonating with the diurnal cycle of radiative heating in highly organized states.
- 4) Mesoscale organization reduces the rate of work performed by moist convection, in spite of reduced production of entropy by hydrometeor sedimentation.

Objectives:

- 1) Collect aircraft observations of thermodynamics and dynamics in the air mass within 80 n mi (150 km) of the RV Meteor
- 2) Collect airborne TDR observations of convection within a disturbance coincident with targeted SEA-POL radar observations from RV Meteor
- 3) Collect aircraft observations of thermodynamics and dynamics within convection along the edge of the ITCZ.

P-3/G-IV Pattern #1: This pattern uses a Hybrid Figure 4/Circumnavigation centered on the RV Meteor to sample precipitation features, measure the mesoscale thermodynamic environment and circulation with dropsondes, and permit combined TDR and SEA-POL radar (on board RV Meteor) analysis. This pattern complements the ORCESTRA/PICCOLO goals to understand the mechanisms of mesoscale convective organization and characterize the entropy budget. This is also known as the *ORCESTRA “Deep Convection” Module*.

P-3/G-IV Pattern #2: This pattern takes place during the ferry to the disturbance of interest (or the ferry to RV Meteor in conjunction with P-3 Pattern #1). Time permitting, the ferry includes sampling along the Northern edge of the ITCZ, defined by the 50 mm precipitable water contour. This characterizes the precipitation features and thermodynamic and kinematic environment along the Northern edge of the ITCZ. This pattern complements the ORCESTRA/PERCUSSION and ORCESTRA/BOW-TIE goals to understand how mesoscale processes impact the structure of the Atlantic ITCZ. This pattern is also known as the *ORCESTRA “Precipitable Water” Module*.

GENESIS STAGE EXPERIMENT

Science Description

References:

- Bony, S., and B. Stevens, 2019: Measuring Area-Averaged Vertical Motions with Dropsondes. *J. Atmos. Sci.*, **76**, 767–783, <https://doi.org/10.1175/JAS-D-18-0141.1>.
- Brammer A., and C. D. Thorncroft, 2015: Variability and Evolution of African Easterly Wave Structures and Their Relationship with Tropical Cyclogenesis over the Eastern Atlantic. *Mon. Wea. Rev.*, **143**, 4975–4995.
- Brammer A., and C. D. Thorncroft, 2017: Spatial and Temporal Variability of the Three-Dimensional Flow around African Easterly Waves. *Mon. Wea. Rev.*, **145**, 2879–2895.
- Dunion, J.P., 2011: Re-writing the climatology of the tropical North Atlantic and Caribbean Sea atmosphere. *J. Climate*, **24**, 893–908.
- Dunion, J. P., and C. S. Velden, 2004: The Impact of the Saharan Air Layer on Atlantic Tropical Cyclone Activity. *Bull. Amer. Meteor. Soc.*, **85**, 353-366, <https://doi.org/10.1175/bams-85-3-353>.
- Dunkerton, T. J., M. T. Montgomery, and Z. Wang, 2009: Tropical cyclogenesis in a tropical wave critical layer: Easterly waves. *Atmos. Chem. Phys.*, **9**, 5587–5646.
- Dunn, G. E., 1940: Cyclogenesis in the tropical Atlantic. *Bull. Amer. Meteor. Soc.*, **21**, 215–229.
- Freismuth, T. M., Rutherford, B., Boothe, M. A., and Montgomery, M. T., 2016: Why did the storm ex-Gaston (2010) fail to redevelop during the PREDICT experiment? *Atmos. Chem. Phys.*, **16**, 8511–8519.
- George, G., and Coauthors, 2021: JOANNE: Joint dropsonde Observations of the Atmosphere in tropical North atlantic meso-scale Environments. *Earth Syst. Sci. Data*, **13**, 5253-5272, <https://doi.org/10.5194/essd-13-5253-2021>.
- Goldenberg, S. B., and L. J. Shapiro, 1996: Physical Mechanisms for the Association of El Niño and West African Rainfall with Atlantic Major Hurricane Activity. *J. Climate*, **9**, 1169-1187, [https://doi.org/10.1175/1520-0442\(1996\)009<1169:Pmftao>2.0.Co;2](https://doi.org/10.1175/1520-0442(1996)009<1169:Pmftao>2.0.Co;2).
- Halperin, D. J., H. E. Fuelberg, R. E. Hart, J. H. Cossuth, P. Sura, and R. J. Pasch, 2013: An evaluation of tropical cyclone genesis forecasts from global numerical models. *Wea. Forecasting*, **28**, 1423–1445, doi:10.1175/WAF-D-13-00008.1
- Hopsch, S. B., C. D. Thorncroft, and K. R. Tyle, 2010: Analysis of African Easterly Wave Structures and Their Role in Influencing Tropical Cyclogenesis. *Mon. Wea. Rev.*, **138**, 1399-1419, <https://doi.org/10.1175/2009mwr2760.1>.
- Landsea, C. W., 1993: A climatology of intense (or major) Atlantic hurricanes. *Mon. Wea. Rev.*, **121**, 1703–1713.
- Riehl, H., 1945: "Waves in the easterlies and the polar front in the tropics" Misc. Rep. No. 17, Department of Meteorology, University of Chicago, 79 pp.